American College of Radiology ACR Appropriateness Criteria® Workup of Noncerebral Systemic Arterial Embolic Source

<u>Variant: 1</u> Known upper extremity arterial occlusion. Suspected embolic etiology. Next imaging study to determine source.

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	0
US echocardiography transthoracic resting	Usually Appropriate	0
MRA chest without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without IV contrast	Usually Appropriate	0
CTA chest with IV contrast	Usually Appropriate	���
CT heart function and morphology with IV contrast	Usually Appropriate	૽ ૽
MRA chest without IV contrast	May Be Appropriate	0
US duplex Doppler abdomen	Usually Not Appropriate	0
Aortography chest	Usually Not Appropriate	∵ •••

<u>Variant: 2</u> Known arterial occlusion in the mesenteric or renal arterial system or renal infarcts. Suspected embolic etiology. Next imaging study to determine source.

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	0
US echocardiography transthoracic resting	Usually Appropriate	0
MRA chest and abdomen without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without IV contrast	Usually Appropriate	0
CTA chest with IV contrast	Usually Appropriate	���
CT heart function and morphology with IV contrast	Usually Appropriate	∵
CTA chest and abdomen with IV contrast	Usually Appropriate	∵ ∵ ∵ ∵
US duplex Doppler abdomen	May Be Appropriate	0
MRA chest and abdomen without IV contrast	May Be Appropriate	0
MRA chest without and with IV contrast	May Be Appropriate	0
MRA chest without IV contrast	May Be Appropriate	0
Aortography chest and abdomen	Usually Not Appropriate	※※※

<u>Variant: 3</u> Known lower extremity arterial occlusion. Suspected embolic etiology. Next imaging study to determine source.

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	0
US echocardiography transthoracic resting	Usually Appropriate	0
MRA chest abdomen pelvis without and with IV contrast	Usually Appropriate	0
MRA chest without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without IV contrast	Usually Appropriate	0

CTA chest with IV contrast	Usually Appropriate	⊗ ⊗
CT heart function and morphology with IV contrast	Usually Appropriate	
CTA chest abdomen pelvis with IV contrast	Usually Appropriate	
MRA chest abdomen pelvis without IV contrast	May Be Appropriate	0
MRA chest without IV contrast	May Be Appropriate	0
US duplex Doppler abdomen	Usually Not Appropriate	0
Aortography chest abdomen pelvis	Usually Not Appropriate	∵ ∵ ∵

<u>Variant: 4</u> Known multiorgan system arterial occlusions. Suspected embolic etiology. Next imaging study to determine source.

Procedure	Appropriateness Category	Relative Radiation Level
US echocardiography transesophageal	Usually Appropriate	0
US echocardiography transthoracic resting	Usually Appropriate	0
MRA chest abdomen pelvis without and with IV contrast	Usually Appropriate	0
MRA chest abdomen pelvis without IV contrast	Usually Appropriate	0
MRA chest without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without and with IV contrast	Usually Appropriate	0
MRI heart function and morphology without IV contrast	Usually Appropriate	0
CTA chest with IV contrast	Usually Appropriate	૽ ૽
CT heart function and morphology with IV contrast	Usually Appropriate	⊗⊗⊗
CTA chest abdomen pelvis with IV contrast	Usually Appropriate	⊗⊗⊗⊗
US duplex Doppler abdomen	May Be Appropriate	0
MRA chest without IV contrast	May Be Appropriate	0

Panel Members

Summary of Literature Review

Introduction/Background

Special Imaging Considerations

Initial Imaging Definition

Discussion of Procedures by Variant

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

A. Aortography Chest

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

B. CT Heart Function and Morphology With IV Contrast

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

C. CTA Chest With IV Contrast

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

D. MRA Chest Without and With IV Contrast

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

E. MRA Chest Without IV Contrast

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

F. MRI Heart Function and Morphology Without and With IV Contrast

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

G. MRI Heart Function and Morphology Without IV Contrast

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

H. US Duplex Doppler Abdomen

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

I. US Echocardiography Transesophageal

Variant 1: The variant assumes that an upper extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

J. US Echocardiography Transthoracic Resting

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR

Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]). A. Aortography Chest and Abdomen

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

B. CT Heart Function and Morphology With IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

C. CTA Chest With IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

D. CTA Chest and Abdomen With IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

E. MRA Chest Without and With IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

F. MRA Chest Without IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

G. MRA Chest and Abdomen Without and With IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

H. MRA Chest and Abdomen Without IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

I. MRI Heart Function and Morphology Without and With IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA,

although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

J. MRI Heart Function and Morphology Without IV Contrast

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

K. US Duplex Doppler Abdomen

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

L. US Echocardiography Transesophageal

Variant 2: The variant assumes that a mesenteric/renal arterial occlusion or renal infarct has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Imaging of Mesenteric Ischemia" [55]).

M. US Echocardiography Transthoracic Resting

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

A. Aortography Chest, Abdomen, and Pelvis

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

B. CT Heart Function and Morphology With IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

C. CTA Chest With IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

D. CTA Chest, Abdomen, and Pelvis With IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the

clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]). E. MRA Chest Without and With IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

F. MRA Chest Without IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

G. MRA Chest, Abdomen, and Pelvis Without and With IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

H. MRA Chest, Abdomen, and Pelvis Without IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

I. MRI Heart Function and Morphology Without and With IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

J. MRI Heart Function and Morphology Without IV Contrast

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

K. US Duplex Doppler Abdomen

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

L. US Echocardiography Transesophageal

Variant 3: The variant assumes that a lower extremity arterial occlusion has already been established. Typically, this diagnosis is made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used (see the ACR Appropriateness Criteria® topic on "Sudden Onset of Cold, Painful Leg" [59]).

M. US Echocardiography Transthoracic Resting

Variant 4: The variant assumes that multiorgan arterial occlusions have already been

established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

A. CT Heart Function and Morphology With IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

B. CTA Chest With IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

C. CTA Chest, Abdomen, and Pelvis With IV contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

D. MRA Chest Without and With IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

E. MRA Chest Without IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

F. MRA Chest, Abdomen, and Pelvis Without and With IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

G. MRA Chest, Abdomen, and Pelvis Without IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

H. MRI Heart Function and Morphology Without and With IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

I. MRI Heart Function and Morphology Without IV Contrast

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

J. US Duplex Doppler Abdomen

Variant 4: The variant assumes that multiorgan arterial occlusions have already been

established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used. K. US Echocardiography Transesophageal

Variant 4: The variant assumes that multiorgan arterial occlusions have already been established. Typically, these diagnoses are made by CTA, arteriography, or MRA, although the clinical examination or another imaging study could also be used.

L. US Echocardiography Transthoracic Resting

Summary of Recommendations

Supporting Documents

The evidence table, literature search, and appendix for this topic are available at https://acsearch.acr.org/list. The appendix includes the strength of evidence assessment and the final rating round tabulations for each recommendation.

For additional information on the Appropriateness Criteria methodology and other supporting documents, please go to the ACR website at https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Appropriateness-Criteria.

Appropriateness Category Names and Definitions

Appropriateness Category Name	Appropriateness Rating	Appropriateness Category Definition
Usually Appropriate	7, 8, or 9	The imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable riskbenefit ratio for patients.
May Be Appropriate	4, 5, or 6	The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal.
May Be Appropriate (Disagreement)	5	The individual ratings are too dispersed from the panel median. The different label provides transparency regarding the panel's recommendation. "May be appropriate" is the rating category and a rating of 5 is assigned.
Usually Not Appropriate	1, 2, or 3	The imaging procedure or treatment is unlikely to be

	indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

Relative Radiation Level Information

References

- **1.** Weiss S, Buhlmann R, von Allmen RS, et al. Management of floating thrombus in the aortic arch. Journal of Thoracic & Cardiovascular Surgery. 152(3):810-7, 2016 09.
- **2.** Kumagai T, Matsuura Y, Yamamoto T, Ugawa Y, Fukushima T. Risk factors for left atrial thrombus from transesophageal echocardiography findings in ischemic stroke patients. Fukushima J Med Sci. 60(2):154-8, 2014.
- **3.** Wysokinski WE, Ammash N, Sobande F, Kalsi H, Hodge D, McBane RD. Predicting left atrial thrombi in atrial fibrillation. Am Heart J. 159(4):665-71, 2010 Apr.
- **4.** Yamamoto M, Seo Y, Kawamatsu N, et al. Complex left atrial appendage morphology and left atrial appendage thrombus formation in patients with atrial fibrillation. Circ Cardiovasc Imaging. 7(2):337-43, 2014 Mar.
- **5.** Gianstefani S, Douiri A, Delithanasis I, et al. Incidence and predictors of early left ventricular thrombus after ST-elevation myocardial infarction in the contemporary era of primary percutaneous coronary intervention. Am J Cardiol 2014;113:1111-6.
- **6.** McCarthy CP, Vaduganathan M, McCarthy KJ, Januzzi JL, Jr., Bhatt DL, McEvoy JW. Left Ventricular Thrombus After Acute Myocardial Infarction: Screening, Prevention, and Treatment. JAMA Cardiol 2018;3:642-49.
- **7.** Sordelli C, Fele N, Mocerino R, et al. Infective Endocarditis: Echocardiographic Imaging and New Imaging Modalities. J Cardiovasc Echogr 2019;29:149-55.
- **8.** Wintersperger BJ, Becker CR, Gulbins H, et al. Tumors of the cardiac valves: imaging findings in magnetic resonance imaging, electron beam computed tomography, and echocardiography. Eur Radiol 2000;10:443-9.
- **9.** O'Connell JB, Quinones-Baldrich WJ. Proper evaluation and management of acute embolic versus thrombotic limb ischemia. Semin Vasc Surg 2009;22:10-6.
- **10.** Yoo SM, Lee HY, White CS. MDCT evaluation of acute aortic syndrome. [Review] [62 refs]. Radiol Clin North Am. 48(1):67-83, 2010 Jan.
- **11.** Tsilimparis N, Hanack U, Pisimisis G, Yousefi S, Wintzer C, Ruckert RI. Thrombus in the non-aneurysmal, non-atherosclerotic descending thoracic aorta--an unusual source of arterial embolism. Eur J Vasc Endovasc Surg. 41(4):450-7, 2011 Apr.
- **12.** Klang E, Kerpel A, Soffer S, et al. CT imaging features of symptomatic and asymptomatic floating aortic thrombus. Clin Radiol. 73(3):323.e9-323.e14, 2018 03.
- **13.** Pagni S, Trivedi J, Ganzel BL, et al. Thoracic aortic mobile thrombus: is there a role for early surgical intervention?. Ann Thorac Surg. 91(6):1875-81, 2011 Jun.
- 14. Boufi M, Mameli A, Compes P, Hartung O, Alimi YS. Elective stent-graft treatment for the

- management of thoracic aorta mural thrombus. European Journal of Vascular & Endovascular Surgery. 47(4):335-41, 2014 Apr.
- **15.** American College of Radiology. ACR–NASCI–SIR–SPR Practice Parameter for the Performance and Interpretation of Body Computed Tomography Angiography (CTA). Available at: https://gravitas.acr.org/PPTS/GetDocumentView?docId=164+&releaseId=2.
- **16.** Fleischmann D, Mitchell RS, Miller DC. Acute aortic syndromes: new insights from electrocardiographically gated computed tomography. [Review] [8 refs]. Semin Thorac Cardiovasc Surg. 20(4):340-7, 2008.
- **17.** Holloway BJ, Rosewarne D, Jones RG. Imaging of thoracic aortic disease. [Review]. Br J Radiol. 84 Spec No 3:S338-54, 2011 Dec.
- **18.** Hur J, Kim YJ, Lee HJ, et al. Left atrial appendage thrombi in stroke patients: detection with two-phase cardiac CT angiography versus transesophageal echocardiography. Radiology. 251(3):683-90, 2009 Jun.
- **19.** Hur J, Kim YJ, Lee HJ, et al. Dual-enhanced cardiac CT for detection of left atrial appendage thrombus in patients with stroke: a prospective comparison study with transesophageal echocardiography. Stroke. 42(9):2471-7, 2011 Sep.
- **20.** Hur J, Kim YJ, Lee HJ, et al. Cardioembolic stroke: dual-energy cardiac CT for differentiation of left atrial appendage thrombus and circulatory stasis. Radiology. 263(3):688-95, 2012 Jun.
- **21.** Teunissen C, Habets J, Velthuis BK, Cramer MJ, Loh P. Double-contrast, single-phase computed tomography angiography for ruling out left atrial appendage thrombus prior to atrial fibrillation ablation. Int J Cardiovasc Imaging. 33(1):121-128, 2017 Jan.
- **22.** Kapa S, Martinez MW, Williamson EE, et al. ECG-gated dual-source CT for detection of left atrial appendage thrombus in patients undergoing catheter ablation for atrial fibrillation. J Interv Card Electrophysiol. 29(2):75-81, 2010 Nov.
- **23.** Zhai Z, Tang M, Zhang S, et al. Transoesophageal echocardiography prior to catheter ablation could be avoided in atrial fibrillation patients with a low risk of stroke and without filling defects in the late-phase MDCT scan: A retrospective analysis of 783 patients. Eur Radiol. 28(5):1835-1843, 2018 May.
- **24.** Ikegami Y, Tanimoto K, Inagawa K, et al. Identification of Left Atrial Appendage Thrombi in Patients With Persistent and Long-Standing Persistent Atrial Fibrillation Using Intra-Cardiac Echocardiography and Cardiac Computed Tomography. Circ J. 82(1):46-52, 2017 12 25.
- **25.** Kantarci M, Ogul H, Sade R, Aksakal E, Colak A, Tanboga IH. Circulatory Stasis or Thrombus in Left Atrial Appendage, An Easy Diagnostic Solution. J Comput Assist Tomogr. 43(3):406-409, 2019 May/Jun.
- **26.** Bilchick KC, Mealor A, Gonzalez J, et al. Effectiveness of integrating delayed computed tomography angiography imaging for left atrial appendage thrombus exclusion into the care of patients undergoing ablation of atrial fibrillation. Heart Rhythm. 13(1):12-9, 2016 Jan.
- **27.** Martinez MW, Kirsch J, Williamson EE, et al. Utility of nongated multidetector computed tomography for detection of left atrial thrombus in patients undergoing catheter ablation of atrial fibrillation. JACC Cardiovasc Imaging 2009;2:69-76.
- **28.** Romero J, Husain SA, Kelesidis I, Sanz J, Medina HM, Garcia MJ. Detection of left atrial appendage thrombus by cardiac computed tomography in patients with atrial fibrillation: a

- meta-analysis. [Review]. Circ Cardiovasc Imaging. 6(2):185-94, 2013 Mar 01.
- **29.** Vira T, Pechlivanoglou P, Connelly K, Wijeysundera HC, Roifman I. Cardiac computed tomography and magnetic resonance imaging vs. transoesophageal echocardiography for diagnosing left atrial appendage thrombi. Europace. 21(1):e1-e10, 2019 Jan 01.
- **30.** Zou H, Zhang Y, Tong J, Liu Z. Multidetector computed tomography for detecting left atrial/left atrial appendage thrombus: a meta-analysis. [Review]. Intern Med J. 45(10):1044-53, 2015 Oct.
- **31.** Choi BH, Ko SM, Hwang HK, et al. Detection of left atrial thrombus in patients with mitral stenosis and atrial fibrillation: retrospective comparison of two-phase computed tomography, transoesophageal echocardiography and surgical findings. Eur Radiol. 23(11):2944-53, 2013 Nov.
- **32.** Korhonen M, Muuronen A, Arponen O, et al. Left atrial appendage morphology in patients with suspected cardiogenic stroke without known atrial fibrillation. PLoS ONE. 10(3):e0118822, 2015.
- **33.** Hozawa M, Morino Y, Matsumoto Y, et al. 3D-computed tomography to compare the dimensions of the left atrial appendage in patients with normal sinus rhythm and those with paroxysmal atrial fibrillation. Heart Vessels. 33(7):777-785, 2018 Jul.
- **34.** Dieker W, Behnes M, Fastner C, et al. Impact of left atrial appendage morphology on thrombus formation after successful left atrial appendage occlusion: Assessment with cardiac-computed-tomography. Sci. rep.. 8(1):1670, 2018 01 26.
- **35.** Bittencourt MS, Achenbach S, Marwan M, et al. Left ventricular thrombus attenuation characterization in cardiac computed tomography angiography. J Cardiovasc Comput Tomogr 2012;6:121-6.
- **36.** Kim IC, Chang S, Hong GR, et al. Comparison of Cardiac Computed Tomography With Transesophageal Echocardiography for Identifying Vegetation and Intracardiac Complications in Patients With Infective Endocarditis in the Era of 3-Dimensional Images. Circ Cardiovasc Imaging. 11(3):e006986, 2018 03.
- **37.** Erba PA, Pizzi MN, Roque A, et al. Multimodality Imaging in Infective Endocarditis: An Imaging Team Within the Endocarditis Team. Circulation 2019;140:1753-65.
- **38.** Feuchtner GM, Stolzmann P, Dichtl W, et al. Multislice computed tomography in infective endocarditis: comparison with transesophageal echocardiography and intraoperative findings. J Am Coll Cardiol. 2009;53(5):436-444.
- **39.** Araoz PA, Mulvagh SL, Tazelaar HD, Julsrud PR, Breen JF. CT and MR imaging of benign primary cardiac neoplasms with echocardiographic correlation. Radiographics. 2000;20(5):1303-19.
- **40.** Kassop D, Donovan MS, Cheezum MK, et al. Cardiac Masses on Cardiac CT: A Review. Curr Cardiovasc Imaging Rep 2014;7:9281.
- **41.** Mesurolle B, Qanadli SD, Merad M, El Hajjam M, Mignon F, Lacombe P. Dual-slice helical CT of the thoracic aorta. J Comput Assist Tomogr. 24(4):548-56, 2000 Jul-Aug.
- **42.** Ryoo S, Chung JW, Lee MJ, et al. An Approach to Working Up Cases of Embolic Stroke of Undetermined Source. Journal of the American Heart Association. 5(3):e002975, 2016 Mar 22.

- **43.** Krishnam MS, Tomasian A, Malik S, Desphande V, Laub G, Ruehm SG. Image quality and diagnostic accuracy of unenhanced SSFP MR angiography compared with conventional contrast-enhanced MR angiography for the assessment of thoracic aortic diseases. Eur Radiol. 20(6):1311-20, 2010 Jun.
- **44.** Gebker R, Gomaa O, Schnackenburg B, Rebakowski J, Fleck E, Nagel E. Comparison of different MRI techniques for the assessment of thoracic aortic pathology: 3D contrast enhanced MR angiography, turbo spin echo and balanced steady state free precession. Int J Cardiovasc Imaging. 23(6):747-56, 2007 Dec.
- **45.** Chen J, Zhang H, Zhu D, Wang Y, Byanju S, Liao M. Cardiac MRI for detecting left atrial/left atrial appendage thrombus in patients with atrial fibrillation: Meta-analysis and systematic review. Herz. 44(5):390-397, 2019 Aug.
- **46.** Srichai MB, Junor C, Rodriguez LL, et al. Clinical, imaging, and pathological characteristics of left ventricular thrombus: a comparison of contrast-enhanced magnetic resonance imaging, transthoracic echocardiography, and transesophageal echocardiography with surgical or pathological validation. Am Heart J 2006;152:75-84.
- **47.** Rustemli A, Bhatti TK, Wolff SD. Evaluating cardiac sources of embolic stroke with MRI. Echocardiography 2007;24:301-8; discussion 08.
- **48.** Motwani M, Kidambi A, Herzog BA, Uddin A, Greenwood JP, Plein S. MR imaging of cardiac tumors and masses: a review of methods and clinical applications. [Review]. Radiology. 268(1):26-43, 2013 Jul.
- **49.** Weinsaft JW, Kim J, Medicherla CB, et al. Echocardiographic Algorithm for Post-Myocardial Infarction LV Thrombus: A Gatekeeper for Thrombus Evaluation by Delayed Enhancement CMR. JACC Cardiovasc Imaging. 9(5):505-15, 2016 05.
- **50.** He YQ, Liu L, Zhang MC, Zeng H, Yang P. Dual-Energy Computed Tomography-Enabled Material Separation in Diagnosing Left Atrial Appendage Thrombus. Tex Heart Inst J. 46(2):107-114, 2019 Apr.
- **51.** Kumar V, Nanda NC. Is it time to move on from two-dimensional transesophageal to three-dimensional transthoracic echocardiography for assessment of left atrial appendage? Review of existing literature. [Review]. Echocardiography. 29(1):112-6, 2012.
- **52.** Nakanishi K, Homma S. Role of echocardiography in patients with stroke. [Review]. J Cardiol. 68(2):91-9, 2016 08.
- **53.** de Bruijn SF, Agema WR, Lammers GJ, et al. Transesophageal echocardiography is superior to transthoracic echocardiography in management of patients of any age with transient ischemic attack or stroke. Stroke 2006;37:2531-4.
- **54.** Pearson AC, Labovitz AJ, Tatineni S, Gomez CR. Superiority of transesophageal echocardiography in detecting cardiac source of embolism in patients with cerebral ischemia of uncertain etiology. J Am Coll Cardiol 1991;17:66-72.
- **55.** Ginsburg M, Obara P, Lambert DL, et al. ACR Appropriateness Criteria® Imaging of Mesenteric Ischemia. J Am Coll Radiol 2018;15:S332-S40.
- **56.** Labruto F, Blomqvist L, Swedenborg J. Imaging the intraluminal thrombus of abdominal aortic aneurysms: techniques, findings, and clinical implications. [Review]. J Vasc Interv Radiol. 22(8):1069-75; quiz 1075, 2011 Aug.

- **57.** Litmanovich D, Bankier AA, Cantin L, Raptopoulos V, Boiselle PM. CT and MRI in diseases of the aorta. AJR Am J Roentgenol 2009;193:928-40.
- **58.** Nguyen VL, Leiner T, Hellenthal FA, et al. Abdominal aortic aneurysms with high thrombus signal intensity on magnetic resonance imaging are associated with high growth rate. Eur J Vasc Endovasc Surg. 48(6):676-84, 2014 Dec.
- **59.** Expert Panel on Vascular Imaging:, Weiss CR, Azene EM, et al. ACR Appropriateness Criteria R Sudden Onset of Cold, Painful Leg. [Review]. J. Am. Coll. Radiol.. 14(5S):S307-S313, 2017 May.
- **60.** American College of Radiology. ACR Appropriateness Criteria® Radiation Dose Assessment Introduction. Available at: https://edge.sitecorecloud.io/americancoldf5f-acrorgf92a-productioncb02-3650/media/ACR/Files/Clinical/Appropriateness-Criteria/ACR-Appropriateness-Criteria-Radiation-Dose-Assessment-Introduction.pdf.

Disclaimer

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination