

**American College of Radiology
ACR Appropriateness Criteria®
Nonvariceal Upper Gastrointestinal Bleeding**

Variant: 1 Adult. Suspected nonvariceal upper gastrointestinal bleeding; no endoscopy performed. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
CTA abdomen and pelvis without and with IV contrast	Usually Appropriate	⦿⦿⦿⦿
Arteriography visceral	May Be Appropriate	⦿⦿⦿
Fluoroscopy upper GI series	Usually Not Appropriate	⦿⦿⦿
MR enterography	Usually Not Appropriate	○
CT abdomen and pelvis with IV contrast	Usually Not Appropriate	⦿⦿⦿
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	⦿⦿⦿
CT abdomen with IV contrast	Usually Not Appropriate	⦿⦿⦿
CT abdomen without IV contrast	Usually Not Appropriate	⦿⦿⦿
CTA abdomen with IV contrast	Usually Not Appropriate	⦿⦿⦿
CTA chest with IV contrast	Usually Not Appropriate	⦿⦿⦿
CTA chest without and with IV contrast	Usually Not Appropriate	⦿⦿⦿
RBC scan abdomen and pelvis	Usually Not Appropriate	⦿⦿⦿
CT abdomen and pelvis without and with IV contrast	Usually Not Appropriate	⦿⦿⦿⦿
CT abdomen without and with IV contrast	Usually Not Appropriate	⦿⦿⦿⦿
CT enterography	Usually Not Appropriate	⦿⦿⦿⦿
CTA abdomen and pelvis with IV contrast	Usually Not Appropriate	⦿⦿⦿⦿
CTA abdomen without and with IV contrast	Usually Not Appropriate	⦿⦿⦿⦿

Variant: 2 Adult. Endoscopy confirms nonvariceal upper gastrointestinal bleeding with a clear source, but treatment not possible or continued bleeding after endoscopic treatment. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
Arteriography visceral	Usually Appropriate	⦿⦿⦿
CTA abdomen and pelvis without and with IV contrast	Usually Appropriate	⦿⦿⦿⦿
CTA chest without and with IV contrast	May Be Appropriate	⦿⦿⦿
CTA abdomen without and with IV contrast	May Be Appropriate	⦿⦿⦿⦿
Fluoroscopy upper GI series	Usually Not Appropriate	⦿⦿⦿
MR enterography	Usually Not Appropriate	○
CT abdomen and pelvis with IV contrast	Usually Not Appropriate	⦿⦿⦿
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	⦿⦿⦿
CT abdomen with IV contrast	Usually Not Appropriate	⦿⦿⦿
CT abdomen without IV contrast	Usually Not Appropriate	⦿⦿⦿
CTA abdomen with IV contrast	Usually Not Appropriate	⦿⦿⦿
CTA chest with IV contrast	Usually Not Appropriate	⦿⦿⦿
RBC scan abdomen and pelvis	Usually Not Appropriate	⦿⦿⦿
CT abdomen and pelvis without and with IV contrast	Usually Not Appropriate	⦿⦿⦿⦿

CT abdomen without and with IV contrast	Usually Not Appropriate	☢☢☢☢
CT enterography	Usually Not Appropriate	☢☢☢☢
CTA abdomen and pelvis with IV contrast	Usually Not Appropriate	☢☢☢☢

Variant: 3 Adult. Endoscopy confirms nonvariceal upper gastrointestinal bleeding without a clear source. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
CTA abdomen and pelvis without and with IV contrast	Usually Appropriate	☢☢☢☢
Arteriography visceral	May Be Appropriate	☢☢☢
CTA chest without and with IV contrast	May Be Appropriate	☢☢☢
RBC scan abdomen and pelvis	May Be Appropriate	☢☢☢
CT enterography	May Be Appropriate	☢☢☢☢☢
Fluoroscopy upper GI series	Usually Not Appropriate	☢☢☢
MR enterography	Usually Not Appropriate	0
CT abdomen and pelvis with IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen with IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen without IV contrast	Usually Not Appropriate	☢☢☢
CTA abdomen with IV contrast	Usually Not Appropriate	☢☢☢
CTA chest with IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen and pelvis without and with IV contrast	Usually Not Appropriate	☢☢☢☢☢
CT abdomen without and with IV contrast	Usually Not Appropriate	☢☢☢☢☢
CTA abdomen and pelvis with IV contrast	Usually Not Appropriate	☢☢☢☢☢
CTA abdomen without and with IV contrast	Usually Not Appropriate	☢☢☢☢☢

Variant: 4 Adult. Nonvariceal upper gastrointestinal bleeding; negative endoscopy. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
CT enterography	Usually Appropriate	☢☢☢☢
CTA abdomen and pelvis without and with IV contrast	Usually Appropriate	☢☢☢☢
MR enterography	May Be Appropriate	0
RBC scan abdomen and pelvis	May Be Appropriate	☢☢☢
CT abdomen and pelvis without and with IV contrast	May Be Appropriate	☢☢☢☢☢
CTA abdomen without and with IV contrast	May Be Appropriate (Disagreement)	☢☢☢☢☢
Arteriography visceral	Usually Not Appropriate	☢☢☢
Fluoroscopy upper GI series	Usually Not Appropriate	☢☢☢
CT abdomen and pelvis with IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen with IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen without IV contrast	Usually Not Appropriate	☢☢☢
CTA abdomen with IV contrast	Usually Not Appropriate	☢☢☢
CTA chest with IV contrast	Usually Not Appropriate	☢☢☢
CTA chest without and with IV contrast	Usually Not Appropriate	☢☢☢
CT abdomen without and with IV contrast	Usually Not Appropriate	☢☢☢☢☢

CTA abdomen and pelvis with IV contrast	Usually Not Appropriate	☢☢☢☢
---	-------------------------	------

Variant: 5 Adult. Postsurgical or traumatic causes of nonvariceal upper gastrointestinal bleeding. Endoscopy is contraindicated. Initial imaging.

Procedure	Appropriateness Category	Relative Radiation Level
Arteriography visceral	Usually Appropriate	☢☢☢
CTA abdomen and pelvis without and with IV contrast	Usually Appropriate	☢☢☢☢☢
CT abdomen and pelvis with IV contrast	May Be Appropriate	☢☢☢☢
CTA chest without and with IV contrast	May Be Appropriate	☢☢☢☢
CT abdomen and pelvis without and with IV contrast	May Be Appropriate	☢☢☢☢☢☢
CT enterography	May Be Appropriate	☢☢☢☢☢☢
Fluoroscopy upper GI series	Usually Not Appropriate	☢☢☢☢
CT abdomen and pelvis without IV contrast	Usually Not Appropriate	☢☢☢☢
CT abdomen with IV contrast	Usually Not Appropriate	☢☢☢☢☢
CT abdomen without IV contrast	Usually Not Appropriate	☢☢☢☢☢
CTA abdomen with IV contrast	Usually Not Appropriate	☢☢☢☢☢
CTA chest with IV contrast	Usually Not Appropriate	☢☢☢☢☢
RBC scan abdomen and pelvis	Usually Not Appropriate	☢☢☢☢☢
CT abdomen without and with IV contrast	Usually Not Appropriate	☢☢☢☢☢☢
CTA abdomen and pelvis with IV contrast	Usually Not Appropriate	☢☢☢☢☢☢
CTA abdomen without and with IV contrast	Usually Not Appropriate	☢☢☢☢☢☢

Panel Members

Summary of Literature Review

Introduction/Background

Special Imaging Considerations

Initial Imaging Definition

Discussion of Procedures by Variant

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

A. Arteriography Visceral

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

B. CT Abdomen and Pelvis With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

C. CT Abdomen and Pelvis Without and With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

D. CT Abdomen and Pelvis Without IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

E. CT Abdomen With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

F. CT Abdomen Without and With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

G. CT Abdomen Without IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

H. CT Enterography

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

I. CTA Abdomen and Pelvis With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

J. CTA Abdomen and Pelvis Without and With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

K. CTA Abdomen With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

L. CTA Abdomen Without and With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

M. CTA Chest With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

N. CTA Chest Without and With IV Contrast

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

O. Fluoroscopy Upper GI Series

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

P. MR Enterography

Variant 1: Endoscopy is the usual first test in patients presenting with overt or occult UGIB [6]. This variant is applicable to a clinical scenario in which a patient presents clinically with overt UGIB and initial endoscopy was not performed due to large volume bleeding or clinical instability.

Q. RBC Scan Abdomen and Pelvis

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive

treatment of the bleeding was not possible or there is continued bleeding after treatment.

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

A. Arteriography Visceral

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

B. CT Abdomen and Pelvis With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

C. CT Abdomen and Pelvis Without and With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

D. CT Abdomen and Pelvis Without IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

E. CT Abdomen With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

F. CT Abdomen Without and With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

G. CT Abdomen Without IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

H. CT Enterography

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

I. CTA Abdomen and Pelvis With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.

J. CTA Abdomen and Pelvis Without and With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.
K. CTA Abdomen With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.
L. CTA Abdomen Without and With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.
M. CTA Chest With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.
N. CTA Chest Without and With IV Contrast

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.
O. Fluoroscopy Upper GI Series

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.
P. MR Enterography

Variant 2: This variant is applicable in a clinical scenario when the patient had endoscopy performed, which diagnosed the upper GI tract as the source of bleed, but definitive treatment of the bleeding was not possible or there is continued bleeding after treatment.
Q. RBC Scan Abdomen and Pelvis

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.
A. Arteriography Visceral

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.
B. CT Abdomen and Pelvis With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these

patients typically present with overt GIB.

C. CT Abdomen and Pelvis Without and With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

D. CT Abdomen and Pelvis Without IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

E. CT Abdomen With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

F. CT Abdomen Without and With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

G. CT Abdomen Without IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

H. CT Enterography

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

I. CTA Abdomen and Pelvis With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

J. CTA Abdomen and Pelvis Without and With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

K. CTA Abdomen With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

L. CTA Abdomen Without and With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

M. CTA Chest With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

N. CTA Chest Without and With IV Contrast

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

O. Fluoroscopy Upper GI Series

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

P. MR Enterography

Variant 3: This variant is applicable to clinical scenario in which endoscopy shows UGIB but the site or source of the bleeding cannot be determined on endoscopy. Clinically, these patients typically present with overt GIB.

Q. RBC Scan Abdomen and Pelvis

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

A. Arteriography Visceral

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

B. CT Abdomen and Pelvis With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

C. CT Abdomen and Pelvis Without and With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite

complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

D. CT Abdomen and Pelvis Without IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

E. CT Abdomen With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

F. CT Abdomen Without and With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

G. CT Abdomen Without IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

H. CT Enterography

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

I. CTA Abdomen and Pelvis With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

J. CTA Abdomen and Pelvis Without and With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which

may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

K. CTA Abdomen With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

L. CTA Abdomen Without and With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

M. CTA Chest With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

N. CTA Chest Without and With IV Contrast

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

O. Fluoroscopy Upper GI Series

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

P. MR Enterography

Variant 4: This variant is applicable to patients with no clear source of bleeding despite complete endoscopic evaluation. Clinically, these patients can have obscure bleeding (which may be noted in the form of visible passage of blood or melena or occult bleeding, unexplained iron deficiency anemia, or guaiac-positive stools without visible passage of blood). Small bowel pathology is the frequent source of bleeding in these patients.

Q. RBC Scan Abdomen and Pelvis

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

A. Arteriography Visceral

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

B. CT Abdomen and Pelvis With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

C. CT Abdomen and Pelvis Without and With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

D. CT Abdomen and Pelvis Without IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

E. CT Abdomen With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

F. CT Abdomen Without and With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

G. CT Abdomen Without IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

H. CT Enterography

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

I. CTA Abdomen and Pelvis With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

J. CTA Abdomen and Pelvis Without and With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

K. CTA Abdomen With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

L. CTA Abdomen Without and With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

M. CTA Chest With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

N. CTA Chest Without and With IV Contrast

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

O. Fluoroscopy Upper GI Series

Variant 5: This variant is applicable to postsurgical or trauma patients with UGIB, contraindicated to upper GI endoscopy.

P. RBC Scan Abdomen and Pelvis

Summary of Recommendations

Supporting Documents

The evidence table, literature search, and appendix for this topic are available at <https://acsearch.acr.org/list>. The appendix includes the strength of evidence assessment and the final rating round tabulations for each recommendation.

For additional information on the Appropriateness Criteria methodology and other supporting documents, please go to the ACR website at <https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Appropriateness-Criteria>.

Appropriateness Category Names and Definitions

Appropriateness Category Name	Appropriateness Rating	Appropriateness Category Definition
Usually Appropriate	7, 8, or 9	The imaging procedure or treatment is indicated in the specified clinical scenarios at a favorable risk-benefit ratio for patients.
May Be Appropriate	4, 5, or 6	The imaging procedure or treatment may be indicated in the specified clinical scenarios as an alternative to imaging procedures or treatments with a more favorable risk-benefit ratio, or the risk-benefit ratio for patients is equivocal.
May Be Appropriate (Disagreement)	5	The individual ratings are too dispersed from the panel median. The different label provides transparency regarding the panel's recommendation. "May be appropriate" is the rating category and a rating of 5 is assigned.
Usually Not Appropriate	1, 2, or 3	The imaging procedure or treatment is unlikely to be indicated in the specified clinical scenarios, or the risk-benefit ratio for patients is likely to be unfavorable.

Relative Radiation Level Information

References

1. Lanas A, Dumonceau JM, Hunt RH, et al. Non-variceal upper gastrointestinal bleeding. *Nat Rev Dis Primers* 2018;4:18020.
2. Gerson LB, Fidler JL, Cave DR, Leighton JA. ACG Clinical Guideline: Diagnosis and Management of Small Bowel Bleeding. *Am J Gastroenterol.* 110(9):1265-87; quiz 1288, 2015 Sep.
3. Ichita C, Sasaki A, Sumida C, et al. Clinical and endoscopic features of aorto-duodenal fistula resulting in its definitive diagnosis: an observational study. *BMC Gastroenterol.* 21(1):45, 2021 Feb 01.
4. Feng W, Yue D, ZaiMing L, ZhaoYu L, Wei L, Qiyong G. Hemobilia following laparoscopic cholecystectomy: computed tomography findings and clinical outcome of transcatheter arterial embolization. *Acta Radiol.* 58(1):46-52, 2017 Jan.
5. Guglielmo FF, Wells ML, Bruining DH, et al. Gastrointestinal Bleeding at CT Angiography and CT Enterography: Imaging Atlas and Glossary of Terms. *Radiographics.* 41(6):1632-1656, 2021 Oct. *Radiographics.* 41(6):1632-1656, 2021 Oct.
6. Mullady DK, Wang AY, Waschke KA. AGA Clinical Practice Update on Endoscopic Therapies for Non-Variceal Upper Gastrointestinal Bleeding: Expert Review. [Review]. *Gastroenterology.* 159(3):1120-1128, 2020 09. *Gastroenterology.* 159(3):1120-1128, 2020 09.
7. Wells ML, Hansel SL, Bruining DH, et al. CT for Evaluation of Acute Gastrointestinal Bleeding. *Radiographics.* 2018;38(4):1089-1107.
8. Laine L, Barkun AN, Saltzman JR, Martel M, Leontiadis GI. Correction to: ACG Clinical Guideline: Upper Gastrointestinal and Ulcer Bleeding. *Am J Gastroenterol* 2021;116:2309.
9. Kim CY, Pinchot JW, Ahmed O, et al. ACR Appropriateness Criteria® Radiologic Management of Gastric Varices. *J Am Coll Radiol* 2020;17:S239-S54.
10. Pinchot JW, Kalva SP, Majdalany BS, et al. ACR Appropriateness Criteria® Radiologic Management of Portal Hypertension. *J Am Coll Radiol* 2021;18:S153-S73.
11. Trabzonlu TA, Mozaffary A, Kim D, Yaghmai V. Dual-energy CT evaluation of gastrointestinal bleeding. [Review]. *Abdom Radiol.* 45(1):1-14, 2020 01.
12. Mohammadinejad P, Kwapisz L, Fidler JL, et al. The utility of a dual-phase, dual-energy CT protocol in patients presenting with overt gastrointestinal bleeding. *Acta Radiol Open.* 10(7):20584601211030658, 2021 Jul.
13. American College of Radiology. ACR–NASCI–SIR–SPR Practice Parameter for the Performance and Interpretation of Body Computed Tomography Angiography (CTA). Available at: <https://gravitas.acr.org/PPTS/GetDocumentView?docId=164+&releaseId=2>.
14. Kim G, Soto JA, Morrison T. Radiologic Assessment of Gastrointestinal Bleeding. [Review]. *Gastroenterol Clin North Am.* 47(3):501-514, 2018 Sep.
15. Choi C, Lim H, Kim MJ, et al. Relationship between angiography timing and angiographic visualization of extravasation in patients with acute non-variceal gastrointestinal bleeding. *BMC Gastroenterol.* 20(1):426, 2020 Dec 14.
16. Chevallier O, Comby PO, Guillen K, et al. Efficacy, safety and outcomes of transcatheter

- arterial embolization with N-butyl cyanoacrylate glue for non-variceal gastrointestinal bleeding: A systematic review and meta-analysis. *Diagnostic and Interventional Imaging*. 102(7-8):479-487, 2021 Jul-Aug. *Diagn Interv Imaging*. 102(7-8):479-487, 2021 Jul-Aug.
17. Lai HY, Wu KT, Liu Y, Zeng ZF, Zhang B. Angiography and transcatheter arterial embolization for non-variceal gastrointestinal bleeding. *Scandinavian Journal of Gastroenterology*. 55(8):931-940, 2020 Aug. *Scand J Gastroenterol*. 55(8):931-940, 2020 Aug.
 18. Shotar E, Soyer P, Barat M, et al. Diagnosis of acute overt gastrointestinal bleeding with CT-angiography: Comparison of the diagnostic performance of individual acquisition phases. *Diagnostic and Interventional Imaging*. 98(12):857-863, 2017 Dec. *Diagn Interv Imaging*. 98(12):857-863, 2017 Dec.
 19. Garcia-Blazquez V, Vicente-Bartulos A, Olavarria-Delgado A, Plana MN, van der Winden D, Zamora J. Accuracy of CT angiography in the diagnosis of acute gastrointestinal bleeding: systematic review and meta-analysis. *Eur Radiol*. 2013;23(5):1181-1190.
 20. Fidler JL, Gunn ML, Soto JA, et al. Society of abdominal radiology gastrointestinal bleeding disease-focused panel consensus recommendations for CTA technical parameters in the evaluation of acute overt gastrointestinal bleeding. *Abdom Radiol (NY)*. 2019 Sep;44(9):2957-2962.
 21. Sun H, Hou XY, Xue HD, et al. Dual-source dual-energy CT angiography with virtual non-enhanced images and iodine map for active gastrointestinal bleeding: image quality, radiation dose and diagnostic performance. *European Journal of Radiology*. 84(5):884-91, 2015 May. *Eur J Radiol*. 84(5):884-91, 2015 May.
 22. Chan V, Tse D, Dixon S, et al. Outcome following a negative CT Angiogram for gastrointestinal hemorrhage. *Cardiovasc Intervent Radiol*. 38(2):329-35, 2015 Apr.
 23. Tse JR, Shen J, Shah R, Fleischmann D, Kamaya A. Extravasation Volume at Computed Tomography Angiography Correlates With Bleeding Rate and Prognosis in Patients With Overt Gastrointestinal Bleeding. *Investigative Radiology*. 56(6):394-400, 2021 06 01. *Invest Radiol*. 56(6):394-400, 2021 06 01.
 24. Stewart K, Sharma AK. The utilization of CTA in management of gastrointestinal bleeding in a tertiary care center ED. Are we using it enough?. *American Journal of Emergency Medicine*. 39:60-64, 2021 01. *Am J Emerg Med*. 39:60-64, 2021 01.
 25. Hsu MJ, Dinh DC, Shah NA, et al. Time to conventional angiography in gastrointestinal bleeding: CT angiography compared to tagged RBC scan. *Abdominal Radiology*. 45(2):307-311, 2020 02. *Abdom Radiol*. 45(2):307-311, 2020 02.
 26. Alrashidi I, Kim TH, Shin JH, Alreshidi M, Park M, Jang EB. Efficacy and safety of transcatheter arterial embolization for active arterial esophageal bleeding: a single-center experience. *Diagnostic & Interventional Radiology*. 27(4):519-523, 2021 Jul. *Diagn Interv Radiol*. 27(4):519-523, 2021 Jul.
 27. Aoki M, Tokue H, Koyama Y, Tsushima Y, Oshima K. Transcatheter arterial embolization with N-butyl cyanoacrylate for arterial esophageal bleeding in esophageal cancer patients. *World Journal of Surgical Oncology*. 14:54, 2016 Feb 24. *World J Surg Oncol*. 14:54, 2016 Feb 24.
 28. Kohli DR, Shah RD, Komorowski DJ, Smallfield GB. Periesophageal Pseudoaneurysms: Rare Cause of Refractory Bleeding Treated with Transarterial Embolization. *Case Rep Gastrointest Med*. 2016:1456949, 2016.

29. Zheng L, Lee IJ, Shin JH, Chu HH, Li HL. Endovascular Management of Gastric Conduit Hemorrhage following Transthoracic Esophagectomy. *Journal of Vascular & Interventional Radiology*. 32(8):1144-1149, 2021 Aug. *J Vasc Interv Radiol*. 32(8):1144-1149, 2021 Aug.
30. Cho SB, Hur S, Kim HC, et al. Transcatheter arterial embolization for advanced gastric cancer bleeding: A single-center experience with 58 patients. *Medicine*. 99(15):e19630, 2020 Apr. *Medicine (Baltimore)*. 99(15):e19630, 2020 Apr.
31. Koo HJ, Shin JH, Shin S, Yoon HK, Ko GY, Gwon DI. Efficacy and Clinical Outcomes of Transcatheter Arterial Embolization for Gastrointestinal Bleeding from Gastrointestinal Stromal Tumor. *J Vasc Interv Radiol*. 26(9):1297-304.e1, 2015 Sep.
32. Sun CJ, Wang CE, Wang YH, Xie LL, Liu TH, Ren WC. Transcatheter arterial embolization of acute gastrointestinal tumor hemorrhage with Onyx. *Indian Journal of Cancer*. 51 Suppl 2:e56-9, 2015 Feb. *Indian J Cancer*. 51 Suppl 2:e56-9, 2015 Feb.
33. Lee SM, Jeong SY, Shin JH, et al. Transcatheter arterial embolization for gastrointestinal bleeding related to pancreatic adenocarcinoma: clinical efficacy and predictors of clinical outcome. *European Journal of Radiology*. 123:108787, 2020 Feb. *Eur J Radiol*. 123:108787, 2020 Feb.
34. Zandrino F, Tettoni SM, Gallesio I, Summa M. Emergency arterial embolization of upper gastrointestinal and jejunal tumors: An analysis of 12 patients with severe bleeding. *Diagnostic and Interventional Imaging*. 98(1):51-56, 2017 Jan. *Diagn Interv Imaging*. 98(1):51-56, 2017 Jan.
35. Nykanen T, Peltola E, Kylanpaa L, Udd M. Bleeding gastric and duodenal ulcers: case-control study comparing angioembolization and surgery. *Scandinavian Journal of Gastroenterology*. 52(5):523-530, 2017 May. *Scand J Gastroenterol*. 52(5):523-530, 2017 May.
36. Laursen SB, Jakobsen M, Nielsen MM, Hovendal C, Schaffalitzky de Muckadell OB. Transcatheter arterial embolization is the first-line therapy of choice in peptic ulcer bleeding not responding to endoscopic therapy. *Scandinavian Journal of Gastroenterology*. 50(3):264-71, 2015 Mar. *Scand J Gastroenterol*. 50(3):264-71, 2015 Mar.
37. Mille M, Huber J, Wlasak R, et al. Prophylactic Transcatheter Arterial Embolization After Successful Endoscopic Hemostasis in the Management of Bleeding Duodenal Ulcer. *Journal of Clinical Gastroenterology*. 49(9):738-45, 2015 Oct. *J Clin Gastroenterol*. 49(9):738-45, 2015 Oct.
38. Yao Z, Tian W, Xu X, et al. Transcatheter Arterial Embolization in the Treatment of Abdominal Bleeding in Patients Being Treated with Open Abdomen Due to Duodenal Fistula. *World J Surg*. 44(8):2562-2571, 2020 08.
39. Wen F, Dong Y, Lu ZM, Liu ZY, Li W, Guo QY. Hemobilia After Laparoscopic Cholecystectomy: Imaging Features and Management of an Unusual Complication. *Surg Laparosc Endosc Percutan Tech*. 26(1):e18-24, 2016 Feb.
40. Lee NJ, Shin JH, Lee SS, Park DH, Lee SK, Yoon HK. Transcatheter arterial embolization for iatrogenic bleeding after endoscopic ultrasound-guided pancreaticobiliary drainage. *Diagnostic and Interventional Imaging*. 99(11):717-724, 2018 Nov. *Diagn Interv Imaging*. 99(11):717-724, 2018 Nov.
41. Huprich JE, Barlow JM, Hansel SL, Alexander JA, Fidler JL. Multiphase CT enterography evaluation of small-bowel vascular lesions. *AJR Am J Roentgenol* 2013;201:65-72.

42. Zhang, Sun, Xue, Li XG, Jin ZY. Computed Tomography Signs for Active Severe Gastrointestinal Bleeding. *Chung Kuo I Hsueh Ko Hsueh Yuan Hsueh Pao*. 38(3):322-6, 2016 06 10.
43. Tseng CM, Lin IC, Chang CY, et al. Role of computed tomography angiography on the management of overt obscure gastrointestinal bleeding. *PLoS ONE* [Electronic Resource]. 12(3):e0172754, 2017.*PLoS ONE*. 12(3):e0172754, 2017.
44. Kokoroskos N, Naar L, Peponis T, et al. Provocative Angiography, Followed by Therapeutic Interventions, in the Management of Hard-To-Diagnose Gastrointestinal Bleeding. *World Journal of Surgery*. 44(9):2944-2949, 2020 09.*World J Surg*. 44(9):2944-2949, 2020 09.
45. Shin JH. Refractory gastrointestinal bleeding: role of angiographic intervention. *Clin Endosc* 2013;46:486-91.
46. Wildgruber M, Wrede CE, Zorger N, et al. Computed tomography versus digital subtraction angiography for the diagnosis of obscure gastrointestinal bleeding. *European Journal of Radiology*. 88:8-14, 2017 Mar.*Eur J Radiol*. 88:8-14, 2017 Mar.
47. Kawabata H, Kawakatsu Y, Sone D, et al. A rare case of Goodpasture syndrome concomitant with bleeding jejunal Dieulafoy's lesion. *Clinical Journal of Gastroenterology*. 13(3):382-385, 2020 Jun.*Clin J Gastroenterol*. 13(3):382-385, 2020 Jun.
48. Batouli A, Kazemi A, Hartman MS, Heller MT, Midian R, Lupetin AR. Dieulafoy lesion: CT diagnosis of this lesser-known cause of gastrointestinal bleeding. *Clin Radiol* 2015;70:661-6.
49. Aksoy T. Obscure and occult gastrointestinal bleeding: role of radionuclide imaging. *Abdom Imaging* 2012;37:309-10; author reply 11-2.
50. Shukla PA, Zybulewski A, Kolber MK, Berkowitz E, Silberzweig J, Hayim M. No catheter angiography is needed in patients with an obscure acute gastrointestinal bleed and negative CTA. *Clinical Imaging*. 43:106-109, 2017 May - Jun.*Clin Imaging*. 43:106-109, 2017 May - Jun.
51. Pai M, Frampton AE, Virk JS, et al. Preoperative superselective mesenteric angiography and methylene blue injection for localization of obscure gastrointestinal bleeding. *JAMA surgery* 2013;148:665-8.
52. Agrawal JR, Travis AC, Morteale KJ, et al. Diagnostic yield of dual-phase computed tomography enterography in patients with obscure gastrointestinal bleeding and a non-diagnostic capsule endoscopy. *Journal of gastroenterology and hepatology* 2012;27:751-9.
53. Chu Y, Wu S, Qian Y, et al. Complimentary Imaging Modalities for Investigating Obscure Gastrointestinal Bleeding: Capsule Endoscopy, Double-Balloon Enteroscopy, and Computed Tomographic Enterography. *Gastroenterol Res Pract* 2016;2016:8367519.
54. Huprich JE, Fletcher JG, Fidler JL, et al. Prospective blinded comparison of wireless capsule endoscopy and multiphase CT enterography in obscure gastrointestinal bleeding. *Radiology*. 2011;260(3):744-751.
55. Jeon SR, Jin-Oh K, Gun KH, et al. Is there a difference between capsule endoscopy and computed tomography as a first-line study in obscure gastrointestinal bleeding? *The Turkish journal of gastroenterology : the official journal of Turkish Society of Gastroenterology* 2014;25:257-63.
56. He B, Gong S, Hu C, et al. Obscure gastrointestinal bleeding: diagnostic performance of 64-

section multiphase CT enterography and CT angiography compared with capsule endoscopy. *The British journal of radiology* 2014;87:20140229.

57. Chang WC, Tsai SH, Chang WK, et al. The value of multidetector-row computed tomography for localization of obscure acute gastrointestinal bleeding. *Eur J Radiol.* 80(2):229-35, 2011 Nov.
58. Yen HH, Chen YY, Yang CW, Liu CK, Soon MS. Clinical impact of multidetector computed tomography before double-balloon enteroscopy for obscure gastrointestinal bleeding. *World J Gastroenterol* 2012;18:692-7.
59. Casciani E, Nardo GD, Chin S, et al. MR Enterography in paediatric patients with obscure gastrointestinal bleeding. *Eur J Radiol.* 93:209-216, 2017 Aug.
60. Bocker U, Dinter D, Litterer C, et al. Comparison of magnetic resonance imaging and video capsule enteroscopy in diagnosing small-bowel pathology: localization-dependent diagnostic yield. *Scand J Gastroenterol.* 45(4):490-500, 2010 Apr.
61. Wiarda BM, Heine DG, Mensink P, et al. Comparison of magnetic resonance enteroclysis and capsule endoscopy with balloon-assisted enteroscopy in patients with obscure gastrointestinal bleeding. *Endoscopy.* 2012;44(7):668-673.
62. Tabibian JH, Wong Kee Song LM, Enders FB, Aguet JC, Tabibian N. Technetium-labeled erythrocyte scintigraphy in acute gastrointestinal bleeding. *Int J Colorectal Dis.* 2013;28(8):1099-1105.
63. Dolezal J, Vizda J, Kopacova M. Single-photon emission computed tomography enhanced Tc-99m-pertechnetate disodium-labelled red blood cell scintigraphy in the localization of small intestine bleeding: a single-centre twelve-year study. *Digestion.* 2011;84(3):207-211.
64. Otomi Y, Otsuka H, Terazawa K, et al. The diagnostic ability of SPECT/CT fusion imaging for gastrointestinal bleeding: a retrospective study. *BMC Gastroenterol.* 18(1):183, 2018 Dec 10.
65. Park S, Jeong B, Shin JH, Jang EH, Hwang JH, Kim JH. Transarterial embolisation for gastroduodenal bleeding following endoscopic resection. *British Journal of Radiology.* 94(1122):20210062, 2021 Jun 01. *Br J Radiol.* 94(1122):20210062, 2021 Jun 01.
66. Hsia CC, Wang CY, Huang JF, et al. Diagnostic Accuracy of Computed Tomography for the Prediction of the Need for Laparotomy for Traumatic Hollow Viscus Injury: Systematic Review and Meta-Analysis. *J Pers Med* 2021;11.
67. Murugesan SD, Sathyanesan J, Lakshmanan A, et al. Massive hemobilia: a diagnostic and therapeutic challenge. *World Journal of Surgery.* 38(7):1755-62, 2014 Jul. *World J Surg.* 38(7):1755-62, 2014 Jul.
68. Rebibo L, Fuks D, Blot C, et al. Gastrointestinal bleeding complication of gastric fistula after sleeve gastrectomy: consider pseudoaneurysms. *Surg Endosc* 2013;27:2849-55.
69. Wu CH, Wang LJ, Wong YC, et al. Contrast-enhanced multiphasic computed tomography for identifying life-threatening mesenteric hemorrhage and transmural bowel injuries. *J Trauma.* 71(3):543-8, 2011 Sep.
70. Luo J, Tang W, Wang M, Xiao Y, Tan M, Jiang C. Case series of aortoenteric fistulas: a rare cause of gastrointestinal bleeding. *BMC Gastroenterology.* 21(1):49, 2021 Feb 02. *BMC Gastroenterol.* 21(1):49, 2021 Feb 02.
71. Deijen CL, Smulders YM, Coveliers HME, Wisselink W, Rauwerda JA, Hoksbergen AWJ. The

Importance of Early Diagnosis and Treatment of Patients with Aortoenteric Fistulas Presenting with Herald Bleeds. *Annals of Vascular Surgery*. 36:28-34, 2016 Oct. *Ann Vasc Surg*. 36:28-34, 2016 Oct.

72. American College of Radiology. ACR Appropriateness Criteria® Radiation Dose Assessment Introduction. Available at: <https://edge.sitecorecloud.io/americancoldf5f-acrorgf92a-productioncb02-3650/media/ACR/Files/Clinical/Appropriateness-Criteria/ACR-Appropriateness-Criteria-Radiation-Dose-Assessment-Introduction.pdf>.

Disclaimer

The ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those examinations generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

^aUniversity of Wisconsin School of Medicine and Public Health, Madison, Wisconsin. ^bNYU Grossman School of Medicine, New York, New York. ^cPanel Chair, Brigham & Women's Hospital, Boston, Massachusetts. ^dPanel Chair, University of California San Diego, San Diego, California. ^ePanel Vice-Chair, Massachusetts General Hospital and Harvard Medical School, Boston, Massachusetts. ^fGlobal Advanced Imaging, PLLC, Little Rock, Arkansas; Commission on Nuclear Medicine and Molecular Imaging. ^gUniversity of Texas Health Science Center at Houston and McGovern Medical School, Houston, Texas; American Gastroenterological Association. ^hMayo Clinic, Rochester, Minnesota; Society for Cardiovascular Magnetic Resonance. ⁱOchsner Hospital, Baton Rouge, Louisiana. ^jUMass Memorial Health and UMass Chan Medical School, Worcester, Massachusetts; Committee on Emergency Radiology-GSER. ^kUniversity of North Carolina at Chapel Hill, Chapel Hill, North Carolina. ^lVA Puget Sound Health Care System and University of Washington, Seattle, Washington. ^mUniversity of California San Diego, San Diego, California. ⁿSpecialty Chair, University of Wisconsin School of Medicine and Public Health, Madison, Wisconsin. ^oSpecialty Chair, Brigham & Women's Hospital, Boston, Massachusetts.