

# ACR PRACTICE PARAMETER FOR THE PERFORMANCE OF PREOPERATIVE IMAGE-GUIDED LOCALIZATION IN THE BREAST

The American College of Radiology, with more than 40,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

## PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care<sup>1</sup>. For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

---

<sup>1</sup> *Iowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing*, 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the "ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures (Revised 2008)" sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, *Stanley v. McCarver*, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

## I. INTRODUCTION

Preoperative image-guided localization of breast abnormalities before surgical excision is currently the standard of care for nonpalpable lesions. Localization devices guide appropriate excision and provide a surgeon with the best means to ensure successful removal of the target tissue. Preoperative localization with image-guided wire placement has been a standard of breast imaging diagnosis and treatment since its development in the 1970s and remains a reliable and safe method for localization [1]. Several recent advancements in nonwire localization (NWL) techniques minimize limitations of wire localization and potentially improve patient care and clinical workflow. New technological approaches and devices are continually being developed and introduced that will expand the variety of localization tools.

## II. INDICATIONS

Presurgical localization in the breast may be performed for patients with selected indications including, but not limited to:

1. Biopsy-proven cancer
2. Biopsy-proven metastatic lymphadenopathy
3. High-risk lesions diagnosed at percutaneous biopsy
4. Imaging-pathological discordance at core-needle biopsy
5. Cases in which core-needle biopsy is not an option or fails to provide a definitive histological diagnosis

## III. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

Interpreting physicians, medical physicists, and radiological technologists who work in breast imaging must meet the requirements that are appropriate to the scope of their practice as outlined in the following documents or practice parameters:

1. Mammography Quality Standards Act Final Regulations [2]
2. [ACR Practice Parameter for the Performance of Stereotactic/Tomosynthesis-Guided Breast Interventional Procedures](#) [3]
3. [ACR Practice Parameter for the Performance of Ultrasound-Guided Percutaneous Breast Interventional Procedures](#) [4]
4. [ACR Practice Parameter for the Performance of a Diagnostic Breast Ultrasound Examination](#) [5]
5. [ACR Practice Parameter for the Performance of Contrast-Enhanced Magnetic Resonance Imaging \(MRI\) of the Breast](#) [6]
6. [ACR Practice Parameter for the Performance of Magnetic Resonance Imaging-Guided Breast Interventional Procedures](#) [7]
7. [ACR Practice Parameter for the Performance of Screening and Diagnostic Mammography](#) [8]

## IV. SPECIFICATIONS OF THE PROCEDURE

Prior to localization, the radiologist should review all pertinent imaging examinations to determine the extent of the target. Review should determine whether biopsy markers, if deployed at the time of prior biopsy, were placed in the appropriate position or whether they have migrated. In patients who have undergone neoadjuvant therapy, the original extent of disease and the visible residual disease are both important to consider. The localization may target a biopsy marker and/or the spectrum of breast imaging abnormalities: mass, calcifications, asymmetry, architectural distortion. If there is known malignancy, it is necessary to understand the extent of malignancy and its location with respect to previously placed biopsy marker(s). More than one guidance device may be used to bracket the extent of disease [9,10]. The use of multiple localizing devices can decrease the number of procedures required to obtain clear lumpectomy margins and increase the rate of breast conservation versus mastectomy [9].

Benefits, limitations, and risks of the procedure as well as alternative procedures should be discussed with the patient. Informed consent should be obtained and documented. Adherence to the Joint Commission's Universal Protocol for Preventing Wrong Site, Wrong Procedure, Wrong Person Surgery™ is required for procedures in nonoperating room settings, including bedside procedures. The organization should have processes and systems in place for reconciling differences in staff responses during the time-out. The breast, imaging equipment, modality in which the procedure is to be performed, and physician performing the procedure should be prepared in conformity with the principles of infection control.

## A. Localization Techniques

### 1. Wire localizations

Preoperative wire localization using mammographic, sonographic, or magnetic resonance imaging (MRI) guidance is optimally performed the day of surgery. The wire may be placed at the breast lesion, adjacent to the biopsy marker (if it lies at the site of the lesion), or at the postbiopsy hematoma if the lesion itself cannot be visualized and if a biopsy marker is not present. If ultrasound guidance is used to place the wire, marking the location of the lesion on the overlying skin with the patient in the supine operative position and measuring the depth of the lesion can assist the surgeon during excision. Tomosynthesis-guided localization is an emerging alternative that may be used by some practices and differs slightly in technique from standard orthogonal mammographic practice. Limitations of wire localizations include need for placement on the day of surgery, imprecision that results from variability in positioning by the radiologist, and inadvertent wire displacement (during patient transfer, postprocedure mammography, or surgical positioning) [11]. An important feature for some wire localization devices is the potential opportunity for repositioning or removal and reinsertion with another wire, if improper localization has occurred. There are certain wires that have FDA approval for deployment under MRI guidance.

### 2. NWL Devices

Alternative forms of preoperative localization that do not use wire and mitigate wire localization limitations are now available and include radioactive seed, radiofrequency reflector, magnetic seed, and radiofrequency identification (RFID), among others. These localization devices lack a component external to the breast after placement (as is present with the proximal wire segment when a wire is placed). The absence of an external component in NWL offers increased patient comfort and decreased risk of displacement or transection of the localizing device compared with wires [12]. In addition, the devices can be placed before the day of surgery. This uncoupling of localization from the day of surgery provides flexibility to both the radiologist and surgeon in the localization and surgical procedures and less waiting time for patients [13-15]. All forms of NWL typically have two components: a single-use sterilized device preloaded into a needle introducer and a console with a handheld probe for the detection of deployment by the radiologist and for surgical guidance by the surgeon. The localizing device may be placed at the breast lesion, adjacent to the biopsy marker (if it lies at the site of the lesion), or at the postbiopsy hematoma if the lesion itself cannot be visualized and if a biopsy marker is not present. In addition to some type-specific limitations (outlined below), NWL may be subject to imprecise positioning during placement or deployment. An important limitation of NWL is that once deployed, they currently cannot be repositioned.

#### a. Radioactive seed localization (RSL)

The radioactive seed placed for localization is currently composed of a titanium capsule containing iodine-125. The radioactive seed is inserted through a needle with sonographic or mammographic guidance. Because the iodine-125-labeled seed half-life is sufficiently long (59.4 days), preoperative RSL theoretically can be performed up to several weeks prior to surgery but is usually performed within a few days before surgery to minimize radiation exposure. Different facilities may have different maximum length of time a seed can stay inserted, depending on the source strength of their seeds.

RSL programs require environmental safety processes and adherence to regulations for

nuclear materials under the Nuclear Regulatory Commission (NRC) or under Agreement States. A lost radioactive seed is a reportable medical event, and an established protocol is needed to manage the event. In addition, a migrated radioactive seed in the breast must be recovered [1,11,15-19]. A written directive, signed by an authorized user, may be required for each procedure by regulation. The information on the written directives may include the patient's name, the treatment site, the activity and type of radionuclide, the number of sources implanted, the total source strength, and the exposure time.

b. Radar reflector

Radar localization technology was introduced in 2014. The reflector is inserted via a needle under sonographic or mammographic guidance. The reflector is passive until activated with infrared light from a handheld probe. Once activated, the device reflects the radar signal, which is detected by the probe and recorded by the console. The console provides audible and visual indicators of the probe's proximity to the reflector. The reflector can be placed at any time prior to surgery. There is no limitation on the length of time the reflector can stay in the breast, providing the opportunity for placement prior to initiation of neoadjuvant therapy. Reflectors placed at significant depth or within a postbiopsy hematoma may not produce a detectable signal to the skin [1,19-21].

c. Magnetic seed

Magnetic seed localization technology was introduced in 2016. The localization device is currently made from stainless steel. The seed is not magnetic, but it is induced to become a magnet under the influence of the handheld probe that transiently magnetizes the seed. The seed is inserted via a needle under sonographic or mammographic guidance. The console provides audible and visual indications of the probe's proximity to the seed. The seed can be placed any time prior to surgery. There is no restriction on the length of time the seed can remain in the breast, providing the opportunity for placement prior to initiation of neoadjuvant therapy. Following deployment, the patient can have an MRI, albeit with significant artifact. Nonmagnetic surgical tools need to be used while the probe is in use, and certain stainless-steel instruments may not be compatible with the system. Compared with wires and other NWL devices, the magnetic seed is more resistant to damage on deployment, following implantation, or with electrocautery during surgery. There is no limitation on depth placement of magnetic seeds for detectable signal [1,19,22-24].

d. RFID tag

RFID technology was introduced in 2017. The RFID tag currently is made of a copper-wrapped ferrite rod and microprocessor within a glass casing enclosed within a polypropylene sheath to prevent migration [25]. The device absorbs, modifies, and re-emits a 134.2-kHz radiofrequency signal that is sent by the handheld reader device [25]. An integrated loop probe on the device has a 6-cm detection range while the disposable sterile surgical pencil-sized probe has a 3-cm detection range. The RFID tag is inserted via a needle under sonographic or mammographic guidance. The console provides audible and visual indicators that indicate the probe's proximity to the seed. A unique identification number for an individual tag is able to be displayed on the device reader, which is useful to distinguish between tags if multiple devices are implanted within the same breast.

In each of the NWL methods, more than one localizing device may be placed to bracket the full extent of disease in patients with large masses, masses with satellite nodules or accompanying microcalcifications extending from the mass, or segmental or linearly distributed microcalcifications alone. When two or more localizing devices are used, each manufacturer recommends a specific minimum distance between the devices in order to discriminate between them.

In general, postlocalization preoperative orthogonal mammograms should be obtained to

depict the localization and to guide the surgeon in the operative procedure. However, in the rare occasion of young women undergoing ultrasound-guided localizations, some practices will only use ultrasound to document placement of localizing device. A radiation detector (eg, a Geiger counter or a gamma probe) can be utilized to confirm the presence of a radioactive seed that was inserted using ultrasound alone. Confirmation with a radiation detector is usually possible when more than one seed is placed, although it may not be able to distinguish two seeds close together [18]. Radiologists may elect to annotate the target and specify the final relation of the localization device to the target for the surgeon on the postlocalization mammogram. In all forms of preoperative localization, communication with the surgeon may avoid misunderstanding and may take the form of a telephone call, written comments, or annotation of the images.

## B. Specimen Imaging

Specimen radiography is essential to document removal of the target and localization device and provide guidance to the surgeon as to the adequacy of excision [26,27]. This should occur while the patient is still in the operating room so that the surgeon can remove more tissue if warranted.

If the lesion is a single mass, particularly if it was mammographically occult, ultrasound of the specimen can be used to document mass removal [28]. If the lesion contains microcalcifications, either extending from a mass or alone, specimen radiography is better to evaluate the adequacy of excision. When tumor can be seen extending to the specimen margins on the specimen radiograph, there is a high positive predictive value for residual tumor in the breast. Conversely, the negative predictive value of clear margins on specimen radiography is low but may be improved with the addition of orthogonal or tomosynthesis views [29]. Therefore, even though the tumor may not appear to extend to the margins of the resected specimen on the specimen radiograph, the residual tumor may still be present in the breast. This may be particularly true for noncalcified ductal carcinoma in situ and infiltrating lobular carcinoma [30,31]. When the radiologist reviews the specimen radiograph, any unexpected findings should be communicated to the surgeon.

## C. Targeted axillary dissection

In an effort to minimize morbidity from complete axillary dissection, targeted axillary dissection with removal of sentinel lymph node and the index biopsy-proven metastatic node has been reported in patients who have undergone neoadjuvant therapy [32]. If a patient converts to node-negative status after therapy, the patient can potentially be spared complete nodal dissection. In this procedure, a pathology-proven metastatic lymph node with biopsy marker may be localized following neoadjuvant therapy. Routine intraoperative lymphatic mapping is performed along with removal of the localized metastatic lymph node. This combined sentinel lymph node dissection with localized removal of metastatic lymph node has a false-negative rate for axillary staging below 5% and provides a potentially safe way to limit axillary surgery [33].

Identifying and localizing a biopsy-proven clipped metastatic axillary node with imaging can be technically challenging following neoadjuvant therapy, particularly if the patient has had complete imaging response. There is increasing interest in taking advantage of the benefit of extended duration of implantation of several NWLs (radar reflector, magnetic seed, RFID tag) and placing the device within a biopsy-proven metastatic axillary node prior to initiating or early in the course of neoadjuvant therapy in order to facilitate accurate identification and localization for targeted axillary dissection.

## V. DOCUMENTATION

Reporting should be in accordance with the [ACR Practice Parameter for Communication of Diagnostic Imaging Findings](#) [34].

Permanent records of image-guided breast localizations should be documented in a retrievable image storage format.

A. Image labeling should include permanent identification containing the following:

1. Patient's first and last names
2. Identifying number and/or date of birth
3. Examination date
4. Facility name and location
5. Designation of left or right breast
6. Annotation of mammographic view (eg, craniocaudal, mediolateral oblique, 90° mediolateral)
7. Annotation of sonographic image: Anatomic location using clockface notation; a labeled diagram of the breast may be included, distance from the nipple to the lesion in centimeters, and transducer orientation
8. Technologist's identification number or initials

B. The physician's report of image-guided localizations should include the following:

1. Procedure performed
2. Designation of the left or right breast
3. Description and location of the lesion
4. Safety time-out having been performed
5. Approach used
6. Type and amount of local anesthesia
7. Skin incision, if made
8. Type of localization device
9. Confirmation of postprocedure mammogram documenting accurate localizing device placement, location, and number of devices with respect to the targeted lesion
10. Complications and treatment, if any
11. Confirmation of specimen imaging (if not detailed in a separate report)

C. Retention of the procedure images, including specimen images if obtained, should be consistent with the facility's policies for retention of mammograms and in compliance with federal and state regulations.

## **VI. QUALITY CONTROL, AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION**

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *ACR Position Statement on Quality Control and Improvement, Safety, Infection Control and Patient Education* on the ACR website (<https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement>).

## **VII. EQUIPMENT**

Equipment requirements are outlined in the following practice parameters:

1. [ACR Practice Parameter for the Performance of Stereotactic/Tomosynthesis-Guided Breast Interventional Procedures](#) [3]
2. [ACR Practice Parameter for the Performance of Ultrasound-Guided Percutaneous Breast Interventional Procedures](#) [4]
3. [ACR Practice Parameter for the Performance of Magnetic Resonance Imaging-Guided Breast Interventional Procedure](#) [7]

## **ACKNOWLEDGEMENTS**

This practice parameter was developed according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (<https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards>) by the Committee on Practice Parameters – Breast Imaging of the ACR Commission on Breast Imaging.

Writing Committee

Karla A. Sepulveda, MD, FACR, Chair

Mary S. Newell, MD, FACR

Phoebe Freer, MD

Haydee Ojeda-Fournier, MD

Catherine S. Giess, MD

Dana H. Smetherman, MD

Cindy Lee, MD

Karen S. Zheng, MD

Linda Moy, MD

#### Committee on Practice Parameters – Breast Imaging

(ACR Committee responsible for sponsoring the draft through the process)

Linda Moy, MD, Chair

Madison Kocher, MD

Catherine S. Giess, MD, Vice-Chair

Cindy S. Lee, MD

Shadi Aminololama-Shakeri, MD

John M. Lewin, MD, FACR

Catherine Margaret Appleton, MD

Vilert A Loving, MD

Stamatia V Destounis, MD, FACR

Haydee Ojeda-Fournier, MD

Dipti Gupta, MD

Roberta M. Strigel, MD

Carolyn A. Haerr, MD, FACR

Dana H. Smetherman, MD, FACR, Chair, Commission on Breast Imaging

David B. Larson, MD, MBA, Chair, Commission on Quality and Safety

Mary S. Newell, MD, FACR, Chair, Committee on Practice Parameters and Technical Standards

#### Comment Reconciliation Committee

David C. Youmans, MD, FACR, Chair

Cindy Lee, MD



## Comment Reconciliation Committee

Taj Kattapuram, MD, Co-Chair

John M. Lewin, MD, FACR

Richard Duszak Jr., MD, FACR

Linda Moy, MD

Samuel A. Einstein, PhD

Mary S. Newell, MD, FACR

Phoebe Freer, MD

Haydee Ojeda-Fournier, MD

Catherine S. Giess, MD

Karla A. Sepulveda, MD, FACR

Amy Kotsenas, MD, FACR

Dana H. Smetherman, MD, MPH, MBA, FACR

David B. Larson, MD, MBA

Roland Wong, ScM

Paul A. Larson, MD, FACR

Karen S. Zheng, MD

Carol H. Lee, MD, FACR

## **REFERENCES**

1. Jeffries DO, Dossett LA, Jorns JM. Localization for Breast Surgery: The Next Generation. Arch Pathol Lab Med 2017;141:1324-29.
2. Food and Drug Administration. Mammography quality standards; final rule. Available at: <https://www.fda.gov/Radiation-EmittingProducts/MammographyQualityStandardsActandProgram/Regulations/ucm110906.htm>. Accessed January 22, 2019.
3. American College of Radiology. ACR practice parameter for the performance of stereotactic-guided breast interventional procedures Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Stereo-Breast.pdf>. Accessed January 22, 2019.
4. American College of Radiology. ACR practice parameter for the performance of ultrasound-guided percutaneous breast interventional procedures Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-GuidedBreast.pdf?la=en>. Accessed January 22, 2019.
5. American College of Radiology. ACR practice parameter for the performance of a breast ultrasound examination Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-Breast.pdf>. Accessed January 22, 2019.
6. American College of Radiology. ACR practice parameter for the performance of contrast-enhanced magnetic resonance imaging (MRI) of the breast Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Contrast-Breast.pdf>. Accessed January 22, 2019.
7. American College of Radiology. ACR practice parameter for the performance of magnetic resonance imaging-guided breast interventional procedures Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Guided-Breast.pdf>. Accessed January 22, 2019.
8. American College of Radiology. ACR practice parameter for the performance of screening and diagnostic mammography Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Screen-Diag-Mammo.pdf>. Accessed January 22, 2019.



9. Kirstein LJ, Rafferty E, Specht MC, et al. Outcomes of multiple wire localization for larger breast cancers: when can mastectomy be avoided? *J Am Coll Surg* 2008;207:342-6.
10. Liberman L, Kaplan J, Van Zee KJ, et al. Bracketing wires for preoperative breast needle localization. *AJR Am J Roentgenol* 2001;177:565-72.
11. Sharek D, Zuley ML, Zhang JY, Soran A, Ahrendt GM, Ganott MA. Radioactive seed localization versus wire localization for lumpectomies: a comparison of outcomes. *AJR Am J Roentgenol* 2015;204:872-7.
12. Cheang E, Ha R, Thornton CM, Mango VL. Innovations in image-guided preoperative breast lesion localization. *Br J Radiol* 2018;91:20170740.
13. Hughes JH, Mason MC, Gray RJ, et al. A multi-site validation trial of radioactive seed localization as an alternative to wire localization. *Breast J* 2008;14:153-7.
14. Jakub JW, Gray RJ, Degnim AC, Boughey JC, Gardner M, Cox CE. Current status of radioactive seed for localization of non palpable breast lesions. *Am J Surg* 2010;199:522-8.
15. McGhan LJ, McKeever SC, Pockaj BA, et al. Radioactive seed localization for nonpalpable breast lesions: review of 1,000 consecutive procedures at a single institution. *Ann Surg Oncol* 2011;18:3096-101.
16. Alderliesten T, Loo CE, Pengel KE, Rutgers EJ, Gilhuijs KG, Vrancken Peeters MJ. Radioactive seed localization of breast lesions: an adequate localization method without seed migration. *Breast J* 2011;17:594-601.
17. Dryden MJ, Dogan BE, Fox P, et al. Imaging Factors That Influence Surgical Margins After Preoperative 125I Radioactive Seed Localization of Breast Lesions: Comparison With Wire Localization. *AJR Am J Roentgenol* 2016;206:1112-8.
18. Goudreau SH, Joseph JP, Seiler SJ. Preoperative Radioactive Seed Localization for Nonpalpable Breast Lesions: Technique, Pitfalls, and Solutions. *Radiographics* 2015;35:1319-34.
19. Hayes MK. Update on Preoperative Breast Localization. *Radiol Clin North Am* 2017;55:591-603.
20. Cox CE, Russell S, Prowler V, et al. A Prospective, Single Arm, Multi-site, Clinical Evaluation of a Nonradioactive Surgical Guidance Technology for the Location of Nonpalpable Breast Lesions during Excision. *Ann Surg Oncol* 2016;23:3168-74.
21. Mango V, Ha R, Gomberawalla A, Wynn R, Feldman S. Evaluation of the SAVI SCOUT Surgical Guidance System for Localization and Excision of Nonpalpable Breast Lesions: A Feasibility Study. *AJR Am J Roentgenol* 2016;207:W69-W72.
22. Harvey JR, Lim Y, Murphy J, et al. Safety and feasibility of breast lesion localization using magnetic seeds (Magseed): a multi-centre, open-label cohort study. *Breast Cancer Res Treat* 2018;169:531-36.
23. Price ER, Khoury AL, Esserman LJ, Joe BN, Alvarado MD. Initial Clinical Experience With an Inducible Magnetic Seed System for Preoperative Breast Lesion Localization. *AJR Am J Roentgenol* 2018;210:913-17.
24. Lamb LR, Bahl M, Specht MC, D'Alessandro HA, Lehman CD. Evaluation of a Nonradioactive Magnetic Marker Wireless Localization Program. *AJR Am J Roentgenol* 2018;211:940-45.
25. Kapoor MM, Patel MM, Scoggins ME. The Wire and Beyond: Recent Advances in Breast Imaging Preoperative Needle Localization. *Radiographics* 2019;39:1886-906.
26. Graham RA, Homer MJ, Sigler CJ, et al. The efficacy of specimen radiography in evaluating the surgical margins of impalpable breast carcinoma. *AJR Am J Roentgenol* 1994;162:33-6.
27. Lee CH, Carter D. Detecting residual tumor after excisional biopsy of impalpable breast carcinoma: efficacy of comparing preoperative mammograms with radiographs of the biopsy specimen. *AJR Am J Roentgenol* 1995;164:81-6.
28. Mesurolle B, El-Khoury M, Hori D, et al. Sonography of postexcision specimens of nonpalpable breast lesions: value, limitations, and description of a method. *AJR Am J Roentgenol* 2006;186:1014-24.
29. Rebner M, Pennes DR, Baker DE, Adler DD, Boyd P. Two-view specimen radiography in surgical biopsy of nonpalpable breast masses. *AJR Am J Roentgenol* 1987;149:283-5.
30. Goldfeder S, Davis D, Cullinan J. Breast specimen radiography: can it predict margin status of excised breast carcinoma? *Acad Radiol* 2006;13:1453-9.
31. McCormick JT, Keleher AJ, Tikhomirov VB, Budway RJ, Caushaj PF. Analysis of the use of specimen mammography in breast conservation therapy. *Am J Surg* 2004;188:433-6.
32. Caudle AS, Yang WT, Mittendorf EA, et al. Selective surgical localization of axillary lymph nodes containing metastases in patients with breast cancer: a prospective feasibility trial. *JAMA Surg* 2015;150:137-43.
33. Shin K, Caudle AS, Kuerer HM, et al. Radiologic Mapping for Targeted Axillary Dissection: Needle Biopsy to Excision. *AJR Am J Roentgenol* 2016;207:1372-79.

Adopted 2021 (Resolution 28)  
Amended 2023 (Resolution 2c)  
College of Radiology. ACR practice parameter for communication of diagnostic imaging findings  
Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CommunicationDiag.pdf?la=en>.  
Accessed January 22, 2019.

\*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised, or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

#### Development Chronology for this Practice Parameter

2021 (Resolution 28)

Amended 2023 (Resolution 2c)