ACR-ASSR-SPR-SSR PRACTICE PARAMETER FOR THE PERFORMANCE OF SPINE RADIOGRAPHY

The American College of Radiology, with more than 40,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care 1. For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question. The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

1 lowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing, 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the "ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures (Revised 2008)" sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, Stanley v. McCarver, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

I. INTRODUCTION

This practice parameter was revised collaboratively by the American College of Radiology (ACR), the American Society of Spine Radiology (ASSR), the Society for Pediatric Radiology (SPR), and the Society of Skeletal Radiology (SSR).

Radiography of the spine is a proven and useful procedure for evaluating the vertebrae, disk spaces, facet and uncovertebral joints, neural foramina, and paravertebral soft tissues. This practice parameter outlines the principles for performing high-quality radiography of the cervical, thoracic, lumbar, sacral, and coccygeal spine and related osseous and soft-tissue structures to the extent they are visualized with radiography.

In many circumstances, especially when there is significant risk for spine injury, computed tomography (CT) or magnetic resonance imaging (MRI) is the initial imaging modality [1].

In patients with a clinical suspicion for spinal cord injury or compromise as well as ligamentous injuries, particularly in the cervical spine, MRI is preferred over CT and radiography.

The goal of these radiographic examinations is to identify or exclude anatomic abnormalities or disease processes of the spine and related tissues. The examinations should be performed with the minimum radiation exposure [2,3] necessary to produce a diagnostic study.

All radiographic examinations should be performed in accordance with the <u>ACR-AAPM-SIIM-SPR Practice</u> <u>Parameter for Digital Radiography</u> [4].

II. INDICATIONS

Indications include, but are not limited to, the evaluation of the spine for [5-26]:

- 1. Spinal Trauma and Fracture
 - a. Pain
 - b. Neurologic symptoms
 - c. Instability
 - d. Limitation of motion
 - e. Nontraumatic (insufficiency and stress) fractures
 - f. Pathologic fractures
- 2. Spinal arthropathy
 - a. Degenerative arthropathy
 - b. Inflammatory arthropathy
 - c. Neuropathic arthropathy
 - d. Crystal-induced arthropathy
- 3. Spinal infection
 - a. Discitis
 - b. Osteomyelitis
- 4. Spinal neoplasms and tumor-like conditions
 - a. Primary bone and soft tissue tumors
 - b. Metastatic disease
 - c. Multiple myeloma
 - d. Paget's disease
 - e. Aneurysmal bone cysts
 - f. Sarcoidosis, Langerhans cell histiocytosis, and other granulomatous conditions
- 5. Metabolic disorders
 - a. Osteoporosis
 - b. Renal osteodystrophy and complications related to dialysis
- 6. Congenital and acquired disorders
 - a. Scoliosis, kyphosis, flatback, truncal imbalance
 - b. Spondylolysis and spondylolisthesis
 - c. Congenital anomalies
 - d. Basilar invagination
- 7. Surgical and preprocedure planning, including intraoperative localizing images
- 8. Postoperative and postprocedural evaluation

For the pregnant or potentially pregnant patient, see the <u>ACR-SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation</u> [27].

III. QUALIFICATIONS AND RESPONSIBILITIES

See the ACR-AAPM-SIIM-SPR Practice Parameter for Digital Radiography [4].

IV. SPECIFICATIONS OF THE EXAMINATION

The written or electronic request for a Spine Radiograph should provide sufficient information to demonstrate the medical necessity of the examination and allow for its proper performance and interpretation.

Documentation that satisfies medical necessity includes 1) signs and symptoms and/or 2) relevant history (including known diagnoses). Additional information regarding the specific reason for the examination or a provisional diagnosis would be helpful and may at times be needed to allow for the proper performance and interpretation of the examination.

The request for the examination must be originated by a physician or other appropriately licensed health care provider. The accompanying clinical information should be provided by a physician or other appropriately licensed health care provider familiar with the patient's clinical problem or question and consistent with the state's scope of practice requirements. (ACR Resolution 35 adopted in 2006 – revised in 2016, Resolution 12-b)

This section discusses radiographic evaluation of the spine. Spinal radiography should be performed with appropriate collimation. Further imaging examinations may be indicated based on the clinical assessment and/or evaluation of the radiographs.

IV. SPECIFICATIONS OF THE EXAMINATION

A. Cervical Spine Examination

1. Adults [22,23,26,28-31]

- a. Routine examination consists of anteroposterior (AP) and lateral views. More limited examinations may be performed for specific indications. A swimmer's lateral view should be performed if necessary to assess the lower cervical segments and cervicothoracic junction alignment.
- b. In patients who have had trauma, and for whom cervical spine CT is nondiagnostic or otherwise unavailable, the entire cervical spine from the craniocervical junction to at least the superior end plate of T1 should be performed to assess for multiple fractures or associated traumatic listhesis [31]. Upright views are preferred but may not be possible if the patient's condition does not permit.
- c. In some clinical circumstances, additional evaluation may include some or all of the following: open mouth view (for assessment of dens and atlantoaxial association), closed mouth odontoid AP view (Fuchs view), oblique views (for assessment of the neural foramina), pillar views (for assessment of the facets), and flexion and extension lateral views (for assessment of cervical instability).
- d. If the patient has limited cervical range of motion on physical examination, flexion and extension radiographs may be inadequate to exclude instability and MRI should be obtained.
- e. If a cervical spine collar is present, it is the responsibility of the referring physician or referring physician's designee to remove the cervical spine collar and replace as appropriate.

2. Children [1,8,19,20,23-25,32-39]

- a. Routine examination includes AP and lateral views. Lateral radiographs should be obtained in true lateral position with the neck in extension if possible, and preferably during inspiration. Some pediatric centers omit the frontal view.
- b. Oblique views are not recommended due to the added radiation and low diagnostic yield.
- c. Flexion and extension lateral views are often not possible in younger children but may be useful to assess for ligament laxity in older children [40].
- d. Odontoid views are difficult to acquire in children younger than 5 years because of their short necks and imposition of the mandible on the spine and are not recommended [23].
- e. Cervical spine injury in young children (younger than 9 years old) [41] most commonly occurs from the occiput through C3 and has a propensity for ligamentous or cartilaginous rather than osseous injury. Therefore, normal cervical spine radiographs do not exclude ligamentous or spinal cord injury [24 25]
- f. In older children with chronic cervical instability (especially those with Down syndrome), lateral radiographs of the cervical spine centered at the craniocervical junction are taken in 3 positions: active flexion, active extension, and the standard neutral view [8,33-36].
- g. In the event additional imaging is needed, MRI, or occasionally CT, may be considered [12,29,42].

Radiation dose, magnetic safety, and potential sedation needs should be considered if these alternate imaging modalities are used.

IV. SPECIFICATIONS OF THE EXAMINATION

B. Thoracic Spine Examination

1. Adults

- a. A standard routine examination includes AP and lateral views. Lower cervical or upper lumbar anatomy should be visualized to assure accurate numbering of thoracic levels.
- b. Additional evaluation may be needed in some clinical circumstances and may include some or all of the following: swimmer's lateral view of the upper thoracic region, oblique views, flexion-extension lateral views, lateral bending views, and coned view of the thoracolumbar junction.

2. Children

- a. Routine examination includes AP and lateral views.
- b. Additional views may be obtained for specific clinical indications [16].

IV. SPECIFICATIONS OF THE EXAMINATION

C. Lumbosacral Spine Examination in Adults and Children

1. Adults

- a. A Standard examination includes AP and lateral views. Some may choose a posterior/anterior (PA) view instead of an AP view to reduce radiation dosage.
- b. In many adults and occasionally in older children, additional evaluation may be needed and may include some or all of the following: Both oblique views, coned lateral view of the lumbosacral junction, angled AP view of the lumbosacral junction, and upright flexion and extension lateral views may be particularly helpful to assess for abnormal motion.
- c. he upper part of the sacrum is included in the standard lumbosacral examination. When a more complete evaluation of the sacrum, coccyx, or sacroiliac joints is needed, a cephalad-angled AP (Ferguson) view of the sacrum and bilateral oblique/sacroiliac views may be obtained [21,43]. In select patients, dynamic coccygeal views or lateral seated position radiographs may demonstrate hypermobility or ligament laxity [44].

2. Children

- a. Standard examination includes AP and lateral views. A PA view may be used to reduce radiation dose.
- b. Oblique views are not recommended because of the added radiation and low diagnostic yield.
- c. Additional evaluation may be obtained for specific clinical indications.

IV. SPECIFICATIONS OF THE EXAMINATION

D. Scoliosis/Spine Deformity Examination in Adults and Children

1. Adults

- a. Erect PA (or AP) views of the entire thoracolumbar spine should be obtained, either with a single long computed radiography (CR) or screen-film cassette or with separate computed radiography, digital radiography, or screen-film captures physically or electronically stitched together. Some centers may use biplanar low-dose radiography to avoid stitching error and reduce whole body dose [45,46], which is becoming standard of care.
- b. Additional evaluation may be obtained for operative planning or post-operative evaluation.

2. Children

a. See the ACR–SPR–SSR Practice Parameter for the Performance of Radiography for Scoliosis in Children [47].

IV. SPECIFICATIONS OF THE EXAMINATION

E. Examination of Infants

- 1. Neonates and infants are usually evaluated with ultrasound [48,49] (see the ACR-AIUM-SPR-SRU Practice Parameter for the Performance of an Ultrasound Examination of the Neonatal and Infant Spine) [50] or MRI if congenital abnormality or trauma is highly suspected clinically or based on other imaging [51].
- 2. Interpretation of cervical spine radiography is difficult in infants because of epiphyseal variants, incomplete ossification of synchondroses including the apex of the odontoid, normal ligamentous laxity resulting in pseudosubluxation of C2 on C3, and the propensity of ligamentous rather than osseous injury. Normal lack of ossification of the anterior arch of C1 precludes radiographic evaluation of the atlantodental interval. MRI should be considered if there is concern for cervical spine injury.
- 3. Frontal and lateral views of the cervical spine, and combined frontal and lateral views of the thoracic and lumbar spine may be performed. These views are most frequently used in the setting of a skeletal survey for nonaccidental trauma or in the evaluation of skeletal dysplasia or congenital vertebral anomalies.

IV. SPECIFICATIONS OF THE EXAMINATION

F. Limited Examinations

- 1. For some clinical indications (eg, intraoperative or postsurgical follow-up), a limited examination of the area of clinical concern may provide sufficient information.
- 2. The goal is to limit patient radiation exposure.

V. DOCUMENTATION

Reporting should be in accordance with the <u>ACR Practice Parameter for Communication of Diagnostic Imaging Findings</u> [52].

VI. EQUIPMENT SPECIFICATIONS

See the <u>ACR-AAPM-SIIM-SPR Practice Parameter for Digital Radiography</u> [4].

Equipment performance monitoring should be in accordance with the <u>ACR-AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Radiographic Equipment [40]</u>. Radiographic Quality Control in Adults and Children

- 1. Examinations of the spine should completely demonstrate the designated portion(s) of the spine or the levels of clinical interest in a limited examination.
- 2. Images not of diagnostic quality should be repeated.
- 3. Each image should be permanently labeled in accordance with the <u>ACR-AAPM-SIIM-SPR Practice</u> <u>Parameter for Digital Radiography</u> [4].

VII. RADIATION SAFETY IN IMAGING

Radiologists, medical physicists, non-physician radiology providers, radiologic technologists, and all supervising physicians have a responsibility for safety in the workplace by keeping radiation exposure to staff, and to society as a whole, "as low as reasonably achievable" (ALARA) and to assure that radiation doses to individual patients are appropriate, taking into account the possible risk from radiation exposure and the diagnostic image quality necessary to achieve the clinical objective. All personnel who work with ionizing radiation must understand the key principles of occupational and public radiation protection (justification, optimization of protection, application of dose constraints and limits) and the principles of proper management of radiation dose to patients (justification, optimization including the use of dose reference levels). https://www-pub.iaea.org/MTCD/Publications/PDF/PUB1775_web.pdf

Nationally developed guidelines, such as the <u>ACR's Appropriateness Criteria</u>®, should be used to help choose the most appropriate imaging procedures to prevent unnecessary radiation exposure.

Facilities should have and adhere to policies and procedures that require ionizing radiation examination protocols (radiography, fluoroscopy, interventional radiology, CT) to vary according to diagnostic requirements and patient body habitus to optimize the relationship between appropriate radiation dose and adequate image quality. Automated dose reduction technologies

available on imaging equipment should be used, except when inappropriate for a specific exam. If such technology is not available, appropriate manual techniques should be used.

Additional information regarding patient radiation safety in imaging is available from the following websites – Image Gently® for children (www.imagegently.org) and Image Wisely® for adults (www.imagewisely.org). These advocacy and awareness campaigns provide free educational materials for all stakeholders involved in imaging (patients, technologists, referring providers, medical physicists, and radiologists).

Radiation exposures or other dose indices should be periodically measured by a Qualified Medical Physicist in accordance with the applicable ACR Technical Standards. Monitoring or regular review of dose indices from patient imaging should be performed by comparing the facility's dose information with national benchmarks, such as the ACR Dose Index Registry and relevant publications relying on its data, applicable ACR Practice Parameters, NCRP Report No. 172, Reference Levels and Achievable Doses in Medical and Dental Imaging: Recommendations for the United States or the Conference of Radiation Control Program Director's National Evaluation of X-ray Trends; 2006, 2009, amended 2013, revised 2023 (Res. 2d).

VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *Position Statement on Quality Control & Improvement, Safety, Infection Control, and Patient Education* on the ACR website (https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement).

ACKNOWLEDGEMENTS

This practice parameter was revised according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards) by the Committee on Body Imaging (Musculoskeletal) of the ACR Commission on Body Imaging and the Committee on Practice Parameters – Pediatric Radiology of the ACR Commission on Pediatric Radiology in collaboration with the ASSR, the SPR and the SSR.

https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards) by the Committee on Body Imaging (Musculoskeletal) of the ACR Commission on Pediatric Radiology in collaboration with the ASSR, the SPR and the SSR.

https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards) by the Committee on Body Imaging (Musculoskeletal) of the ACR Commission on Pediatric Radiology in collaboration with the ASSR, the SPR and the SSR.

https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards) by the Committee on Practice Parameters – Pediatric Radiology of the ACR Commission on Pediatric Radiology in collaboration with the ASSR, the SPR and the SSR.

https://www.acr.org/Clinical-Resources/Parameters-and-Technical-Standards) by the Committee on Practice Parameters – Pediatric Radiology of the ACR Commission on Pediatric Radiology in collaboration with the ASSR, the SPR and the SPR and the Parameters – Pediatric Radiology in Collaboration with the ASSR and the Parameters – Pediatric

<u>ACR</u> <u>ASSR</u>

Mary K. Jesse, MD, Chair

Felix Gonzalez, MD

Jennifer A. Knight, MD

Vikas Agarwal, MD

Allison Grayev, MD

Troy Hutchins, MD

Vinil Shah, MD

<u>SPR</u> <u>SSR</u>

Kirsten Ecklund, MD Leah Davis, DO

Richard Jones, MD Padmaja Jonnalagadda, MD

Amisha Shah, MD

<u>Committee on Practice Parameters – Musculoskeletal Body Imaging</u>

(ACR Committee responsible for sponsoring the draft through the process)

Catherine C. Roberts, MD, Chair

Jeffrey M. Brody, MD, FACR

Suzanne S. Long, MD

Bethany U. Casagranda, DO

Kambiz Motamedi, MD

Felix Gonzalez, MD

Carlos A. Rivera, BSc

Elaine S. Gould, MD, FACR Aleksandr Rozenberg, MD

Mary K. Jesse, MD Naveen Subhas, MD

<u>Committee on Practice Parameters – Pediatric Radiology</u>

(ACR Committee responsible for sponsoring the draft through the process)

Terry L. Levin, MD, FACR, Chair

Jane Sun Kim, MD

Jennifer A Knight, MD

Jesse Berman, MD

Jessica Kurian, MD

Tara M. Catanzano, MB, BCh Matthew P. Lungren, MD, MPH

Harris L. Cohen, MD, FACR Helen R. Nadel, MD
Kassa Darge, MD, PhD Erica Poletto, MD

Dorothy L. Gilbertson-Dahdal, MD Richard B. Towbin, MD, FACR

Lauren P. Golding, MD

Safwan S. Halabi, MD

Esben S. Vogelius, MD

Jason Higgins, DO

Andrew B. Rosenkrantz, MD, Chair, Commission on Body Imaging
Richard A. Barth, MD, FACR, Chair, Commission on Pediatric Radiology
David B. Larson, MD, MBA, Chair, Commission on Quality and Safety
Mary S. Newell, MD, FACR, Chair, Committee on Practice Parameters and Technical Standards

Comments Reconciliation Committee

Kurt Schoppe, MD– CSC Chair Padmaja Jonnalagadda, MD
Neil Lall, MD– CSC Co-Chair Jennifer A. Knight, MD

Vikas Agarwal, MD Amy L. Kotsenas, MD, FACR Hamza Alizai, MD David B. Larson, MD, MBA

Richard A. Barth, MD, FACR Paul Larson, MD

Nicholas M. Beckmann, MD

Terry L. Levin, MD, FACR

Rebecca Milman, Ph.D

Nirvikar Dahiya, MD, FAIUM, FSRU

Leah Davis, DO

Felix Gonzalez, MD

Terry L. Levin, MD, FACR

Rebecca Milman, Ph.D

Mary S. Newell, MD, FACR

Catherine C. Roberts, MD

Andrew B Rosenkrantz, MD

Felix Gonzalez, MD

Andrew B Rosenkrantz, MD

Allison Grayev, MD

Marla B. K. Sammer, MD

Troy Hutchins, MD Vinil Shah, MD

Mary K. Jesse, MD Jonelle M. Thomas, MD

REFERENCES

- **1.** Beckmann NM, West OC, Nunez D, Jr., et al. ACR Appropriateness Criteria® Suspected Spine Trauma. J Am Coll Radiol 2019;16:S264-S85.
- **2.** Almén AJ, Mattsson S. Dose distribution at radiographic examination of the spine in pediatric radiology. Spine (Phila Pa 1976). 1996 Mar 15;21(6):750-6.
- **3.** Gialousis GI, Yakoumakis EN, Papadopoulou DI, et al. Differences in effective dose and energy imparted estimation from PA-AP, RLAT-LLAT projections in pediatric full spine x-ray examination using the Monte Carlo technique. Phys Med Biol. 2006 Jan 21;51(2):287-97.
- **4.** American College of Radiology. ACR-AAPM-SIIM Practice Parameter for Digital Radiography. Available at https://gravitas.acr.org/PPTS/GetDocumentView?docld=135+&releaseId=2
- **5.** Hadley MN, Walters BC, Grabb PA, et al. Radiographic assessment of the cervical spine in asymptomatic trauma patients. Neurosurgery. 2002 Mar;50(3 Suppl):S30-5.

- **6.** Berry GE, Adams S, Harris MB, et al. Are plain radiographs of the spine necessary during evaluation after blunt trauma? Accuracy of screening torso computed tomography in thoracic/lumbar spine fracture diagnosis. J Trauma. 59(6):1410-3; discussion 1413, 2005 Dec.
- **7.** Brockmeyer D. Down syndrome and craniovertebral instability. Topic review and treatment recommendations. Pediatr Neurosurg. 1999 Aug;31(2):71-7.
- **8.** Freemyer B, Knopp R, Piche J, Wales L, Williams J. Comparison of five-view and three-view cervical spine series in the evaluation of patients with cervical trauma. Ann Emerg Med. 1989 Aug;18(8):818-21.
- **9.** Gale SC, Gracias VH, Reilly PM, Schwab CW. The inefficiency of plain radiography to evaluate the cervical spine after blunt trauma. Journal of Trauma-Injury Infection & Critical Care. 59(5):1121-5, 2005 Nov.
- **10.** Griffen MM, Frykberg ER, Kerwin AJ, et al. Radiographic clearance of blunt cervical spine injury: plain radiograph or computed tomography scan?. Journal of Trauma-Injury Infection & Critical Care. 55(2):222-6; discussion 226-7, 2003 Aug.
- **11.** Holmes JF, Akkinepalli R. Computed tomography versus plain radiography to screen for cervical spine injury: a meta-analysis. [Review] [21 refs]. Journal of Trauma-Injury Infection & Critical Care. 58(5):902-5, 2005 May.
- **12.** Insko EK, Gracias VH, Gupta R, Goettler CE, Gaieski DF, Dalinka MK. Utility of flexion and extension radiographs of the cervical spine in the acute evaluation of blunt trauma. Journal of Trauma-Injury Infection & Critical Care. 53(3):426-9, 2002 Sep.
- **13.** McNamara RM, Heine E, Esposito B. Cervical spine injury and radiography in alert, high-risk patients. J Emerg Med. 8(2):177-82, 1990 Mar-Apr.
- **14.** Mirvis SE, Diaconis JN, Chirico PA, Reiner BI, Joslyn JN, Militello P. Protocol-driven radiologic evaluation of suspected cervical spine injury: efficacy study. Radiology. 170(3 Pt 1):831-4, 1989 Mar.
- **15.** Mower WR, Hoffman JR, Pollack CV, et al. Use of plain radiography to screen for cervical spine injuries. Ann Emerg Med. 2001 Jul;38(1):1-7.
- **16.** Offerman SR, Holmes JF, Katzberg RX, Richards JR. Utility of supine oblique radiographs in detecting cervical spine injury. J Emerg Med. 2006 Feb;30(2):189-95.
- **17.** Pollack CV Jr, Hendey GW, Martin DR, Hoffman JR, Mower WR. Use of flexion-extension radiographs of the cervical spine in blunt trauma. Annals of Emergency Medicine. 38(1):8-11, 2001 Jul.
- **18.** Ralston ME, Chung K, Barnes PD, Emans JB, Schutzman SA. Role of flexion-extension radiographs in blunt pediatric cervical spine injury. Acad Emerg Med. 2001 Mar;8(3):237-45.
- **19.** Ralston ME, Ecklund K, Emans JB, Torrey SB, Bailey MC, Schutzman SA. Role of oblique radiographs in blunt pediatric cervical spine injury. [Review] [28 refs]. Pediatr Emerg Care. 19(2):68-72, 2003 Apr.
- **20.** Reinus WR, Strome G, Zwemer FL. Use of lumbosacral spine radiographs in a level II emergency department. AJR Am J Roentgenol. 1998 Feb;170(2):443-7.
- **21.** Ross SE, O'Malley KF, DeLong WG, Born CT, Schwab CW. Clinical predictors of unstable cervical spinal injury in multiply injured patients. Injury. 1992; 23(5):317-319.
- **22.** Schousboe JT, Rosen HR, Vokes TJ, et al. Prediction models of prevalent radiographic vertebral fractures among older women. J Clin Densitom 2014;17:378-85.
- **23.** Swischuk LE, John SD, Hendrick EP. Is the open-mouth odontoid view necessary in children under 5 years?. Pediatr Radiol. 2000 Mar;30(3):186-9.
- **24.** Platzer P, Jaindl M, Thalhammer G, et al. Cervical spine injuries in pediatric patients. J Trauma. 2007 Feb;62(2):389-96; discussion 394-6.
- **25.** Smart PJ, Hardy PJ, Buckley DM, et al. Cervical spine injuries to children under 11: should we use radiography more selectively in their initial assessment?. Emerg Med J. 2003 May;20(3):225-7.
- **26.** Michaleff ZA, Maher CG, Verhagen AP, Rebbeck T, Lin CW. Accuracy of the Canadian C-spine rule and NEXUS to screen for clinically important cervical spine injury in patients following blunt trauma: a systematic review. [Review]. CMAJ. 184(16):E867-76, 2012 Nov 06.
- **27.** American College of Radiology. ACR-SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation. Available at:
- https://gravitas.acr.org/PPTS/GetDocumentView?docId=23+&releaseId=2.
- **28.** Dreizin D, Letzing M, Sliker CW, et al. Multidetector CT of blunt cervical spine trauma in adults. [Review]. Radiographics. 34(7):1842-65, 2014 Nov-Dec.
- **29.** McCulloch PT, France J, Jones DL, et al. Helical computed tomography alone compared with plain radiographs with adjunct computed tomography to evaluate the cervical spine after high-energy trauma. J Bone Joint Surg Am. 2005 Nov;87(11):2388-94.
- **30.** Inaba K, Nosanov L, Menaker J, et al. Prospective derivation of a clinical decision rule for thoracolumbar spine evaluation after blunt trauma: An American Association for the Surgery of Trauma Multi-Institutional Trials Group

- Study. J Trauma Acute Care Surg. 78(3):459-65; discussion 465-7, 2015 Mar.
- **31.** Nelson DW, Martin MJ, Martin ND, Beekley A. Evaluation of the risk of noncontiguous fractures of the spine in blunt trauma. The Journal of Trauma and Acute Care Surgery. 75(1):135-9, 2013 Jul.
- **32.** Rozzelle CJ, Aarabi B, Dhall SS, et al. Management of pediatric cervical spine and spinal cord injuries. Neurosurgery. 2013 Mar;72 Suppl 2():205-26.
- **33.** American Academy of Pediatrics. Committee on Sports Medicine. Atlantoaxial instability in Down Syndrome. Pediatrics. 1984 Jul;74(1):152-4.
- **34.** Atlantoaxial instability in Down syndrome: subject review. American Academy of Pediatrics Committee on Sports Medicine and Fitness. Pediatrics. 1995 Jul;96(1 Pt 1):151-4.
- 35. Bull MJ, Health supervision for children with Down syndrome. Pediatrics. 2011 Aug;128(2):393-406.
- **36.** Wyckoff AS. AAP updates guidance on caring for children with Down syndrome. AAP News 2011;32:10.
- **37.** McAllister AS, Nagaraj U, Radhakrishnan R. Emergent Imaging of Pediatric Cervical Spine Trauma. Radiographics. 2019;39(4):1126-1142.
- **38.** Arbuthnot M, Mooney DP. The sensitivity and negative predictive value of a pediatric cervical spine clearance algorithm that minimizes computerized tomography. Journal of Pediatric Surgery. 52(1):130-135, 2017 Jan.
- **39.** Cui LW, Probst MA, Hoffman JR, Mower WR. Sensitivity of plain radiography for pediatric cervical spine injury. Emergency Radiology. 23(5):443-8, 2016 Oct.
- **40.** Tat ST, Mejia MJ, Freishtat RJ. Imaging, clearance, and controversies in pediatric cervical spine trauma. [Review]. Pediatr Emerg Care. 30(12):911-5; quiz 916-8, 2014 Dec.
- **41.** Beckmann NM, Chinapuvvula NR, Zhang X, West OC. Epidemiology and Imaging Classification of Pediatric Cervical Spine Injuries: 12-Year Experience at a Level 1 Trauma Center. AJR. American Journal of Roentgenology. 214(6):1359-1368, 2020 06.
- **42.** Waddell VA, Connelly S. Decreasing Radiation Exposure in Pediatric Trauma Related to Cervical Spine Clearance: A Quality Improvement Project. J Trauma Nurs. 2018;25(1):38-44.
- **43.** Battistone MJ, Manaster BJ, Reda DJ, Clegg DO. Radiographic diagnosis of sacroiliitis--are sacroiliac views really better? J Rheumatol. 1998;25(12):2395-2401.
- **44.** Maigne JY, Tamalet B. Standardized radiologic protocol for the study of common coccygodynia and characteristics of the lesions observed in the sitting position. Clinical elements differentiating luxation, hypermobility, and normal mobility. Spine (Phila Pa 1976). 1996 Nov 15;21(22):2588-93.
- **45.** Celestre PC, Dimar JR, Glassman SD. Spinopelvic Parameters: Lumbar Lordosis, Pelvic Incidence, Pelvic Tilt, and Sacral Slope: What Does a Spine Surgeon Need to Know to Plan a Lumbar Deformity Correction?. Neurosurg Clin N Am. 2018 Jul;29(3):S1042-3680(18)30837-4.
- **46.** Melhem E, Assi A, El Rachkidi R, Ghanem I. EOS(®) biplanar X-ray imaging: concept, developments, benefits, and limitations. J Child Orthop. 2016 Feb;10(1):1-14.
- **47.** American College of Radiology. ACR-SPR-SSR Practice Parameter for the Performance of Radiography for Scoliosis in Children. Available at: https://gravitas.acr.org/PPTS/GetDocumentView?docId=44+&releaseId=2
- **48.** Lowe LH, Johanek AJ, Moore CW. Sonography of the neonatal spine: part 1, Normal anatomy, imaging pitfalls, and variations that may simulate disorders. AJR Am J Roentgenol. 2007;188(3):733-738.
- **49.** Fotter R, Sorantin E, Schneider U, Ranner G, Fast C, Schober P. Ultrasound diagnosis of birth-related spinal cord trauma: neonatal diagnosis and follow-up and correlation with MRI. Pediatr Radiol. 1994;24(4):241-4.
- **50.** American College of Radiology. ACR-AIUM-SPR-SRU Practice Parameter for the Performance of an Ultrasound Examination of the Neonatal and Infant Spine. Available at
- https://gravitas.acr.org/PPTS/GetDocumentView?docId=191+&releaseId=2
- **51.** Rossi A, Martinetti C, Morana G, Severino M, Tortora D. Diagnostic Approach to Pediatric Spine Disorders. Magn Reson Imaging Clin N Am. 2016 Aug;24(3):S1064-9689(16)30015-0.
- **52.** American College of Radiology. ACR Practice Parameter for Communication of Diagnostic Imaging Findings. Available at https://gravitas.acr.org/PPTS/GetDocumentView?docId=74+&releaseId=2
- *Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

Amended 2009 (Resolution 11)
Revised 2012 (Resolution 32)
Amended 2014 (Resolution 39)
Revised 2017 (Resolution 7)
Revised 2022 (Resolution 37)
Amended 2023 (Resolution 2c, 2d)