# ACR-SIR-SPR PRACTICE PARAMETER FOR THE PERFORMANCE OF INFERIOR VENA CAVA (IVC) FILTER PLACEMENT FOR THE PREVENTION OF PULMONARY EMBOLISM

The American College of Radiology, with more than 40,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

#### **PREAMBLE**

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care  $\frac{1}{2}$ . For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question. The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

1 lowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing, 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the "ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures (Revised 2008)" sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, Stanley v. McCarver, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

#### I. INTRODUCTION

This practice parameter was revised collaboratively by the ACR, the Society of Interventional Radiology (SIR), and the Society for Pediatric Radiology (SPR).

"Pulmonary embolism (PE) continues to be a major cause of morbidity and mortality in the United States. Estimates of the incidence of nonfatal PE range from 400,000 to 630,000 cases per year and from 50,000 to

200,000 fatalities per year are directly attributable to PE" [1].

Historically, inferior vena cava (IVC) filters were designed for permanent placement, but most commercially available IVC filter are now retrievable. "The (retrievable) filters can be left in place as a permanent implant but can also be removed when the indication for filter placement resolves (for detailed information regarding each of these filters, the reader can refer to several reviews [1-20])." Device selection is dependent on many factors, including, but not limited to, clinical indication, anatomy of the iliocaval veins and available venous access sites, device-specific outcomes data, magnetic resonance imaging (MRI) compatibility, ease of placement, and experience of the operator [1].

"The IVC should be assessed with imaging prior to placement of a filter. The current preferred method is by vena cavography" [1]. Intravascular ultrasound (IVUS) has been utilized as an adjunct or primary tool for IVC assessment in certain situations. However, the use of IVUS is beyond the scope of this document [21-24]. The length and diameter of the IVC should be assessed, the location and number of renal veins determined, IVC variants defined (eg, duplication), and intrinsic IVC disease, such as preexisting thrombus or extrinsic compression, should be assessed [1,25]. If available, prior imaging studies (such as contrast-enhanced computed tomography [CT] or MRI of the abdomen and pelvis) should be reviewed to evaluate the anatomy of the IVC (size, patency, and anatomical variants) and to determine intrinsic IVC disease and extrinsic compression [1]. The recommended location for filter placement for preventing PE from a lower extremity or pelvic venous source is the infrarenal IVC [1,25]. It is a generally accepted belief that the apex or superior aspect of any filtration device should be at or immediately inferior to the level of the renal veins, according to the manufacturer's recommendations [1,25]. In specific clinical circumstances, other target locations, such as the suprarenal IVC, iliac veins, or both sides of a duplicated IVC, may be appropriate [1].

When a retrievable filter is placed, the patient should be clinically reassessed periodically to weigh the benefits of continued filtration (need for PE prophylaxis) against the associated risks (eg, recurrent deep venous thrombosis [DVT], IVC thrombosis, symptomatic penetration, or mechanical failure) and uncertainties (given limited data on long-term mechanical stability and integrity of some devices). The current recommendation is to consider removal of the IVC filter as soon as mechanical protection against PE is no longer needed [26]. Filters placed with the intent of subsequent retrieval may be left in place permanently for any of several reasons (eg, continuing need for filtration, thrombus on the filter, or inability to retrieve the filter). Data for the feasibility of filter retrieval vary widely among devices and centers.

These practice parameters are intended to be used in quality improvement programs to assess percutaneous interruption of the IVC to prevent PE. The most important aspects of care are 1) patient selection, 2) performing the procedure, and 3) patient monitoring. The outcome measures or indicators for these processes are indications, success rates, and complication rates. Outcome measures are assigned threshold levels.

## **II. INDICATIONS AND CONTRAINDICATIONS [1,25,27-38]**

Indications include, but are not limited to, the following:

- A. Therapeutic (Documented Thromboembolic Disease)
  - 1. Patients with evidence of pulmonary embolus or DVT involving the IVC, iliac, or femoral-popliteal veins and 1 or more of the following:
    - a. An absolute contraindication to anticoagulation [39]
    - b. Failure of anticoagulation [27]
      - i. Recurrent symptomatic PE despite therapeutic anticoagulation
      - ii. Propagation/progression of DVT on therapeutic anticoagulation
  - 2. Perioperative patients with a history of prior venous thromboembolism (VTE) for whom anticoagulation must be interrupted [27,39]
  - 3. Prior to pulmonary thromboendarterectomy in patients with chronic pulmonary thromboembolic pulmonary hypertension
  - 4. During or after surgical or endovascular thrombolysis/thrombectomy for massive PE [40-42]
  - 5. PE in the presence of severe cardiopulmonary disease (eg, cor pulmonale with pulmonary hypertension, congestive heart failure, PE requiring ventilator support) [40,43,44]
- B. Prophylactic (No Current Thromboembolic Disease)
  - There is a limited role for prophylactic placement of IVC filters in the absence of acute thromboembolic disease. Multidisciplinary discussion of the appropriateness of an IVC filter should be performed with a

focus on use in patients with increased mortality risk from recurrent VTE and inability to continue anticoagulation [39].

## C. Suprarenal Filter Placement [1,37]

Suprarenal caval filter placement may be considered when any of the following situations exist in addition to the indications listed above:

- 1. Presence of IVC thrombus precluding placement of a filter in the infrarenal IVC
- 2. Filter placement during pregnancy
- 3. Thrombus extending above a previously placed infrarenal filter or inability to safely or appropriately place the IVC filter in an infrarenal location
- 4. Gonadal vein thrombosis [45]
- 5. Anatomic variants: duplication of the IVC, short length of infrarenal IVC
- 6. Significant extrinsic compression of the infrarenal IVC
- 7. Intrinsic narrowing of the infrarenal IVC
- 8. Patients with an intra-abdominal or pelvic mass who will undergo surgery and in whom operative IVC mobilization is contemplated

## D. Filters Placed for Temporary Use and Possible Future Retrieval

The indications for filter placement remain the same as described above, except that the mechanical protection for PE is required for the short term because of its transient inability to anticoagulate a patient [1,32].

The use of retrievable filters should also be considered in pediatric and young adult patients because the long-term effects and durability of permanent IVC filters are not precisely known. Currently, there are no filters specifically designed for use in children [1]. The safety of vena cava filters in children is unknown [1,25,46,47].

"The threshold for these indications is 95%. When <95% of procedures are performed for these indications, the process of patient selection should be reviewed according to institutional policy" [1].

# E. Relative Contraindications

1. Bacteremia or untreated infection [1,25]

"Clinical judgment should be applied in these situations, weighing the theoretical risk of implant infection versus the risk of PE" [1].

## III. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

#### A. Physician

Examinations must be performed under the supervision of and interpreted by a physician who has the following qualifications:

- Certification in Radiology, Diagnostic Radiology or Interventional Radiology/Diagnostic Radiology (IR/DR), by the American Board of Radiology, the American Osteopathic Board of Radiology, the Royal College of Physicians and Surgeons of Canada (RCPSC), or the Collège des Médecins du Québec and has performed (with supervision) a sufficient number of IVC filter procedures to demonstrate competency as attested by the supervising physician(s).
- 2. Completion of a residency program approved by the Accreditation Council for Graduate Medical Education (ACGME), the Royal College of Physicians and Surgeons of Canada (RCPSC), the Collège des Médecins du Québec, or the American Osteopathic Association (AOA) and has performed (with supervision) a sufficient number of IVC filter procedures to demonstrate competency as attested by the supervising physician(s).

or

3. Physicians whose residency or fellowship training did not include the above may still be considered qualified to perform IVC filter procedures provided that the following can be demonstrated:

The physician must have endovascular procedural experience, during which the physician was supervised and during which the physician performed percutaneous vascular procedures, including IVC filter placement procedures with outcomes within the quality improvement thresholds of this document.

Physicians meeting any of the qualifications in 1, 2, and 3 above must also have written substantiation that they are familiar with all of the following:

- a. Indications and contraindications for the procedure
- b. Periprocedural and intraprocedural assessment, monitoring, and management of the patient and potential complications
- c. Where applicable, pharmacology of moderate-sedation medications and recognition and treatment of adverse reactions and complications
- d. Appropriate use and operation of fluoroscopic and radiographic equipment, mechanical injectors, digital image capture devices, and electronic imaging systems
- e. Where applicable, principles of radiation protection, hazards of radiation, and radiation monitoring requirements as they apply to both patients and personnel
- f. Where applicable, pharmacology of contrast agents and recognition and treatment of potential adverse reactions
- g. Percutaneous needle and catheter introduction techniques
- h. Technical aspects of performing the procedure, including the use of alternative catheter and guidewire systems, contrast materials, and filming sequences
- i. Anatomy, physiology, and pathophysiology of peripheral and central venous vasculature
- j. Postprocedure patient management, especially recognition and initial management of complications
- k. Postprocedure management of puncture sites
- I. Principles, operation, and imaging findings of vascular ultrasound, where applicable
  The written substantiation should come from the chief of interventional radiology, the chair of
  the department of radiology, or his or her designee at the institution in which the physician
  will be providing these services. Substantiation could also come from a prior institution in
  which the physician provided the services, but only at the discretion of the chair of the
  department of radiology or his or her designee who solicits the additional input.

## Maintenance of Competence

Physicians must perform a sufficient number of filter placement procedures to maintain their skills, with acceptable success and complication rates as presented in this document (see section X). Continued competence should depend on participation in a quality improvement program that monitors these rates.

# **Continuing Medical Education**

The physician's continuing education should be in accordance with the <u>ACR Practice</u> <u>Parameter for Continuing Medical Education (CME)</u> [48].

# B. Qualified Medical Physicist

For qualifications of the Qualified Medical Physicist, see the <u>ACR—AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Fluoroscopic Equipment</u> and the <u>ACR—AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Radiographic Equipment</u> [49,50].

# C. Non-Physician Radiology Provider (NPRP)

NPRPs are all Non-Physician Providers (eg, RRA, RPA, RA, PA, NP, ...) who assist with or participate in portions of the practice of a radiologist-led team (Radiologists = diagnostic, interventional, neurointerventional radiologists, radiation oncologists, and nuclear medicine physicians). The term "NPRP" does not include radiology, CT, US, NM MRI technologists, or radiation therapists who have specific training for radiology related tasks (eg, acquisition of images, operation of imaging and therapeutic equipment) that are not typically performed by radiologists.

The term 'radiologist-led team' is defined as a team supervised by a radiologist (eg, diagnostic,

interventional, neurointerventional radiologist, radiation oncologist, and nuclear medicine physician) and consists of additional healthcare providers including RRAs, PAs, NPs, and other personnel critical to the provision of the highest quality of healthcare to patients. (ACR Resolution 8, adopted 2020).

#### D. Radiologic Technologist

- 1. The technologist, together with the physician and nursing personnel, should be responsible for patient comfort and safety. The technologist should be able to prepare and position [1] the patient for the filter placement procedure. The technologist should obtain the imaging data in a manner prescribed by the supervising physician. If intravenous contrast material is to be administered, qualifications for technologists performing intravenous injection should be in compliance with current ACR policy [2] and existing operating procedures or manuals at the facility. The technologist should also perform the regular quality control testing of the equipment under the supervision of the physicist.
- 2. Technologists should be certified by the ARRT or have an unrestricted state license with documented training and experience in the imaging modality used for the imaging-guided procedure.

## E. Nursing Services

Nursing services, when deemed appropriate by the performing physician, are an integral part of the team for preprocedure and postprocedure patient management and education and are thus recommended in monitoring the patient during the procedure.

[1] The American College of Radiology approves of the practice of certified and/or licensed radiologic technologists performing fluoroscopy in a facility or department as a positioning or localizing procedure only, and then only if monitored by a supervising physician who is personally and immediately available\*. There must be a written policy or process for the positioning or localizing procedure that is approved by the medical director of the facility or department/service and that includes written authority or policies and processes for designating radiologic technologists who may perform such procedures. (ACR Resolution 26, 1987 – revised in 2007, Resolution 12-m)

\*For the purposes of this guideline, "personally and immediately available" is defined in manner of the "personal supervision" provision of CMS—a physician must be in attendance in the room during the performance of the procedure. Program Memorandum Carriers, DHHS, HCFA, Transmittal B-01-28, April 19, 2001.

[2] See the ACR-SPR Practice Parameter for the Use of Intravascular Contrast Media.

# IV. SPECIFICATIONS OF THE EXAMINATION [1]

"There are several technical requirements to ensure safe and successful filter placement procedures. These include adequate angiographic equipment and institutional facilities, physiologic monitoring equipment, and support personnel" [1].

#### A. Equipment and Facilities for Filter Placement

"The following are considered the minimum equipment requirements for performing vena cavagrams and filter placement. In planning facilities for IVC placement, equipment and facilities more advanced than those outlined (below) may be desired to produce higher-quality studies with reduced risk and time of study" [1]. At a minimum, the facility should include the following:

- 1. A high-resolution image receptor, preferably with a 28-cm to 40-cm field of view, and a digital imaging system with a minimum 1024-image matrix [1]. The imaging system should be capable of pulsed fluoroscopy at a rate of 7.5 frames per second.
  Digital angiographic systems are preferred as they allow for reduced volumes of contrast material and reduced examination times. Images are acquired and stored digitally on computerized storage media. Imaging and image recording must be consistent with the "as low as reasonably achievable" (ALARA) radiation safety guidelines. The use of cineradiography or small-field mobile image intensifiers is inappropriate for the routine recording of the vena cavagram and IVC placement because these methods cause an unacceptably high patient and operator radiation dose. Use of last image hold and pulsed fluoroscopy are recommended for dose reduction [1].
- 2. Adequate angiographic supplies, such as catheters, guidewires, needles, and introducer sheaths [1].

- 3. "An angiographic injector capable of varying injection volumes and rates with appropriate safety mechanisms to prevent overinjection" [1].
- 4. "An angiography suite that is large enough to allow easy transfer of the patient from the bed to the table and allow room for the procedure table, monitoring equipment, and other hardware, such as intravenous pumps, respirators, anesthesia equipment, and oxygen tanks. Ideally there should be adequate space for the operating team to work unencumbered on either side of the patient and for the circulation of other technical staff in the room without contaminating the sterile conditions" [1].
- 5. "An area within the institution appropriate for patient preparation prior to the procedure and for observation of patients after the procedure. This might be within the radiology department, a short-stay unit, a routine nursing unit, or a postanesthesia care unit. At this location, there should be personnel to provide care as outlined below (Patient Care), and there should be immediate access to emergency resuscitation equipment" [1].

## B. Physiologic Monitoring and Resuscitation Equipment

- "Equipment should be present in the procedure suite to allow for monitoring the patient's heart rate, cardiac rhythm, and blood pressure. For facilities using moderate sedation, a pulse oximeter monitor should be available" [1] (See the <u>ACR-SIR Practice Parameter for Minimal and/or Moderate Sedation/Analgesia</u> [51]).
- 2. "Appropriate emergency equipment and medications must be immediately available to treat adverse reactions associated with administered medications and/or procedural complications. The equipment should be maintained and medications inventoried for drug expiration dates on a regular basis. The equipment, medications, and other emergency support must also be appropriate for the range of ages and sizes in the patient population" [1].

## C. Support Personnel

- 1. "Radiologic technologists properly trained in the use of the angiographic equipment should assist in performing the procedure. They should demonstrate appropriate knowledge of patient positioning; angiographic image recording; angiographic contrast injectors; angiographic supplies, including IVC filters; and the physiologic monitoring equipment. Certification as a vascular and interventional radiologic technologist is one measure of appropriate training. The technologist should be trained in basic cardiopulmonary resuscitation and in the function of the resuscitation equipment" [1].
- 2. "If the patient does not receive sedation for the procedure, one of the staff assisting the procedure should be assigned to periodically assess the patient's status. In cases where moderate sedation is used in adults or children or the patient is critically ill, an experienced, licensed provider should be present, whose primary responsibility is monitoring the patient's vital signs, sedation state, and level of comfort or pain. This person should maintain a record of the patient's vital signs, the time and dose of medications given, and other pertinent information" [1] (See the <u>ACR-SIR Practice Parameter for Minimal and/or Moderate Sedation/Analgesia</u> [51]).

## D. Acute Care Support

"Although surgical or other emergency treatment is needed infrequently for serious complications after filter placement procedures, there should be prompt access to surgical and interventional equipment and to specialists familiar with the management of patients with complications in the unlikely event of a lifethreatening complication" [1].

#### E. Patient Care

For additional information see the <u>ACR–SIR–SNIS–SPR Practice Parameter for the Clinical Practice of Interventional Radiology</u> [52].

1. Preprocedure care [1]

For elective filter placement, the following should be documented:

- a. Clinically significant history, including indications for the procedure
- b. Clinically significant physical or diagnostic examination findings, including clinical or medical conditions that may necessitate specific care, such as preprocedure antibiotics and other measures

- c. Clinically indicated laboratory evaluation including, but not limited to, coagulation factors, creatinine, white blood cell count, and previously obtained cultures
- d. Preprocedure documentation should conform to the requirements of the <u>ACR-SIR-SPR</u>

  <u>Practice Parameter for the Reporting and Archiving of Interventional Radiology Procedures</u>

  [53]

Informed consent must be in compliance with all state laws and the <u>ACR-SIR-SPR Practice</u> <u>Parameter on Informed Consent for Image-Guided Procedures</u> [54].

"For emergency procedures, a note should be written summarizing the indication for the study, the pertinent history and physical findings, if available, and the proposed procedure" [1].

#### 2. Procedural care

- a. "Adherence to the Joint Commission's Universal Protocol for Preventing Wrong Site, Wrong Procedure, Wrong Person Surgery™ is required for procedures in nonoperating room settings, including bedside procedures" [1]. The organization should have processes and systems in place for reconciling differences in staff responses during the "time-out."
- b. "All patients should have cardiac monitoring continuously during the procedure, with intermittent blood pressure monitoring. A record of vital signs should be maintained" [1].
- c. "All patients should have intravenous access for the administration of fluids and medications as needed" [1].
- d. "If the patient is to receive sedation for the procedure, pulse oximetry should be used. A registered nurse or other appropriately trained personnel should be present, and their primary responsibility should be to monitor the patient. A record should be kept of medication doses and times of administration" [1]. See the <u>ACR-SIR Practice Parameter for Minimal and/or Moderate Sedation/Analgesia</u> [51].

#### 3. Postprocedure care

- a. "All patients should be on bed rest and observed in the initial postprocedure period. The length of this period of bed rest will depend on the site and size of the venotomy and the patient's medical condition" [1].
- b. "During the initial postprocedure period, skilled nurses or other appropriately trained personnel should periodically monitor the puncture site" [1].
- c. "Initial ambulation of the patient must be carefully supervised. The puncture site stability and independent patient function and mobility must be ensured" [1].
- d. "The operating physician or a qualified designee should evaluate the patient after the procedure, and these findings should be summarized in a progress note" [1]. If moderate sedation was administered prior to and during the procedure, complete recovery from sedation must be documented [1]. "The physician or designee should be available for continuing care during hospitalization and after discharge. The designee may be another physician or a nurse" [1]. See the <u>ACR-SIR Practice Parameter for Minimal and/or Moderate Sedation/Analgesia</u> [51].
- e. Requirement for an IVC filter retrieval program. Patients in whom a retrievable IVC filter is implanted should undergo clinical reassessment for the appropriateness and timing of filter removal during the first 3 to 6 months after placement (depending on the device-specific time window for retrieval). The operating physician should participate in this clinical assessment.

## F. Selection Criteria for Short-Term Observation

"The duration of postprocedure observation must be individualized. IVC filter placement can be performed on some patients with a short period of postprocedure observation (<6 hours) prior to discharge to home; others require overnight care. Short-term observation should only be considered when all the following conditions can be met" [1]:

1. "Those patients capable of independent ambulation prior to the procedure demonstrate stable independent ambulation after the procedure. Nonambulatory patients have adequate assistance after discharge to provide care as needed" [1].

- 2. "The patient is capable of following instructions and detecting changes in symptomatology. Alternatively, patients with impaired mental or neurologic status should have adequate assistance after discharge to provide care as needed" [1].
- 3. "The patient is provided with instructions on how to recognize potential complications and how to obtain medical assistance in the event of such complications. A responsible adult is also provided with information regarding recognition of potential complications and is available to transport the patient and be in attendance during the initial night after discharge" [1].
- 4. "The patient is free of concurrent serious medical illness that might contribute to a significantly increased risk of complication" [1].
- 5. "The patient has recovered from the effects of the sedation" [1].

#### G. Relative Contraindications to Short-Term Observation

- 1. "Several factors must be considered when determining the length of postprocedure skilled nursing care" [1]. Some of the relative contraindications to short-term observation are listed below [1].
- 2. "Patients with coagulopathies or electrolyte abnormalities that require correction should be hospitalized until stable" [1].
- 3. "Insulin-dependent diabetics who have labile serum glucose levels in the periprocedural period should be hospitalized until stable" [1].
- 4. "Complications occurring during or after IVC filter placement, including large hematoma, anuria, and persistent nausea and vomiting, should prompt observation until symptoms resolve" [1].
- 5. "Patients who exhibit hemodynamic instability or significant dysrhythmia during or after the procedure should be hospitalized until stable" [1].
- 6. "Patients with concurrent serious medical illness that might contribute to a significantly increased risk of complication should be hospitalized until stable" [1].
- 7. "Patients with impaired mental or neurologic status who do not have adequate assistance to provide care as needed should be hospitalized until appropriate assistance is available or no longer required" [1].

"The decision for short-term or longer-term postprocedure observation must be individualized, and a patient's care may vary from the above criteria for sound clinical reasons. The decision in each case must be made by the physician who performed the procedure and the referring physician after review of all pertinent data" [1].

# **V. DOCUMENTATION**

Reporting should be in accordance with the <u>ACR–SIR–SPR Practice Parameter for the Reporting and Archiving of Interventional Radiology Procedures</u> [53].

These data should be utilized in conjunction with the thresholds described in section X below to assess filter placement procedural efficacy and complication rates and to trigger institutional review when these thresholds are exceeded.

# **VI. RADIATION SAFETY IN IMAGING**

Radiologists, medical physicists, non-physician radiology providers, radiologic technologists, and all supervising physicians have a responsibility for safety in the workplace by keeping radiation exposure to staff, and to society as a whole, "as low as reasonably achievable" (ALARA) and to assure that radiation doses to individual patients are appropriate, taking into account the possible risk from radiation exposure and the diagnostic image quality necessary to achieve the clinical objective. All personnel who work with ionizing radiation must understand the key principles of occupational and public radiation protection (justification, optimization of protection, application of dose constraints and limits) and the principles of proper management of radiation dose to patients (justification, optimization including the use of dose reference levels). <a href="https://www-pub.iaea.org/MTCD/Publications/PDF/PUB1775\_web.pdf">https://www-pub.iaea.org/MTCD/Publications/PDF/PUB1775\_web.pdf</a>

Nationally developed guidelines, such as the <u>ACR's Appropriateness Criteria</u>®, should be used to help choose the most appropriate imaging procedures to prevent unnecessary radiation exposure.

Facilities should have and adhere to policies and procedures that require ionizing radiation examination protocols (radiography, fluoroscopy, interventional radiology, CT) to vary according to diagnostic requirements and patient body habitus to optimize the relationship between appropriate radiation dose and adequate image quality. Automated dose reduction technologies available on imaging equipment should be used, except when inappropriate for a specific exam. If such technology is not available, appropriate manual techniques should be used.

Additional information regarding patient radiation safety in imaging is available from the following websites – Image Gently® for children (<a href="www.imagegently.org">www.imagegently.org</a>) and Image Wisely® for adults (<a href="www.imagewisely.org">www.imagewisely.org</a>). These advocacy and awareness campaigns provide free educational materials for all stakeholders involved in imaging (patients, technologists, referring providers, medical physicists, and radiologists).

Radiation exposures or other dose indices should be periodically measured by a Qualified Medical Physicist in accordance with the applicable ACR Technical Standards. Monitoring or regular review of dose indices from patient imaging should be performed by comparing the facility's dose information with national benchmarks, such as the ACR Dose Index Registry and relevant publications relying on its data, applicable ACR Practice Parameters, NCRP Report No. 172, Reference Levels and Achievable Doses in Medical and Dental Imaging: Recommendations for the United States or the Conference of Radiation Control Program Director's National Evaluation of X-ray Trends; 2006, 2009, amended 2013, revised 2023 (Res. 2d).

# VII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education, appearing under the heading ACR Position Statement on Quality Control & Improvement, Safety, Infection Control, and Patient Education on the ACR website (https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement).

## **VIII. QUALITY IMPROVEMENT**

## A. Success Rates and Thresholds [1]

Although practicing physicians should strive to achieve perfect outcomes (eg, 100% success, 0% complications), in practice, all physicians will fall short of this ideal to a variable extent. Thus, indicator thresholds may be used to assess the efficacy of ongoing improvement programs.

For the purpose of these practice parameters, a threshold is a specific level of an indicator that should prompt a review. Individual complications may also be associated with complication-specific thresholds. When measures such as indications or success rates fall below a minimum threshold or when complication rates exceed a maximum threshold, a review should be performed to determine causes and to implement changes, if necessary.

Thresholds may vary from those listed here; for example, patient referral patterns and selection factors may dictate a different threshold value for a particular indicator at a particular institution. Thus, setting universal thresholds is very difficult, and each department is urged to alter the thresholds as needed to higher or lower values to meet its own quality improvement program needs [1].

"Complications can be stratified on the basis of outcome. Major complications result in admission to a hospital for therapy (for outpatient procedures), an unplanned increase in the level of care, prolonged hospitalization, permanent adverse sequelae, or death. Minor complications result in no sequelae; they may require nominal therapy or a short hospital stay for observation (generally overnight)" [1]. For further information, see the <a href="Proposal of a New Adverse Event Classification by the Society of Interventional Radiology Standards of Practice Committee">Committee</a>. The complication rates and thresholds below refer to major complications.

"It is expected that the technical success for percutaneously placed IVC filters will be 97% or better in experienced hands. Therefore, the proposed threshold for review of technical failures should be 3%" [1]

- B. Complication Rates and Thresholds [1]
- 1. Complications

"Each currently available filter has been extensively studied as part of the FDA approval process. Few comparative studies have been completed evaluating all filters in one project, and those that have done so have been retrospective analyses. Complication rates are highly variable, depending on the filter being studied. For simplicity, these practice parameters do not suggest threshold rates for each individual filter; rather, filtration devices are considered as a group" [1].

Complications	Reported Rates (%)	Threshold (%)
Death [1,55]	0.12	< 1
Filter embolization [1,2,28,46,47,56-68]	0.1	1
Deployment outside target area [1,27,69,70]	1-9	< 1
Access site thrombosis occlusive [1,71,72]	3-10	3

<sup>&</sup>quot;Published rates for individual types of complications are highly dependent on patient selection and are in some cases based on series comprising several hundred patients, which is a volume larger than most individual practitioners are likely to treat. It is also recognized that a single complication can cause a rate to cross above a complication-specific threshold when the complication occurs within a small patient volume (eg, early in a quality improvement program)" [1].

#### 2. Other trackable events

"Because an IVC filter may be implanted as a permanent device (if not retrieved) and can be used in relatively young patients, several other trackable parameters are appropriate to record in a quality improvement program. The (following) events may or may not be clinically significant in a particular patient. For this reason, thresholds for these events are not included in this document" [1].

#### TABLE 2

Other Trackable Events	Reported Rates (%)
Other Trackable Events	Reported Rates (%)
IVC penetration* [1,3,5,6,8,9,14,27-34,37,54,55,61,62,64,66,68,70-80]	0-100
Migration of filter/filter components* [1,3,4,6,9,11,12,14,17,26-28,30,35- 38,46,55,58,59,66,68,70,73,74,76-89]	0-25
Filter fracture [1-10,14,27,28,31,33,54,56,57,61,63-65,69,74,76-78,80,82,84,85,87-90]	0-50
Recurrent PE [1,28,74,76,82,86]	0.5-6
Access site thrombus – all types [1,55,71,87,88,90]	0-25
IVC occlusion [1,5,7,8,13,19,20,23,27,28,30,36,38,40,46,47,54,59-61,64,72-74,76-81,83,86,89,91,92]	2-65
Insertion problems [1,28,55,64,74,76,82,83,86,88,91,92]	5-23
Other complications [1,79,93]	1-15

The data in the table represents reported outcomes from various publications and not the SIR standard for complications.

[1] Reprinted with modifications from Journal of Vascular Interventional Radiology, Caplin DM, Nikolic B, Kalva SP, et al., Quality Improvement Guidelines for the Performance of Inferior Vena Cava Filter Placement for the Prevention of Pulmonary Embolism, 1499-1506, 2011, with permission from Elsevier.

#### **ACKNOWLEDGEMENTS**

This practice parameter was revised according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (<a href="https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards">https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards</a>) by the Committee on Practice Parameters — Interventional and Cardiovascular Radiology of the ACR Commission on Interventional and Cardiovascular Radiology and the Committee on Practice Parameters — Pediatric Radiology of the ACR Commission on Pediatric Radiology, in collaboration with the SIR and the SPR.

<u>Writing Committee</u> – members represent their societies in the initial and final revision of this practice parameter

<sup>\*</sup> The rate of clinically significant penetration is not precisely known [89].

Douglas M. Coldwell, MD, PhD, Chair Suvranu Ganguli, MD, FSIR Manish N. Patel, DO

Kevin W. Dickey, MD, FACR Sanjeeva P. Kalva, MD, FSIR Abhay S. Srinivasan, MD

Matthew P. Lungren, MD, MPH Ravi N. Srinivasa, MD, FSIR

<u>Committee on Practice Parameters – Interventional and Cardiovascular Radiology</u>

(ACR Committee responsible for sponsoring the draft through the process)

Drew M. Caplin, MD, Chair Margaret Hsin-Shung Lee, MD, FACR

Chaitanya Ahuja, MBBS Mary Lee Jensen, MD, FACR

Douglas M. Coldwell, MD, PhD Claire Kaufman, MD Mandeep S. Dagli, MD Dennis Kay, MD, FACR

Kevin W. Dickey, MD, FACR Kennith F. Layton, MD, FACR

Meredith J. Englander, MD M. Victoria Marx, MD

C. Matthew Hawkins, MD Christopher D Yeisley, MD

Committee on Practice Parameters - Pediatric Radiology

(ACR Committee responsible for sponsoring the draft through the process)

Terry L. Levin, MD, FACR, Chair Jane Sun Kim, MD

Jennifer A Knight, MD John B. Amodio, MD, FACR

Jesse Berman, MD Jessica Kurian, MD

Tara M. Catanzano, MB, BCh Matthew P. Lungren, MD, MPH

Harris L. Cohen, MD, FACR Helen R. Nadel, MD Kassa Darge, MD, PhD Erica Poletto, MD

Dorothy L. Gilbertson-Dahdal, MD Richard B. Towbin, MD, FACR

Lauren P. Golding, MD Andrew T. Trout, MD Safwan S. Halabi, MD Esben S. Vogelius, MD

Jason Higgins, DO

Alan H. Matsumoto, MD, FACR, Chair, Commission on Interventional and Cardiovascular Radiology

Richard A. Barth, MD, FACR, Chair, Commission on Pediatric Radiology David B. Larson, MD, MBA, Chair, Commission on Quality and Safety

Mary S. Newell, MD, FACR, Chair, Committee on Practice Parameters and Technical Standards

**Comments Reconciliation Committee** 

Traci Pritchard, MD, FACR, Chair Terry L. Levin, MD, FACR

Derrick Siebert, MD, Co-Chair Matthew P. Lungren, MD, MPH Richard A. Barth, MD, FACR Alan H. Matsumoto, MD, FACR

Drew M. Caplin, MD Mary S. Newell, MD, FACR

Douglas M. Coldwell, MD, PhD Indravadan Patel, MD Kush Desai, MD Manish N. Patel, DO

Kevin W. Dickey, MD, FACR Andrew Picel, MD

Richard Duszak Jr., MD, FACR Ravi N. Srinivasa, MD, FSIR Suvranu Ganguli, MD, FSIR William F. Sensakovic, PhD

Sanjeeva P. Kalva, MD, FSIR Abhay S. Srinivasan, MD

Amy L. Kotsenas, MD, FACR Ron Winokur, MD

**REFERENCES** 

David B. Larson, MD, MBA

- **1.** Caplin DD, Nikolic BB, Kalva SS, et al. Quality improvement guidelines for the performance of inferior vena cava filter placement for the prevention of pulmonary embolism. J Vasc Interv Radiol 22:1499-506, .
- **2.** Angel LF, Tapson V, Galgon RE, Restrepo MI, Kaufman J. Systematic review of the use of retrievable inferior vena cava filters. J Vasc Interv Radiol. 2011;22(11):1522-1530 e1523.
- **3.** Asch MR. Initial experience in humans with a new retrievable inferior vena cava filter. Radiology. 2002 Dec;225(3):835-44.
- 4. Dorfman GG. Percutaneous inferior vena caval filters. Radiology 174:987-92, .
- **5.** Given MM, McDonald BB, Brookfield PP, et al. Retrievable Gunther Tulip inferior vena cava filter: experience in 317 patients. J Med Imaging Radiat Oncol 52:452-7, .
- **6.** Grassi CC. Inferior vena caval filters: analysis of five currently available devices. AJR Am J Roentgenol 156:813-21, .
- **7.** Greenfield LL, DeLucia AA. Endovascular therapy of venous thromboembolic disease. Surg Clin North Am 72:969-89, .
- **8.** Hoppe HH, Nutting CC, Smouse HH, et al. Günther Tulip filter retrievability multicenter study including CT follow-up: final report. J Vasc Interv Radiol 17:1017-23, .
- **9.** Johnson MM, Nemcek AA, Benenati JJ, et al. The safety and effectiveness of the retrievable option inferior vena cava filter: a United States prospective multicenter clinical study. J Vasc Interv Radiol 21:1173-84, .
- 10. Kinney TB. Update on inferior vena cava filters. J Vasc Interv Radiol. 2003 Apr;14(4):425-40.
- **11.** Le Blanche AA, Benazzouz AA, Reynaud PP, et al. The VenaTech LP permanent caval filter: effectiveness and safety in the prevention of pulmonary embolism--a European multicenter study. J Vasc Interv Radiol 19:509-15, .
- **12.** Linsenmaier UU, Rieger JJ, Schenk FF, Rock CC, Mangel EE, Pfeifer KK. Indications, management, and complications of temporary inferior vena cava filters. Cardiovasc Intervent Radiol 21:464-9, .
- **13.** Lyon SS, Riojas GG, Uberoi RR, et al. Short- and long-term retrievability of the Celect vena cava filter: results from a multi-institutional registry. J Vasc Interv Radiol 20:1441-8, .
- 14. Millward SS. Temporary and retrievable inferior vena cava filters: current status. J Vasc Interv Radiol 9:381-7, .
- **15.** Millward SF, Oliva VL, Bell SD, et al. Günther Tulip Retrievable Vena Cava Filter: results from the Registry of the Canadian Interventional Radiology Association. J Vasc Interv Radiol. 2001 Sep;12(9):1053-8.
- **16.** Oliva VV, Perreault PP, Giroux MM, Bouchard LL, Therasse EE, Soulez GG. Recovery G2 inferior vena cava filter: technical success and safety of retrieval. J Vasc Interv Radiol 19:884-9, .
- **17.** Savader SJ. Inferior vena cava filters. In: Savader SJ, Trerotola SO, eds. Venous Interventional Radiology with Clinical Perspective. New York, NY: Thieme; 1996:367-99.
- **18.** Smouse HB, Mendes R, Bosiers M, Van Ha TG, Crabtree T. The RETRIEVE trial: safety and effectiveness of the retrievable crux vena cava filter. J Vasc Interv Radiol. 2013;24(5):609-621.
- **19.** Yamagami T, Kato T, Iida S, Tanaka O, Nishimura T. Retrievable vena cava filter placement during treatment for deep venous thrombosis. Br J Radiol. 2003 Oct;76(910):712-8.
- **20.** Ziegler JJ, Dietrich GG, Cohen SS, Sterling KK, Duncan JJ, Samotowka MM. PROOF trial: protection from pulmonary embolism with the OptEase filter. J Vasc Interv Radiol 19:1165-70, .
- **21.** Aidinian GG, Fox CC, White PP, Cox MM, Adams EE, Gillespie DD. Intravascular ultrasound--guided inferior vena cava filter placement in the military multitrauma patients: a single-center experience. Vasc Endovascular Surg 43:497-501, .
- **22.** Kassavin DD, Constantinopoulos GG. The transition to IVUS-guided IVC filter deployment in the nontrauma patient. Vasc Endovascular Surg 45:142-5, .
- **23.** Gunn AA, Iqbal SS, Kalva SS, et al. Intravascular ultrasound-guided inferior vena cava filter placement using a single-puncture technique in 99 patients. Vasc Endovascular Surg 47:97-101, .
- **24.** Ganguli S, Hawkins BM, Abtahian F, et al. Comparison of Inferior Vena Cava Filters Placed at the Bedside via Intravenous Ultrasound Guidance Versus Fluoroscopic Guidance. Annals of Vascular Surgery. 39:250-255, 2017 Feb.Ann Vasc Surg. 39:250-255, 2017 Feb.
- **25.** Grassi CC, Swan TT, Cardella JJ, et al. Quality improvement guidelines for percutaneous permanent inferior vena cava filter placement for the prevention of pulmonary embolism. J Vasc Interv Radiol 14:S271-5, .
- **26.** U.S. Food and Drug Administration. Removing retrievable inferior vena cava filters: FDA safety communication. Available at: http://www.fda.gov/MedicalDevices/Safety/AlertsandNotices/ucm396377.htm .
- 27., Baglin TT, Brush JJ, Streiff MM. Guidelines on use of vena cava filters. Br J Haematol 134:590-5, .
- **28.** Ferris EJ, McCowan TC, Carver DK, McFarland DR. Percutaneous inferior vena caval filters: follow-up of seven designs in 320 patients. Radiology. 188(3):851-6, 1993 Sep.Radiology. 188(3):851-6, 1993 Sep.
- **29.** Kalva SS, Chlapoutaki CC, Wicky SS, Greenfield AA, Waltman AA, Athanasoulis CC. Suprarenal inferior vena cava filters: a 20-year single-center experience. J Vasc Interv Radiol 19:1041-7, .

- **30.** Kaufman JJ, Geller SS. When to use an inferior vena cava filter. AJR Am J Roentgenol 164:256-7, .
- **31.** Kaufman JJ, Rundback JJ, Kee SS, et al. Development of a research agenda for inferior vena cava filters: proceedings from a multidisciplinary research consensus panel. J Vasc Interv Radiol 20:697-707, .
- **32.** Kearon C, Akl EA, Comerota AJ, et al. Antithrombotic therapy for VTE disease: Antithrombotic Therapy and Prevention of Thrombosis, 9th ed: American College of Chest Physicians Evidence-Based Clinical Practice Guidelines.[Erratum appears in Chest. 2012 Dec;142(6):1698-1704]. Chest. 141(2 Suppl):e419S-e496S, 2012 Feb.
- **33.** Norris CC, Greenfield LL, Herrmann JJ. Free-floating iliofemoral thrombus. A risk of pulmonary embolism. Arch Surg 120:806-8, .
- 34. Ray CC, Kaufman JJ. Complications of inferior vena cava filters. Abdom Imaging 21:368-74, .
- **35.** Rogers FB, Cipolle MD, Velmahos G, Rozycki G, Luchette FA. Practice management guidelines for the prevention of venous thromboembolism in trauma patients: the EAST practice management guidelines work group. J Trauma. 2002 Jul;53(1):142-64.
- 36. Rutherford RR. Prophylactic indications for vena cava filters: critical appraisal. Semin Vasc Surg 18:158-65, .
- 37. Vaiji K. Vascular and interventional radiology. Philadelphia, Pa: WB Saunders; 1999.
- **38.** Eifler AC, Lewandowski RJ, Gupta R, et al. Optional or permanent: clinical factors that optimize inferior vena cava filter utilization. J Vasc Interv Radiol. 2013;24(1):35-40.
- **39.** Kaufman JA, Barnes GD, Chaer RA, et al. Society of Interventional Radiology Clinical Practice Guideline for Inferior Vena Cava Filters in the Treatment of Patients with Venous Thromboembolic Disease: Developed in collaboration with the American College of Cardiology, American College of Chest Physicians, American College of Surgeons Committee on Trauma, American Heart Association, Society for Vascular Surgery, and Society for Vascular Medicine. Journal of Vascular & Interventional Radiology. 31(10):1529-1544, 2020 Oct. J Vasc Interventional Radiol. 31(10):1529-1544, 2020 Oct.
- 40. Kinney TT. Inferior vena cava filters. Semin Intervent Radiol 23:230-9, .
- **41.** Stein PP, Matta FF. Case fatality rate with pulmonary embolectomy for acute pulmonary embolism. Am J Med 125:471-7, .
- **42.** Stein PD, Matta F, Keyes DC, Willyerd GL. Impact of vena cava filters on in-hospital case fatality rate from pulmonary embolism. Am J Med. 2012;125(5):478-484.
- **43.** Stein PP, Matta FF, Hughes MM. Inferior Vena Cava Filters in Stable Patients With Pulmonary Embolism and Heart Failure. Am J Cardiol 124:292-295, .
- **44.** Wadhwa V, Gutta NB, Trivedi PS, et al. In-Hospital Mortality Benefit of Inferior Vena Cava Filters in Patients With Pulmonary Embolism and Congestive Heart Failure. AJR Am J Roentgenol. 211(3):672-676, 2018 09.
- **45.** Benfayed WW, Torreggiani WW, Hamilton SS. Detection of pulmonary emboli resulting from ovarian vein thrombosis. AJR Am J Roentgenol 181:1430-1, .
- **46.** Chaudry GG, Padua HH, Alomari AA. The use of inferior vena cava filters in young children. J Vasc Interv Radiol 19:1103-6, .
- **47.** Kukreja KK, Gollamudi JJ, Patel MM, Johnson NN, Racadio JJ. Inferior vena cava filters in children: our experience and suggested guidelines. J Pediatr Hematol Oncol 33:334-8, .
- **48.** American College of Radiology. ACR Practice Parameter for Continuing Medical Education. Available at https://gravitas.acr.org/PPTS/GetDocumentView?docId=130+&releaseId=2
- **49.** American College of Radiology. ACR—AAPM TECHNICAL STANDARD FOR DIAGNOSTIC MEDICAL PHYSICS PERFORMANCE MONITORING OF FLUOROSCOPIC EQUIPMENT. Available at
- https://gravitas.acr.org/PPTS/GetDocumentView?docId=117+&releaseId=2
- **50.** American College of Radiology. ACR-AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Radiographic Equipment. Available at
- https://gravitas.acr.org/PPTS/GetDocumentView?docId=120+&releaseId=2
- **51.** American College of Radiology. ACR–SIR Practice Parameter For Minimal and/or Moderate Sedation/Analgesia. Available at https://gravitas.acr.org/PPTS/GetDocumentView?docId=95+&releaseId=2
- **52.** American College of Radiology. ACR–SIR–SNIS–SPR Practice Parameter for the Clinical Practice of Interventional Radiology. Available at https://gravitas.acr.org/PPTS/GetDocumentView?docId=36+&releaseId=2
- **53.** American College of Radiology. ACR–SIR–SPR Practice Parameter for the Reporting and Archiving of Interventional Radiology Procedures. Available at
- https://gravitas.acr.org/PPTS/GetDocumentView?docId=39+&releaseId=2
- **54.** American College of Radiology. ACR–SIR–SPR Practice Parameter on Informed Consent for Image-Guided Procedures. Available at https://gravitas.acr.org/PPTS/GetDocumentView?docld=176+&releaseId=2
- **55.** Becker DM, Philbrick JT, Selby JB. Inferior vena cava filters. Indications, safety, effectiveness. Arch Intern Med. 1992 Oct;152(10):1985-94.

- **56.** Akins CC, Thurer RR, Waltman AA, Margolies MM, Schneider RR. A misplaced caval filter: its removal from the heart without cardiopulmonary bypass. Arch Surg 115:1133, .
- **57.** Athanasoulis CA, Kaufman JA, Halpern EF, Waltman AC, Geller SC, Fan CM. Inferior vena caval filters: review of a 26-year single-center clinical experience. Radiology. 2000 Jul;216(1):54-66.
- **58.** Bach JJ, Zaneuski RR, Lee HH. Cardiac arrhythmias from a malpositioned Greenfield filter in a traumatic quadriplegic. Am J Phys Med Rehabil 69:251-3, .
- **59.** Castaneda FF, Herrera MM, Cragg AA, et al. Migration of a Kimray-Greenfield filter to the right ventricle. Radiology 149:690, .
- **60.** Friedell MM, Goldenkranz RR, Parsonnet VV, et al. Migration of a Greenfield filter to the pulmonary artery: a case report. J Vasc Surg 3:929-31, .
- **61.** Hammond CC, Bakshi DD, Currie RR, et al. Audit of the use of IVC filters in the UK: experience from three centres over 12 years. Clin Radiol 64:502-10, .
- **62.** LaPlante JJ, Contractor FF, Kiproff PP, Khoury MM. Migration of the Simon nitinol vena cava filter to the chest. AJR Am J Roentgenol 160:385-6, .
- **63.** Loesberg AA, Taylor FF, Awh MM. Dislodgment of inferior vena caval filters during "blind" insertion of central venous catheters. AJR Am J Roentgenol 161:637-8, .
- **64.** Magnant JG, Walsh DB, Juravsky LI, Cronenwett JL. Current use of inferior vena cava filters. J Vasc Surg. 1992 Nov;16(5):701-6.
- **65.** Puram BB, Maley TT, White NN, Rotman HH, Miller GG. Acute myocardial infarction resulting from the migration of a Greenfield filter. Chest 98:1510-1, .
- 66. Simon M. Vena cava filters: prevalent misconceptions. J Vasc Interv Radiol. 1999 Sep;10(8):1021-4.
- **67.** Urbaneja AA, Fontaine AA, Bruckner MM, Spigos DD. Evulsion of a Vena Tech filter during insertion of a central venous catheter. J Vasc Interv Radiol 5:783-5, .
- **68.** Villard JJ, Detry LL, Clermont AA, Pinet FF. [8 cases of Greenfield filters in the right heart cavities. Their surgical treatment]. Ann Radiol (Paris) 30:102-4, .
- **69.** Savin MA, Panicker HK, Sadiq S, Albeer YA, Olson RE. Placement of vena cava filters: factors affecting technical success and immediate complications. AJR Am J Roentgenol. 2002 Sep;179(3):597-602.
- **70.** Cipolla J, Wegner NS, Sharma R, et al. Complications of vena cava filters: a comprehensive clinical review. OPUS 12 Scientist 2008;2:11-23.
- **71.** Hicks MM, Middleton WW, Picus DD, Darcy MM, Kleinhoffer MM. Prevalence of local venous thrombosis after transferoral placement of a Bird's Nest vena caval filter. J Vasc Interv Radiol 1:63-8, .
- **72.** Luanow E, Kandarpa K, Chopra et al. Bleeding complications in patients undergoing percutaneous vena cava filter placement using new low profile introduction systems. San Francisco, Calif: American Roentgen Ray Annual Meeting; 1993.
- **73.** Greenfield LL, Cho KK, Proctor MM, et al. Results of a multicenter study of the modified hook-titanium Greenfield filter. J Vasc Surg 14:253-7, .
- **74.** Greenfield LL, Proctor MM. Vena caval filter use in patients with sepsis: results in 175 patients. Arch Surg 138:1245-8, .
- **75.** Laliberte F, Coleman CI, Bookhart B, et al. Weekly risk of venous thromboembolism recurrence in patients receiving oral anticoagulants. Curr Med Res Opin. 30(8):1513-20, 2014 Aug.
- **76.** Simon MM, Athanasoulis CC, Kim DD, et al. Simon nitinol inferior vena cava filter: initial clinical experience. Work in progress. Radiology 172:99-103, .
- **77.** McCowan TT, Ferris EE, Carver DD, Molpus WW. Complications of the nitinol vena caval filter. J Vasc Interv Radiol 3:401-8, .
- **78.** Ricco JJ, Crochet DD, Sebilotte PP, et al. Percutaneous transvenous caval interruption with the "LGM" filter: early results of a multicenter trial. Ann Vasc Surg 2:242-7, .
- **79.** Teitelbaum GG, Jones DD, van Breda AA, et al. Vena caval filter splaying: potential complication of use of the titanium Greenfield filter. Radiology 173:809-14, .
- **80.** Deso SS, Idakoji II, Kuo WW. Evidence-Based Evaluation of Inferior Vena Cava Filter Complications Based on Filter Type. Semin Intervent Radiol 33:93-100, .
- 81. Athanasoulis CC. Complications of vena cava filters. Radiology 188:614-5, .
- **82.** Crochet DD, Stora OO, Ferry DD, et al. Vena Tech-LGM filter: long-term results of a prospective study. Radiology 188:857-60, .
- **83.** Hye RR, Mitchell AA, Dory CC, Freischlag JJ, Roberts AA. Analysis of the transition to percutaneous placement of Greenfield filters. Arch Surg 125:1550-3, .
- 84. Lang W, Schweiger H, Hofmann-Preiss K. Results of long-term venacavography study after placement of a

Greenfield vena caval filter. J Cardiovasc Surg (Torino). 1992;33(5):573-8.

- **85.** Mobin-Uddin K, Smith PE, Martinez LO, et al. A vena caval filter for the prevention of pulmonary embolus. Surg Forum 1967;18:209-11.
- **86.** Millward SS, Peterson RR, Moher DD, et al. LGM (Vena Tech) vena caval filter: experience at a single institution. J Vasc Interv Radiol 5:351-6, .
- **87.** Molgaard CC, Yucel EE, Geller SS, Knox TT, Waltman AA. Access-site thrombosis after placement of inferior vena cava filters with 12-14-F delivery sheaths. Radiology 185:257-61, .
- **88.** Vesely T. Technical problems and complications associated with inferior vena cava filters. Sem Interven Radiol 1994;11:121-33.
- **89.** Stawicki SS, Sims CC, Sharma RR, et al. Vena cava filters: a synopsis of complications and related topics. J Vasc Access 9:102-10, .
- **90.** Hackett C, Parker RA, Gemmete JJ. Inferior Vena Cava (IVC) filter position and its effect on the incidence of deep venous thrombosis. J Vascu Interv Radiol 2013;24:S87.
- **91.** Moore BB, Valji KK, Roberts AA, Bookstein JJ. Transcatheter manipulation of asymmetrically opened titanium Greenfield filters. J Vasc Interv Radiol 4:687-90, .
- **92.** Sweeney TT, Van Aman MM. Deployment problems with the titanium Greenfield filter. J Vasc Interv Radiol 4:691-4, .
- 93. Clagett GG. Basic data related to venous thromboembolism. Ann Vasc Surg 2:402-5, .
- \*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised, or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

<u>Development Chronology for This Practice Parameter</u>

2000 (Resolution 12)

Amended 2004 (Resolution 25)

Revised 2005 (Resolution 40)

Amended 2006 (Resolution 16g, 17, 34, 35, 36)

Amended 2007 (Resolution 12m, 38)

Revised 2010 (Resolution 46)

Amended 2014 (Resolution 39)

Revised 2016 (Resolution 18)

Amended 2020 (Resolution 8)

Revised 2021 (Resolution 8)

Amended 2023 (Resolution 2c, 2d)

# Appendix A

For the purpose of this practice parameter, the following definitions apply [1,25,28,34]:

Permanent placement – deployment in those situations where lifelong mechanical protection against thromboembolic episodes is needed

Temporary placement – deployment in those situations where time-limited mechanical protection against thromboembolic episodes is needed

Procedural success – deployment of a filter in the intended location (target zone) such that the filter is judged suitable for mechanical protection against PE

Postfilter recurrent PE – PE occurring after filter placement and documented by pulmonary arteriography, cross-sectional imaging, significant change in ventilation-perfusion (V/Q) lung scan indicative of recurrent PE, surgical embolectomy, or autopsy

Postfilter thrombotic occlusion of the IVC – presence of an occluding thrombus in the filter and/or the IVC occurring after filter insertion and documented by ultrasound, CT, MRI, venography, or autopsy; this may be symptomatic or asymptomatic.

Filter penetration – penetration of the vein wall by a filter strut with transmural incorporation. For quality improvement reporting purposes, the definition of IVC penetration by a filter is filter strut or anchor devices extending >3 mm outside the wall of the IVC as demonstrated by CT, venography, or autopsy. Acute penetration occurring during placement of the filter is considered an insertion complication (see below).

Filter embolization – postdeployment movement of the filter or its components to a distant anatomic site completely out of the target zone

Filter movement – a change in filter position compared to its deployed position (either cranial or caudal) of >2 cm Revised 2021 (Republic Plant Film imaging, CT, or venography

Filter fracture – any loss of a filter's structural integrity (ie, breakage or separation) documented by imaging or autopsy

Insertion complication – malfunctions of the filter or deployment system such as incomplete filter opening, filter tilt >15° from the IVC axis, misplacement of filter outside the infrarenal IVC when the operator's intent is to place the filter in the infrarenal IVC (eg, when a portion of the filter is within one iliac vein), deploying the filter in the incorrect orientation (eg, upside down in the IVC), or prolapse of filter components. Filter malposition requiring surgical/endovascular removal is considered an insertion problem complication.

Access site thrombus – occlusive or nonocclusive thrombus developing at the venotomy site after filter insertion and documented by ultrasound or other imaging

Access site complications with clinical sequelae – arteriovenous fistula, hematoma, or bleeding requiring transfusion, hospitalization (either admission or extended stay), or further treatment