ACR-SPR-SSR PRACTICE PARAMETER FOR THE PERFORMANCE AND INTERPRETATION OF MAGNETIC RESONANCE IMAGING (MRI) OF THE HIP AND PELVIS FOR MUSCULOSKELETAL DISORDERS

The American College of Radiology, with more than 40,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care $\frac{1}{2}$. For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

<u>1</u> Iowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing, 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the "ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures (Revised 2008)" sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, <u>Stanley v. McCarver</u>, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do

not establish the standard of care.

I. INTRODUCTION

This practice parameter was developed and written collaboratively by the American College of Radiology (ACR), the Society for Pediatric Radiology (SPR), and the Society of Skeletal Radiology (SSR).

This parameter addresses magnetic resonance imaging (MRI) performed to evaluate musculoskeletal disorders of the pelvis and hips and to investigate symptoms that are believed to originate in the musculoskeletal system. Guidelines for pelvic MRI examinations performed to evaluate the male and female genitourinary tracts, bowel, and vasculature are not included herein (see the <u>ACR–SAR–SPR Practice Parameter for the Performance of Magnetic Resonance Imaging (MRI) of the Soft-Tissue Components of the Pelvis [1]).</u>

MRI is a proven, established imaging method for the detection, evaluation, staging, and follow-up of musculoskeletal conditions of the hip and pelvis. Properly performed and interpreted, MRI not only contributes to diagnosis but also serves as an important guide to treatment planning and prognostication [2-11]. However, MRI should be performed only for a valid medical reason and only after careful consideration of alternative imaging methods. The strengths of MRI and other imaging techniques should be weighed as to their suitability in particular patients and clinical conditions.

Radiographs should be the initial imaging study for most suspected abnormalities of the hip and pelvis [12]. Sequential radiographs are a key component in the postoperative evaluation of hip arthroplasty and other orthopedic procedures [13]. Bone scintigraphy is used to screen the entire skeleton for conditions such as metastases. Additionally, with some limitations [14], bone scans can also detect radiographically occult osteonecrosis, fractures [15], and stress fractures [16] in the hips and pelvis. Bone scintigraphy and labeled leukocyte scintigraphy may also have a role in the evaluation of symptomatic hip arthroplasties [20,21]. Because of its superior sensitivity and specificity, however, MRI has largely replaced scintigraphy for these indications [12,17-19]. Ultrasound may be used to detect tendon disorders in the proximal thighs [22-24], bursitis, synovitis, and joint effusion [25-27]. Ultrasound is useful in the assessment of the snapping hip due to external and internal causes [28,29]. Ultrasound is a useful tool for guiding hip and pelvis injections and aspirations. In children, sonography can be used to diagnose developmental dysplasia of the hip, hip effusions, and pelvic apophyseal avulsions [30]. Ultrasound can be used to evaluate developmental hip dysplasia in infants and young children [31]. In adults, the clinical response to intra-articular anesthetic injection in the hip helps predict intra-articular pathology [32,33]. Therapeutic injection of the pubic symphysis cleft may help diagnose the cause of groin pain in athletes [34,35]. Hip arthroscopy, an invasive procedure, provides a detailed examination of the internal structures of the hip joint, allowing the surgeon to treat as well as diagnose many internal derangements [36,37].

Computed tomography (CT), especially with multidetector helical scanners using thin collimation, is often preferred to MRI for detailed evaluation of bony alignment, morphology, and cortical pathology. Multiplanar 2-D reformatting and 3-D volume rendering increase the utility of CT for orthopedic purposes. Typical applications include evaluation of the acetabulum and hip joint after fractures or dislocations [38-40], preoperative planning for complex pelvic osteotomies and arthroplasties [39,41], preoperative planning of complex osteotomies in femoroacetabular impingement and surgical navigation [42,43], and evaluation of osteolysis around hip arthroplasty components [44-46]. MRI has largely replaced CT for detecting femoral head osteonecrosis [47], but CT is still valuable for detecting subchondral fractures in necrotic femoral heads [48]. CT can detect radiographically occult hip fractures, but MRI is more sensitive [49,50]. CT is a reasonable secondary imaging method (after MRI) for soft-tissue disorders such as sports hernias [51] and monoarticular proliferative arthropathies [12]. CT can detect erosions in patients with suspected sacroiliitis, and normal radiographs [52]. Multidetector CT arthrography can detect cartilage and labral lesions [53] in patients with contraindications to MR imaging. Radially oriented multiplanar reformation may be a useful adjunct in the assessment of bone morphology, acetabular cartilage, and labrum [53-55].

Although MRI is often the most sensitive noninvasive diagnostic test for detecting anatomic abnormalities of the hip and pelvis, its findings may be misleading if not closely correlated with the clinical history, physical examination, physiologic tests such as nerve conduction analysis and electromyography, and other imaging

studies. Adherence to the following parameters will enhance the probability of detecting such abnormalities.

II. INDICATIONS

- A. Primary indications for MRI of the hip and pelvis include, but are not limited to, screening, diagnosis, exclusion, grading, and/or prognostication of suspected:
 - 1. Osteonecrosis of the femoral head(s), including staging of osteonecrosis and screening of asymptomatic hips contralateral to a hip with osteonecrosis [2,6,47,56-69]
 - 2. Other marrow abnormalities of the femoral head(s), including transient and migratory osteoporosis of the hip, transient bone marrow edema syndrome, and subchondral insufficiency fractures[1] [66,68,70-74]
 - 3. Radiographically occult traumatic fractures of the proximal femur and pelvis [11,17,18,50,75-82]
 - 4. Stress fractures (fatigue and insufficiency types) of the proximal femur, pelvis, and sacrum, atypical fractures in the setting of osteoporosis therapy [16,19,49,83-92]
 - 5. Childhood hip disorders and their adult sequelae: Legg-Calvé-Perthes disease, slipped capital femoral epiphysis, idiopathic chondrolysis, coxa vara, proximal femoral focal deficiency, and developmental dysplasia of the hip (DDH)², [2] [3,7,41,93-98]
 - 6. Femoroacetabular impingement (cam, pincer, and mixed types), including evaluation of labral, articular cartilage, and bone morphologic abnormalities³ [5,97,99-110]
 - 7. Acetabular labral tears, traumatic and/or degenerative³ [97,99,111-120]
 - 8. Abductor (gluteus medius and minimus) musculotendinous disorders and greater trochanteric bursitis² [121-128]
 - 9. Proximal hamstring musculotendinous disorders and ischial bursitis² [22,129-131]
 - 10. Hip rotator and flexor musculotendinous disorders and iliopsoas bursitis (including quadratus femoris, iliopsoas, rectus femoris) [27,132-137]
 - 11. Athletic pubalgia, including adductor/rectus abdominus musculotendinous disorders and pubic bone, osteitis pubis, and other disorders presenting as "sports hernia" ³ [10,19,34,35,51,88,138-147]
 - 12. Proximal iliotibial band syndrome [148]
 - Osteochondral and chondral abnormalities in the hip joint, including chondral delamination³ [98,102,104,105,111,112,149-152]
 - 14. Ligamentum teres injury (rupture)³ [153,154]
 - 15. Bursitis in and around the pelvis² [27,88,111,122,134,155,156]
 - 16. Sacral plexus abnormalities, nerve entrapment, and piriformis syndrome [91,157,158]
 - 17. Pelvic impingement syndromes, including ischiofemoral and subspine impingement [125] [136,170,171]
 - 18. Osteomyelitis, septic arthritis, and soft-tissue infection of the hip and pelvis² [9,159-164]
- B. MRI of the hip and pelvis may be indicated to further clarify and stage conditions diagnosed clinically and/or suggested by other imaging modalities, including, but not limited to:
 - 1. Hip arthritis and synovitis of unclear etiology: inflammatory, infectious, degenerative, crystalinduced, posttraumatic, proliferative² [160,165-174]
 - 2. Hip joint effusions [27,60,175]
 - 3. Hip joint bodies³ [176]
 - 4. Sacroiliitis² [4,165,177]
 - Primary and secondary bone and soft-tissue tumors of the pelvis, proximal femur, and thigh² [157,178-181] (see also the <u>ACR-SPR-SSR Practice Parameter for the Performance and Interpretation</u> of <u>Magnetic Resonance Imaging (MRI) of Bone and Soft-Tissue Tumors</u> [182])
 - 6. Fractures and dislocations of the hip and pelvis² [30,183,184]
- C. MRI of the hip and pelvis may be useful to evaluate specific clinical scenarios, including, but not limited to:
 - 1. Prolonged, refractory, or unexplained hip, trochanteric, pubic, or pelvic pain³ [88,122,124,126,128,142,147,158,185]

- 1. Pelvic, proximal thigh, or groin pain in athletes³ [19,22,34,35,88,111,138-142,144-147,151,186]
- 2. Acute or chronic hip and pelvis trauma with associated soft-tissue injuries² [40,77,187]
- 3. Pelvic pain after radiation therapy [84,85,87]
- 4. Mechanical symptoms in the hip, including snapping and clicking³ [8,24,37,112,114]
- 5. Following reduction of congenital or acquired hip dislocation in infants and children^{2,3} [67,188-193]
- 6. Symptomatic adults with developmental dysplasia of the hip³ [3,7,41,53,97]
- 7. Patients for whom diagnostic or therapeutic hip arthroscopy is planned³ [33,112,116]
- 8. Hip arthroplasties with suspected soft-tissue or periprosthetic abnormalities^{2,[3]} [13,127,128,194-199]
- 9. Pelvimetry in women with obstructed labor [200,201]

[1] Conditions in which intravenous contrast may be useful

[2] Conditions in which intra-articular contrast (performed by direct intra-articular injection or indirect joint opacification following intravenous administration) may be useful

[3] Conditions in which use of metal artifact reduction pulse sequences (MARS) may be useful

III. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

See the <u>ACR Practice Parameter for Performing and Interpreting Magnetic Resonance Imaging (MRI)</u> [202].

IV. SAFETY GUIDELINES AND POSSIBLE CONTRAINDICATIONS

See the <u>ACR Practice Parameter for Performing and Interpreting Magnetic Resonance Imaging (MRI)</u> [202] and the <u>ACR Guidance Document on MR Safe Practices 2020</u> [203].

Peer-reviewed literature pertaining to MR safety should be reviewed on a regular basis [204,205].

V. SPECIFICATIONS OF THE EXAMINATION

The written or electronic request for MRI of the hip and pelvis for musculoskeletal disorders should provide sufficient information to demonstrate the medical necessity of the examination and allow for its proper performance and interpretation.

Documentation that satisfies medical necessity includes 1) signs and symptoms and/or 2) relevant history (including known diagnoses). Additional information regarding the specific reason for the examination or a provisional

diagnosis would be helpful and may at times be needed to allow for the proper performance and interpretation of

the examination.

The request for the examination must be originated by a physician or other appropriately licensed health care provider. The accompanying clinical information should be provided by a physician or other appropriately licensed health care provider familiar with the patient's clinical problem or question and consistent with the state's scope of practice requirements. (ACR Resolution 35 adopted in 2006 – revised in 2016, Resolution 12-b)

The supervising physician must have adequate understanding of the indications, risks, and benefits of the imaging examination as well as alternative imaging procedures. The physician must be familiar with potential hazards associated with MRI, including potential adverse reactions to contrast media. The physician should be familiar with relevant ancillary studies that the patient may have undergone. The physician performing MRI interpretation must have a clear understanding and knowledge of the anatomy and pathophysiology relevant to the MRI examination.

The supervising physician must also understand the pulse sequences to be used and their effect on the appearance of the images, including the potential generation of image artifacts. Standard imaging protocols may be established and varied on a case-by-case basis when necessary. These protocols should be reviewed and updated periodically.

A. Patient Selection

The physician responsible for the examination should supervise patient selection and preparation and be available in person or by telephone for consultation. Patients must be screened and interviewed prior to the examination to exclude individuals who may be at risk by exposure to the MR environment.

Certain indications may require administration of intravenous (IV) contrast media. IV contrast enhancement should be performed using appropriate injection protocols and in accordance with the institution's policy on IV contrast use (see the <u>ACR–SPR Practice Parameter for the Use of Intravascular Contrast Media</u> [206]). Pediatric patients or patients suffering from anxiety or claustrophobia may require sedation or additional assistance. Administration of moderate sedation or general anesthesia may be needed to achieve a successful examination, particularly in young children. If moderate sedation is necessary, refer to the <u>ACR–SIR Practice Parameter for Moderate Sedation/Analgesia</u> [207].

B. Facility Requirements

Appropriate emergency equipment and medications must be immediately available to treat adverse reactions associated with administered medications. The equipment and medications should be monitored for inventory and drug expiration dates on a regular basis. The equipment, medications, and other emergency support must also be appropriate for the range of ages and sizes in the patient population.

C. Examination Technique

Diagnostic-quality hip and pelvis MRI can be performed with low-, medium-, or high-field systems of either closed-bore or open design. High-field magnets (1.5T and higher) have inherently better signal-to-noise ratios (SNRs) than lower-field systems, providing greater flexibility to obtain high-resolution images in a reasonable amount of time. However, there are circumstances in which lower field strength may be advantageous. These situations include imaging around metallic implants like prostheses and screws [13,208] and imaging in pregnant patients to reduce energy deposition in the fetus [201] (for more information, refer to the ACR Practice Parameter for Performing and Interpreting Magnetic Resonance Imaging (MRI) [202] and the ACR Guidance Document on MR Safe Practices 2020 [203]). Three-tesla (3T) MRI systems have higher SNRs and may offer improvements in assessment of cartilage and functional imaging [209-212].

Although an initial screen for abnormalities may use a body coil [39,62,213,214], high-resolution images require the use of a local coil. Multicoil arrays work best when imaging the entire pelvis and both hips [215,216]. When detailed images of a single hip or proximal femur are needed, several coil choices are available, ranging in configuration from flexible single coils [78,217] to paired loop-gap designs to commercially available or custom-built phased arrays [77,194,218-220].

Patients are typically positioned supine. The feet may be internally rotated and gently immobilized with tape, if necessary. Slight flexion at the knees achieved with padding may be more comfortable for some patients.

Coronal images are a mainstay of pelvic and hip MRI, and coronal T1-weighted and short-tau inversion recovery (STIR) images alone can rapidly screen for fractures [11,78,80] or femoral head osteonecrosis [62], but a complete examination should also include images in at least 1 additional imaging plane. Coronal and transverse images constitute a minimum examination for most indications [89,142,147,221]. The addition of sagittal plane images is useful for quantifying the extent of femoral head osteonecrosis [58,63,222], evaluating the hip joint cartilage and acetabular roof [85,223], investigating abnormalities of the proximal hamstring muscles and tendons [129,130], and assessing anterosuperior labral tears [200,201]. Sagittal images are essential and constitute the most important imaging plane to evaluate labrum and chondrolabral separation in the setting of common conditions of femoroacetabular impingement and hip dysplasia. Additionally, the standard imaging planes may be altered for specific indications. Sensitivity for detection of labral tears with large field of view (FOV) is poor, and small FOV images and arthrography

should be performed for evaluation of labrum [118].

Oblique coronal and transverse images angled parallel to the upper sacrum are useful to evaluate the sacroiliac joints [165,177], whereas images in either direct or oblique coronal and transverse planes can image the sacral plexus [157,224]. Selective use of oblique images along one femoral neck may assist in the diagnosis of subtle fractures [214,217,225] and labral tears [119,226] and in the evaluation of the femoral head-neck junction in femoroacetabular impingement syndrome [101]. Axial oblique images obtained parallel to the arcuate line of the pelvic inlet using a sagittal localizer can provide improved delineation of symphysis pubis anatomy in cases of groin pain or pubalgia [146]. In some practices, radial images acquired either directly [227,228] or via multiplanar reformatting of a volumetric data set [103,116,229,230] may assist the diagnosis of labral and cartilage pathology and potentially provide a more sensitive assessment of femoral head asphericity in cam-type femoroacetabular impingement [103,229], but whether the accuracy for diagnosing labral tears in such patients is improved compared with conventional imaging planes has not been determined [227]. Typically, 3 imaging planes are used for MR arthrography of the hip [114]. These may be oriented orthogonal to the pelvis or parallel to the femoral neck and perpendicular to the acetabular face for evaluating the acetabular labrum and hip joint capsule [231]. 3-D imaging may be useful for delineation and follow-up of femoral head and acetabular cartilage abnormalities [232-234]. 3-D imaging is best performed using isotropic spin-echo technique with prudent use of parallel imaging, and it allows radial reconstructions of hip and labral-cartilage assessment in multiple planes [235]. The pain response to the anesthetic in the arthrogram should be recorded and can provide valuable additional information to the orthopedic surgeon in addition to the MR arthrography findings, with respect to intra- or extra-articular origin of hip pain [236].

The FOV should be tailored to the size of the patient and the structures being examined. To screen the entire pelvis, a 35- to 45-cm FOV is typical, and the images should include enough tissue laterally to encompass the gluteal insertions and trochanteric bursae [123]. Images that include the entire pelvis are useful for making side-to-side comparisons [214]. Even when symptoms are unilateral, it may be advantageous to include at least 1 sequence with a large enough FOV to detect contralateral disease, which is frequently present [86]. Hip or groin pain may originate from several different anatomic structures, and a larger FOV may be required for evaluating the source of pain [146]. Larger FOV images can also frequently detect clinically occult pathology in patients with suspected proximal femur fractures [11]. For screening purposes, 6- to 8-mm-thick sections are adequate. However, higher imaging resolution is necessary to distinguish femoral head osteonecrosis from transient marrow conditions [66], to demonstrate subtle fracture lines, and to quantify the extent of osteonecrosis [58]. High-resolution imaging can be accomplished with a relatively large FOV if thin slices and a high imaging matrix are used (for example, 3- to 4-mm slice thickness and a 512 × 512 matrix) [58], or it can be accomplished by reducing the FOV to 16 to 20 cm and imaging each hip separately [77,237]. MR arthrography for labral or articular cartilage disease often requires even higher spatial resolution, with a small FOV to cover just 1 hip (typically 15 to 22 cm), thin sections (1.5 to 3 mm), and a relatively high matrix (256 phase steps or more) [149,231]. There is generally a trade-off between spatial resolution and imaging time [215]. Parallel imaging, which is available on some MR systems, allows faster image acquisition without substantial loss in image quality [238]. Parallel imaging can also mitigate undesired energy deposition on higher-field MRI systems like 3T scanners [239]. An interslice gap can increase coverage [240] but should be as small as feasible in order not to impair visualization of the imaged structures.

A wide variety of pulse sequences are available to image the pelvis and hips [241]. The choice of sequences, like other aspects of the imaging protocol, can be tailored to optimize the examination to answer specific clinical questions [39] and may vary because of local preferences. Short transition time/echo time (TR/TE) (T1-weighted) images are typically obtained using conventional spin-echo (SE) or fast (turbo) spin-echo (FSE) sequences. Long TR/TE (T2-weighted) and STIR images are frequently obtained using FSE techniques for more rapid image acquisition than SE technique allows [64]. A common higher-resolution 2-D imaging protocol for evaluation of internal derangements includes multiplanar intermediate-weighted fat-suppressed and nonfat-suppressed imaging, whereas T1W imaging is reserved specifically for infection and tumor protocols. Dixon imaging is another option that can be used to obtain multiple contrasts, including in-phase, opposed phase, and fat and water maps, which allow combined assessment of internal derangement and marrow assessment in the same setting [242]. Gradient-recalled sequences tend to produce larger artifacts and result in lower soft-tissue contrast [214] but may be advantageous at lower field strengths [243] and for selected applications, such as demonstration of hemosiderin in hips affected by pigmented villonodular synovitis [167] or evaluation of articular cartilage [150,228]. Gradient-echo and

modified fast (turbo) SE sequences can also be acquired as a 3-D volume, which is partitioned into contiguous thin sections or nonorthogonal imaging planes. Gradient-echo imaging results in suboptimal contrast and resolution as compared with FSE sequences. With current scanners, SE sequence with isotropic resolution can be obtained in 6-7 minutes of scanning time.

An imaging protocol will be composed of 1 or more pulse sequence types. For each sequence, the exact TR, TE, TI (inversion time), and flip angle chosen will depend on the field strength of the magnet and the desired contrast weighting. A typical minimal MRI examination of the pelvis and hips might consist of coronal SE or FSE T1-weighted and fat-suppressed, FSE T2-weighted or STIR images, and transverse T1weighted and T2-weighted sequences [147,214,221]. The T1-weighted images optimally show anatomic details such as fracture lines and bone marrow assessment [58,78,85,87], whereas T2-weighted or STIR images demonstrate fluid collections and edema within the soft tissue and bone marrow [85,175,213]; the combination is an effective screen for a variety of hip and pelvic pathologies [83,142,221]. T1-weighted sequences also have a role in characterizing various stages of hemorrhage [244,245] and muscle pathology [246,247] and in showing enhancement when gadolinium-based IV contrast agents are used [248]. T1weighted images with fat suppression—either 2-D SE or FSE [114,149] or 3-D spoiled gradient-echo [230,231]—are also used when MR arthrography is performed with a gadolinium-based contrast agent. At least 1 T2-weighted sequence should also be performed with MR arthrograms to show additional abnormalities affecting the hip (eg, stress fractures, tendon/muscle injuries, bursitis, and cystic structures that do not communicate with the joint). Additionally, at least 1 T1-weighted sequence without fat suppression is useful for evaluating bone marrow and characterizing soft-tissue lesions. Suppressing the signal from fat may enhance the diagnostic yield of some pulse sequences [241]. Fat

suppression can use spectral suppression of water protons, a phase-dependent method such as the Dixon technique, or a STIR sequence [249,250]. The latter 2 methods may be necessary on low-field systems [251]. Fat suppression increases the conspicuity of marrow abnormalities and soft-tissue edema on fluid-sensitive sequences [4,214] and is useful with a T1-weighted sequence when using gadolinium-based contrast agents such as in MR arthrographic evaluation of hip cartilage or labral tears [104,119]. Selective excitation of water protons is an alternative to fat suppression and has been investigated for evaluating the hip articular cartilage [223].

For specific hip and pelvis disorders, IV contrast may be useful. Contrast enhancement, especially on dynamic contrast-enhanced imaging, suggests femoral head viability in Legg-Calvé-Perthes disease [252] and femoral neck fractures [183,184], and may detect early ischemia [253] and predict future risk of osteonecrosis after closed reduction of developmental hip dysplasia [67]. IV contrast can also aid in the diagnosis of hip joint synovitis [160,164,165], pelvic infections [159], tendon degeneration [145], and tumors, and it may play a role in the evaluation of the interface surrounding hip prosthesis components [13]. MR arthrography is beneficial for evaluating internal hip derangements [33,37] and sports injury [88]. The MR diagnosis of labral, articular cartilage, and joint capsule abnormalities in the hip is enhanced by the addition of intra-articular contrast [97,112,231,254]. Conventional 3T MRI is equivalent or superior to 1.5T MR arthrography for labral tears and cartilage defects. Sensitivity of labral tears with 3T MRI is similar to 3T MR arthrography [255]. For the hip joint, MR arthrography is usually performed following direct imaging-guided, intra-articular injection of dilute gadolinium-based contrast or saline. Adding leg traction has been reported to improve delineation of the cartilage surfaces [150,256] and acetabular labrum [254] during MR arthrography but is not commonly used in clinical practice because of logistical issues.

Although indirect MR arthrography is also possible for hip imaging, because of the size of the joint, a delay after IV contrast administration is necessary to allow adequate contrast diffusion into the joint [254]. Indirect MR arthrography may be more sensitive for labral tears but may not improve diagnosis of chondral abnormalities compared with nonarthrographic MRI [120].

Delayed gadolinium-enhanced MRI of cartilage (dGEMRIC) is a technique for assessing cartilage integrity in osteoarthritis and other hip diseases that result in cartilage damage. It requires the use of an anionically charged gadolinium contrast agent, typically administered intravenously [5,109,110,257], but can also be performed after intra-articular injection [258]. dGEMRIC has been used to diagnose and stage osteoarthritis in patients with femoroacetabular impingement [5] and for preosteotomy evaluation of patients with hip dysplasia [7,92]. Additional MR techniques for cartilage assessment have been tested in patients with hip diseases. T2 mapping has been used to evaluate cartilage in hip dysplasia patients [259]. Leg traction may improve delineation of the cartilage surfaces [150,256] and acetabular labrum [254] during MR arthrography. T1-rho may be a useful imaging technique in the detection of early cartilage degeneration [260,261].

Various techniques are used to reduce artifacts that can reduce imaging quality. When the FOV excludes parts of the pelvis that are within the sensitivity range of the coil (eg, when imaging a single hip), aliasing artifacts can be reduced by phase oversampling or by orienting the phase-encoding direction along the anteroposterior axis [252]. Ensuring patient comfort combined with gentle immobilization when necessary best controls involuntary patient motion [241]. Presaturation pulses and/or gradient moment nulling will reduce ghosting artifacts caused by flowing blood [262,263]. Imaging near metallic implants requires special care to reduce susceptibility to artifacts. Orienting the long axis of the implant along the frequencyencoding gradient [128,195,218,264], avoiding gradient-recalled sequences [195], and substituting STIR for chemical fat suppression [13,127,195,198,265] are important considerations. FSE sequences with short interecho spacing, multiple refocusing pulses (long-echo trains), and tailored RF pulses will further minimize metallic artifacts [127,194,196,218,264,266]. Metal artifact is further reduced by using a wide readout bandwidth and small pixel dimensions, which may require more signals averaged to maintain an adequate SNR and are better performed using MR systems with wider available receiver bandwidths [13,194,198,218,264]. Progress in metal artifact reduction techniques minimize metal-related artifacts while improving depiction of synovium and bone-implant interfaces [267,268]. Such techniques provide a diagnostically meaningful reduction of metal artifact around implants and can be used as an adjunct to optimized 2-D FSE sequences [267]. Lastly, artifacts from metal implants are less prominent on low-field systems compared with high-field systems [13].

It is the responsibility of the supervising physician to determine whether additional or unconventional pulse sequences or imaging techniques would confer added benefit for the diagnosis and management of the patient. Examinations that use techniques not approved by the Food and Drug Administration—such as the intra-articular injection of gadolinium chelates (direct MR arthrography) [269]—should be considered only when they are judged to be medically appropriate.

VI. DOCUMENTATION

Reporting should be in accordance with the <u>ACR Practice Parameter for Communication of Diagnostic Imaging</u> <u>Findings</u> [270].

At a minimum, the report should address any abnormalities in the bone marrow, soft tissues, and joints. In selected cases, a description of findings in specific muscles and tendons, articular cartilage, fibrocartilage, synovium, neurovascular structures, lymph nodes, cortical bone, and surrounding bursae would be appropriate. For MR arthrograms of the hip, the report should also specifically indicate the condition of the acetabular labrum and articular cartilage. Whenever possible, the report should use standard anatomic nomenclature and precise terms and anatomic localization for describing identified abnormalities.

Specific policies and procedures related to MRI safety should be in place along with documentation that is updated annually and compiled under the supervision and direction of the supervising MRI physician. Guidelines should be provided that deal with potential hazards associated with the MRI examination of the patient as well as to others in the immediate area. [204,205,271]. Screening forms must also be provided to detect those patients who may be at risk for adverse events associated with the MRI examination [272].

VII. EQUIPMENT SPECIFICATIONS

Equipment monitoring should be in accordance with the <u>ACR–AAPM Technical Standard for Diagnostic Medical</u> <u>Physics Performance Monitoring of Magnetic Resonance Imaging (MRI) Equipment</u> [273].

The MRI equipment specifications and performance must meet all state and federal requirements. The requirements include, but are not limited to, specifications of maximum static magnetic strength, maximum rate of change of the magnetic field strength (dB/dt), maximum radiofrequency power deposition (specific absorption rate), and maximum acoustic noise levels.

VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection

Control, and Patient Education appearing under the heading ACR Position Statement on Quality Control & Improvement, Safety, Infection Control, and Patient Education on the ACR website (https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement).

ACKNOWLEDGEMENTS

This practice parameter was revised according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (<u>https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards</u>) by the Committee on Practice Parameters – Body Imaging (Musculoskeletal) of the ACR Commission on Body Imaging and the Committee on Practice Parameters – Pediatric Radiology of the ACR Commission on Pediatric Radiology, in collaboration with the SPR and the SSR.

Writing Committee – members represent their societies in the initial and final revision of this practice parameter

ACR	<u>SPR</u>		<u>SSR</u>	
Mary K. Jesse, MD, Chair	Kirsten Ecklund	d, MD	Miriam A. Bredella, MD	
Lauren Golding, MD	Kathleen Emery, MD			
Elaine S. Gould, MD, FACR				
Jason Higgins, MD				
Committee on Body Imaging (Musculoskeletal)				
(ACR Committee responsible for sponsoring the draft through the process)				
Catherine C. Roberts, MD, Chair		Suzanne S. Long, MD		
Jeffrey M. Brody, MD, FACR		Kambiz Motamedi, M	ID	
Bethany U. Casagranda, DO		Carlos A. Rivera, BSc		
Elaine S Gould, MD, FACR		Aleksandr Rozenberg	, MD	
Mary K. Jesse, MD		Naveen Subhas, MD		
Kenneth S. Lee, MD				

<u>Committee on Practice Parameters – Pediatric Radiology</u>

(ACR Committee responsible for sponsoring the draft through the process)

Terry L. Levin, MD, FACR, Chair	Jane Sun Kim, MD		
John B. Amodio, MD, FACR	Jennifer A Knight, MD		
Jesse Berman, MD	Jessica Kurian, MD		
Tara M. Catanzano, MB, BCh	Matthew P. Lungren, MD, MPH		
Harris L. Cohen, MD, FACR	Helen R. Nadel, MD		
Kassa Darge, MD, PhD	Erica Poletto, MD		
Dorothy L. Gilbertson-Dahdal, MD	Richard B. Towbin, MD, FACR		
Lauren P. Golding, MD	Andrew T. Trout, MD		
Safwan S. Halabi, MD	Esben S. Vogelius, MD		
Jason Higgins, DO			
Andrew B. Rosenkrantz, MD, Chair, Commission on Body Imaging			
Richard A. Barth, MD, FACR, Chair, Commission on Pediatric Radiology			
David B. Larson, MD, MBA, Chair, Commission on Quality and Safety			
Mary S. Newell, MD, FACR, Chair, Committee on Prac	ctice Parameters and Technical Standards		
Comment Reconciliation Committee			
Rachel Gerson, MD, Chair	Mary K. Jesse, MD		
Adam Specht, MD, FACR, Co-Chair	Amy Kotsenas, MD, FACR		

Richard A. Barth, MD, FACR

David B. Larson, MD, MBA

Comment Reconciliation Committee

Miriam A. Bredella, MD	Paul A. Larson, MD, FACR
Avneesh Chhabra, MD	Terry L. Levin, MD, FACR
Richard Duszak Jr., MD, FACR	Mary S. Newell, MD, FACR
Kirsten Ecklund, MD	Catherine C. Roberts, MD
Kathleen Emery, MD	Andrew B. Rosenkrantz, MD
Lauren Golding, MD	Michael Ian Rothman, MD, FACR
Elaine S. Gould, MD, FACR	Susanna C Spence, MD
Jason Higgins, MD	Naveen Subhas, MD

REFERENCES

- 1. American College of Radiology. ACR–SAR–SPR practice parameter for the performance of magnetic resonance imaging (MRI) of the soft-tissue components of the pelvis. Available at: <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-SoftTissue-Pel.pdf</u>. Accessed January 14, 2020.
- 2. Bassounas AE, Karantanas AH, Fotiadis DI, Malizos KN. Femoral head osteonecrosis: volumetric MRI assessment and outcome. Eur J Radiol 2007;63:10-5.
- 3. Ueshima K, Takahashi KA, Fujioka M, et al. Relationship between acetabular labrum evaluation by using radial magnetic resonance imaging and progressive joint space narrowing in mild hip dysplasia. Magn Reson Imaging 2006;24:645-50.
- 4. Bennett AN, McGonagle D, O'Connor P, et al. Severity of baseline magnetic resonance imaging-evident sacroiliitis and HLA-B27 status in early inflammatory back pain predict radiographically evident ankylosing spondylitis at eight years. Arthritis Rheum 2008;58:3413-8.
- 5. Jessel RH, Zilkens C, Tiderius C, Dudda M, Mamisch TC, Kim YJ. Assessment of osteoarthritis in hips with femoroacetabular impingement using delayed gadolinium enhanced MRI of cartilage. J Magn Reson Imaging 2009;30:1110-5.
- 6. Zibis AH, Karantanas AH, Roidis NT, et al. The role of MR imaging in staging femoral head osteonecrosis. Eur J Radiol 2007;63:3-9.
- 7. Cunningham T, Jessel R, Zurakowski D, Millis MB, Kim YJ. Delayed gadolinium-enhanced magnetic resonance imaging of cartilage to predict early failure of Bernese periacetabular osteotomy for hip dysplasia. J Bone Joint Surg Am 2006;88:1540-8.
- Neumann G, Mendicuti AD, Zou KH, et al. Prevalence of labral tears and cartilage loss in patients with mechanical symptoms of the hip: evaluation using MR arthrography. Osteoarthritis Cartilage 2007;15:909-17.
- 9. Zalavras CG, Rigopoulos N, Lee J, Learch T, Patzakis MJ. Magnetic resonance imaging findings in hematogenous osteomyelitis of the hip in adults. Clin Orthop Relat Res 2009;467:1688-92.
- 10. Meyers WC, McKechnie A, Philippon MJ, Horner MA, Zoga AC, Devon ON. Experience with "sports hernia" spanning two decades. Ann Surg 2008;248:656-65.
- 11. Sankey RA, Turner J, Lee J, Healy J, Gibbons CE. The use of MRI to detect occult fractures of the proximal

femur: a study of 102 consecutive cases over a ten-year period. J Bone Joint Surg Br 2009;91:1064-8.

- 12. Newberg AH, Newman JS. Imaging the painful hip. Clin Orthop Relat Res 2003:19-28.
- 13. Sugimoto H, Hirose I, Miyaoka E, et al. Low-field-strength MR imaging of failed hip arthroplasty: association of femoral periprosthetic signal intensity with radiographic, surgical, and pathologic findings. Radiology 2003;229:718-23.
- 14. Garcia-Morales F, Seo GS, Chengazi V, Monu JU. Collar osteophytes: a cause of false-positive findings in bone scans for hip fractures. AJR Am J Roentgenol 2003;181:191-4.
- 15. Holder LE, Schwarz C, Wernicke PG, Michael RH. Radionuclide bone imaging in the early detection of fractures of the proximal femur (hip): multifactorial analysis. Radiology 1990;174:509-15.
- 16. Song WS, Yoo JJ, Koo KH, Yoon KS, Kim YM, Kim HJ. Subchondral fatigue fracture of the femoral head in military recruits. J Bone Joint Surg Am 2004;86-A:1917-24.
- 17. Feldman F, Staron RB. MRI of seemingly isolated greater trochanteric fractures. AJR Am J Roentgenol 2004;183:323-9.
- Rubin SJ, Marquardt JD, Gottlieb RH, Meyers SP, Totterman SM, O'Mara RE. Magnetic resonance imaging: a cost-effective alternative to bone scintigraphy in the evaluation of patients with suspected hip fractures. Skeletal Radiol 1998;27:199-204.
- 19. Shin AY, Morin WD, Gorman JD, Jones SB, Lapinsky AS. The superiority of magnetic resonance imaging in differentiating the cause of hip pain in endurance athletes. Am J Sports Med 1996;24:168-76.
- 20. Stumpe KD, Notzli HP, Zanetti M, et al. FDG PET for differentiation of infection and aseptic loosening in total hip replacements: comparison with conventional radiography and three-phase bone scintigraphy. Radiology 2004;231:333-41.
- 21. Palestro CJ. Nuclear medicine and the failed joint replacement: Past, present, and future. World J Radiol 2014;6:446-58.
- 22. Connell DA, Schneider-Kolsky ME, Hoving JL, et al. Longitudinal study comparing sonographic and MRI assessments of acute and healing hamstring injuries. AJR Am J Roentgenol 2004;183:975-84.
- 23. Wahl CJ, Warren RF, Adler RS, Hannafin JA, Hansen B. Internal coxa saltans (snapping hip) as a result of overtraining: a report of 3 cases in professional athletes with a review of causes and the role of ultrasound in early diagnosis and management. Am J Sports Med 2004;32:1302-9.
- 24. Wunderbaldinger P, Bremer C, Matuszewski L, Marten K, Turetschek K, Rand T. Efficient radiological assessment of the internal snapping hip syndrome. Eur Radiol 2001;11:1743-7.
- 25. Sudula SN. Imaging the hip joint in osteoarthritis: A place for ultrasound? Ultrasound 2016;24:111-8.
- 26. Lungu E, Michaud J, Bureau NJ. US Assessment of Sports-related Hip Injuries. Radiographics 2018;38:867-89.
- 27. Wunderbaldinger P, Bremer C, Schellenberger E, Cejna M, Turetschek K, Kainberger F. Imaging features of iliopsoas bursitis. Eur Radiol 2002;12:409-15.
- 28. Deslandes M, Guillin R, Cardinal E, Hobden R, Bureau NJ. The snapping iliopsoas tendon: new mechanisms using dynamic sonography. AJR Am J Roentgenol 2008;190:576-81.
- 29. Choi YS, Lee SM, Song BY, Paik SH, Yoon YK. Dynamic sonography of external snapping hip syndrome. Journal of ultrasound in medicine : official journal of the American Institute of Ultrasound in Medicine 2002;21:753-8.
- 30. Pisacano RM, Miller TT. Comparing sonography with MR imaging of apophyseal injuries of the pelvis in four boys. AJR Am J Roentgenol 2003;181:223-30.
- Muresan S, Marginean MO, Voidazan S, Vlasa I, Sintean I. Musculoskeletal ultrasound: a useful tool for diagnosis of hip developmental dysplasia: One single-center experience. Medicine (Baltimore) 2019;98:e14081.
- 32. Martin RL, Irrgang JJ, Sekiya JK. The diagnostic accuracy of a clinical examination in determining intraarticular hip pain for potential hip arthroscopy candidates. Arthroscopy 2008;24:1013-8.
- 33. Byrd JW, Jones KS. Diagnostic accuracy of clinical assessment, magnetic resonance imaging, magnetic resonance arthrography, and intra-articular injection in hip arthroscopy patients. Am J Sports Med 2004;32:1668-74.
- 34. Schilders E, Bismil Q, Robinson P, O'Connor PJ, Gibbon WW, Talbot JC. Adductor-related groin pain in competitive athletes. Role of adductor enthesis, magnetic resonance imaging, and entheseal pubic cleft injections. J Bone Joint Surg Am 2007;89:2173-8.
- 35. Schilders E, Talbot JC, Robinson P, Dimitrakopoulou A, Gibbon WW, Bismil Q. Adductor-related groin pain in

recreational athletes: role of the adductor enthesis, magnetic resonance imaging, and entheseal pubic cleft injections. J Bone Joint Surg Am 2009;91:2455-60.

- 36. Kelly BT, Williams RJ, 3rd, Philippon MJ. Hip arthroscopy: current indications, treatment options, and management issues. Am J Sports Med 2003;31:1020-37.
- 37. McCarthy JC, Lee JA. Arthroscopic intervention in early hip disease. Clin Orthop Relat Res 2004:157-62.
- 38. Brooks RA, Ribbans WJ. Diagnosis and imaging studies of traumatic hip dislocations in the adult. Clin Orthop Relat Res 2000:15-23.
- 39. Conway WF, Totty WG, McEnery KW. CT and MR imaging of the hip. Radiology 1996;198:297-307.
- 40. Potter HG, Montgomery KD, Heise CW, Helfet DL. MR imaging of acetabular fractures: value in detecting femoral head injury, intraarticular fragments, and sciatic nerve injury. AJR Am J Roentgenol 1994;163:881-6.
- 41. Garbuz DS, Masri BA, Haddad F, Duncan CP. Clinical and radiographic assessment of the young adult with symptomatic hip dysplasia. Clin Orthop Relat Res 2004:18-22.
- 42. Ecker TM, Puls M, Steppacher SD, et al. Computer-assisted femoral head-neck osteochondroplasty using a surgical milling device an in vitro accuracy study. J Arthroplasty 2012;27:310-6.
- 43. Su AW, Hillen TJ, Eutsler EP, et al. Low-Dose Computed Tomography Reduces Radiation Exposure by 90% Compared With Traditional Computed Tomography Among Patients Undergoing Hip-Preservation Surgery. Arthroscopy 2019;35:1385-92.
- 44. Claus AM, Totterman SM, Sychterz CJ, Tamez-Pena JG, Looney RJ, Engh CA, Sr. Computed tomography to assess pelvic lysis after total hip replacement. Clin Orthop Relat Res 2004:167-74.
- 45. Park JS, Ryu KN, Hong HP, Park YK, Chun YS, Yoo MC. Focal osteolysis in total hip replacement: CT findings. Skeletal Radiol 2004;33:632-40.
- 46. Puri L, Wixson RL, Stern SH, Kohli J, Hendrix RW, Stulberg SD. Use of helical computed tomography for the assessment of acetabular osteolysis after total hip arthroplasty. J Bone Joint Surg Am 2002;84-A:609-14.
- 47. Mitchell MD, Kundel HL, Steinberg ME, Kressel HY, Alavi A, Axel L. Avascular necrosis of the hip: comparison of MR, CT, and scintigraphy. AJR Am J Roentgenol 1986;147:67-71.
- 48. Stevens K, Tao C, Lee SU, et al. Subchondral fractures in osteonecrosis of the femoral head: comparison of radiography, CT, and MR imaging. AJR Am J Roentgenol 2003;180:363-8.
- 49. Cabarrus MC, Ambekar A, Lu Y, Link TM. MRI and CT of insufficiency fractures of the pelvis and the proximal femur. AJR Am J Roentgenol 2008;191:995-1001.
- 50. Lubovsky O, Liebergall M, Mattan Y, Weil Y, Mosheiff R. Early diagnosis of occult hip fractures MRI versus CT scan. Injury 2005;36:788-92.
- 51. Fon LJ, Spence RA. Sportsman's hernia. Br J Surg 2000;87:545-52.
- 52. Kozin F, Carrera GF, Ryan LM, Foley D, Lawson T. Computed tomography in the diagnosis of sacroiliitis. Arthritis Rheum 1981;24:1479-85.
- 53. Nishii T, Tanaka H, Nakanishi K, Sugano N, Miki H, Yoshikawa H. Fat-suppressed 3D spoiled gradient-echo MRI and MDCT arthrography of articular cartilage in patients with hip dysplasia. AJR Am J Roentgenol 2005;185:379-85.
- 54. Yamamoto Y, Tonotsuka H, Ueda T, Hamada Y. Usefulness of radial contrast-enhanced computed tomography for the diagnosis of acetabular labrum injury. Arthroscopy 2007;23:1290-4.
- 55. Tobalem F, Dugert E, Verdun FR, et al. MDCT arthrography of the hip: value of the adaptive statistical iterative reconstruction technique and potential for radiation dose reduction. AJR Am J Roentgenol 2014;203:W665-73.
- 56. Beltran J, Knight CT, Zuelzer WA, et al. Core decompression for avascular necrosis of the femoral head: correlation between long-term results and preoperative MR staging. Radiology 1990;175:533-6.
- 57. Cheng EY, Thongtrangan I, Laorr A, Saleh KJ. Spontaneous resolution of osteonecrosis of the femoral head. J Bone Joint Surg Am 2004;86-A:2594-9.
- 58. Cherian SF, Laorr A, Saleh KJ, Kuskowski MA, Bailey RF, Cheng EY. Quantifying the extent of femoral head involvement in osteonecrosis. J Bone Joint Surg Am 2003;85-A:309-15.
- 59. Coleman BG, Kressel HY, Dalinka MK, Scheibler ML, Burk DL, Cohen EK. Radiographically negative avascular necrosis: detection with MR imaging. Radiology 1988;168:525-8.
- 60. Huang GS, Chan WP, Chang YC, Chang CY, Chen CY, Yu JS. MR imaging of bone marrow edema and joint effusion in patients with osteonecrosis of the femoral head: relationship to pain. AJR Am J Roentgenol 2003;181:545-9.
- 61. Iida S, Harada Y, Shimizu K, et al. Correlation between bone marrow edema and collapse of the femoral

head in steroid-induced osteonecrosis. AJR Am J Roentgenol 2000;174:735-43.

- 62. Khanna AJ, Yoon TR, Mont MA, Hungerford DS, Bluemke DA. Femoral head osteonecrosis: detection and grading by using a rapid MR imaging protocol. Radiology 2000;217:188-92.
- 63. Koo KH, Kim R. Quantifying the extent of osteonecrosis of the femoral head. A new method using MRI. J Bone Joint Surg Br 1995;77:875-80.
- 64. May DA, Disler DG. Screening for avascular necrosis of the hip with rapid MRI: preliminary experience. J Comput Assist Tomogr 2000;24:284-7.
- 65. Radke S, Kirschner S, Seipel V, Rader C, Eulert J. Magnetic resonance imaging criteria of successful core decompression in avascular necrosis of the hip. Skeletal Radiol 2004;33:519-23.
- 66. Watson RM, Roach NA, Dalinka MK. Avascular necrosis and bone marrow edema syndrome. Radiol Clin North Am 2004;42:207-19.
- 67. Tiderius C, Jaramillo D, Connolly S, et al. Post-closed reduction perfusion magnetic resonance imaging as a predictor of avascular necrosis in developmental hip dysplasia: a preliminary report. J Pediatr Orthop 2009;29:14-20.
- 68. Ito H, Matsuno T, Minami A. Relationship between bone marrow edema and development of symptoms in patients with osteonecrosis of the femoral head. AJR Am J Roentgenol 2006;186:1761-70.
- 69. Piyakunmala K, Sangkomkamhang T, Chareonchonvanitch K. Is magnetic resonance imaging necessary for normal plain radiography evaluation of contralateral non-traumatic asymptomatic femoral head in high osteonecrosis risk patient. J Med Assoc Thai 2009;92 Suppl 6:S147-51.
- 70. Bloem JL. Transient osteoporosis of the hip: MR imaging. Radiology 1988;167:753-5.
- 71. Malizos KN, Zibis AH, Dailiana Z, Hantes M, Karachalios T, Karantanas AH. MR imaging findings in transient osteoporosis of the hip. Eur J Radiol 2004;50:238-44.
- 72. Neuhold A, Hofmann S, Engel A, et al. Bone marrow edema of the hip: MR findings after core decompression. J Comput Assist Tomogr 1992;16:951-5.
- 73. Vande Berg BC, Malghem JJ, Lecouvet FE, Jamart J, Maldague BE. Idiopathic bone marrow edema lesions of the femoral head: predictive value of MR imaging findings. Radiology 1999;212:527-35.
- 74. Gorbachova T, Amber I, Beckmann NM, et al. Nomenclature of Subchondral Nonneoplastic Bone Lesions. AJR Am J Roentgenol 2019;213:963-82.
- 75. Deutsch AL, Mink JH, Waxman AD. Occult fractures of the proximal femur: MR imaging. Radiology 1989;170:113-6.
- 76. Haramati N, Staron RB, Barax C, Feldman F. Magnetic resonance imaging of occult fractures of the proximal femur. Skeletal Radiol 1994;23:19-22.
- 77. Oka M, Monu JU. Prevalence and patterns of occult hip fractures and mimics revealed by MRI. AJR Am J Roentgenol 2004;182:283-8.
- 78. Quinn SF, McCarthy JL. Prospective evaluation of patients with suspected hip fracture and indeterminate radiographs: use of T1-weighted MR images. Radiology 1993;187:469-71.
- 79. Rizzo PF, Gould ES, Lyden JP, Asnis SE. Diagnosis of occult fractures about the hip. Magnetic resonance imaging compared with bone-scanning. J Bone Joint Surg Am 1993;75:395-401.
- 80. Frihagen F, Nordsletten L, Tariq R, Madsen JE. MRI diagnosis of occult hip fractures. Acta Orthop 2005;76:524-30.
- 81. Hossain M, Barwick C, Sinha AK, Andrew JG. Is magnetic resonance imaging (MRI) necessary to exclude occult hip fracture? Injury 2007;38:1204-8.
- 82. Verbeeten KM, Hermann KL, Hasselqvist M, et al. The advantages of MRI in the detection of occult hip fractures. Eur Radiol 2005;15:165-9.
- 83. Ahovuo JA, Kiuru MJ, Visuri T. Fatigue stress fractures of the sacrum: diagnosis with MR imaging. Eur Radiol 2004;14:500-5.
- 84. Blomlie V, Rofstad EK, Talle K, Sundfor K, Winderen M, Lien HH. Incidence of radiation-induced insufficiency fractures of the female pelvis: evaluation with MR imaging. AJR Am J Roentgenol 1996;167:1205-10.
- 85. Grangier C, Garcia J, Howarth NR, May M, Rossier P. Role of MRI in the diagnosis of insufficiency fractures of the sacrum and acetabular roof. Skeletal Radiol 1997;26:517-24.
- 86. Kiuru MJ, Pihlajamaki HK, Ahovuo JA. Fatigue stress injuries of the pelvic bones and proximal femur: evaluation with MR imaging. Eur Radiol 2003;13:605-11.
- 87. Otte MT, Helms CA, Fritz RC. MR imaging of supra-acetabular insufficiency fractures. Skeletal Radiol 1997;26:279-83.

- 88. Overdeck KH, Palmer WE. Imaging of hip and groin injuries in athletes. Semin Musculoskelet Radiol 2004;8:41-55.
- 89. Slocum KA, Gorman JD, Puckett ML, Jones SB. Resolution of abnormal MR signal intensity in patients with stress fractures of the femoral neck. AJR Am J Roentgenol 1997;168:1295-9.
- 90. Hebert KJ, Laor T, Divine JG, Emery KH, Wall EJ. MRI appearance of chronic stress injury of the iliac crest apophysis in adolescent athletes. AJR Am J Roentgenol 2008;190:1487-91.
- 91. Pecina HI, Boric I, Smoljanovic T, Duvancic D, Pecina M. Surgical evaluation of magnetic resonance imaging findings in piriformis muscle syndrome. Skeletal Radiol 2008;37:1019-23.
- 92. Patel RN, Ashraf A, Sundaram M. Atypical Fractures Following Bisphosphonate Therapy. Semin Musculoskelet Radiol 2016;20:376-81.
- 93. Futami T, Suzuki S, Seto Y, Kashiwagi N. Sequential magnetic resonance imaging in slipped capital femoral epiphysis: assessment of preslip in the contralateral hip. J Pediatr Orthop B 2001;10:298-303.
- 94. Jaramillo D, Kasser JR, Villegas-Medina OL, Gaary E, Zurakowski D. Cartilaginous abnormalities and growth disturbances in Legg-Calve-Perthes disease: evaluation with MR imaging. Radiology 1995;197:767-73.
- 95. Kaniklides C, Lonnerholm T, Moberg A, Sahlstedt B. Legg-Calve-Perthes disease. Comparison of conventional radiography, MR imaging, bone scintigraphy and arthrography. Acta Radiol 1995;36:434-9.
- 96. Zilkens C, Holstein A, Bittersohl B, et al. Delayed gadolinium-enhanced magnetic resonance imaging of cartilage in the long-term follow-up after Perthes disease. J Pediatr Orthop 2010;30:147-53.
- 97. Leunig M, Podeszwa D, Beck M, Werlen S, Ganz R. Magnetic resonance arthrography of labral disorders in hips with dysplasia and impingement. Clin Orthop Relat Res 2004:74-80.
- 98. Laor T, Crawford AH. Idiopathic chondrolysis of the hip in children: early MRI findings. AJR Am J Roentgenol 2009;192:526-31.
- 99. Ito K, Leunig M, Ganz R. Histopathologic features of the acetabular labrum in femoroacetabular impingement. Clin Orthop Relat Res 2004:262-71.
- 100. Ito K, Minka MA, 2nd, Leunig M, Werlen S, Ganz R. Femoroacetabular impingement and the cam-effect. A MRI-based quantitative anatomical study of the femoral head-neck offset. J Bone Joint Surg Br 2001;83:171-6.
- 101. Notzli HP, Wyss TF, Stoecklin CH, Schmid MR, Treiber K, Hodler J. The contour of the femoral head-neck junction as a predictor for the risk of anterior impingement. J Bone Joint Surg Br 2002;84:556-60.
- 102. Schmid MR, Notzli HP, Zanetti M, Wyss TF, Hodler J. Cartilage lesions in the hip: diagnostic effectiveness of MR arthrography. Radiology 2003;226:382-6.
- 103. Rakhra KS, Sheikh AM, Allen D, Beaule PE. Comparison of MRI alpha angle measurement planes in femoroacetabular impingement. Clin Orthop Relat Res 2009;467:660-5.
- 104. Anderson LA, Peters CL, Park BB, Stoddard GJ, Erickson JA, Crim JR. Acetabular cartilage delamination in femoroacetabular impingement. Risk factors and magnetic resonance imaging diagnosis. J Bone Joint Surg Am 2009;91:305-13.
- 105. Pfirrmann CW, Duc SR, Zanetti M, Dora C, Hodler J. MR arthrography of acetabular cartilage delamination in femoroacetabular cam impingement. Radiology 2008;249:236-41.
- 106. Pfirrmann CW, Mengiardi B, Dora C, Kalberer F, Zanetti M, Hodler J. Cam and pincer femoroacetabular impingement: characteristic MR arthrographic findings in 50 patients. Radiology 2006;240:778-85.
- 107. Kassarjian A, Yoon LS, Belzile E, Connolly SA, Millis MB, Palmer WE. Triad of MR arthrographic findings in patients with cam-type femoroacetabular impingement. Radiology 2005;236:588-92.
- 108. James SL, Ali K, Malara F, Young D, O'Donnell J, Connell DA. MRI findings of femoroacetabular impingement. AJR Am J Roentgenol 2006;187:1412-9.
- 109. Bittersohl B, Hosalkar HS, Kim YJ, Werlen S, Siebenrock KA, Mamisch TC. Delayed gadolinium-enhanced magnetic resonance imaging (dGEMRIC) of hip joint cartilage in femoroacetabular impingement (FAI): Are pre- and postcontrast imaging both necessary? Magn Reson Med 2009;62:1362-7.
- 110. Bittersohl B, Steppacher S, Haamberg T, et al. Cartilage damage in femoroacetabular impingement (FAI): preliminary results on comparison of standard diagnostic vs delayed gadolinium-enhanced magnetic resonance imaging of cartilage (dGEMRIC). Osteoarthritis Cartilage 2009;17:1297-306.
- 111. Boutin RD, Newman JS. MR imaging of sports-related hip disorders. Magn Reson Imaging Clin N Am 2003;11:255-81.
- 112. Keeney JA, Peelle MW, Jackson J, Rubin D, Maloney WJ, Clohisy JC. Magnetic resonance arthrography versus arthroscopy in the evaluation of articular hip pathology. Clin Orthop Relat Res 2004:163-9.

- 113. Magee T, Hinson G. Association of paralabral cysts with acetabular disorders. AJR Am J Roentgenol 2000;174:1381-4.
- 114. Petersilge CA, Haque MA, Petersilge WJ, Lewin JS, Lieberman JM, Buly R. Acetabular labral tears: evaluation with MR arthrography. Radiology 1996;200:231-5.
- 115. Schnarkowski P, Steinbach LS, Tirman PF, Peterfy CG, Genant HK. Magnetic resonance imaging of labral cysts of the hip. Skeletal Radiol 1996;25:733-7.
- 116. Chan YS, Lien LC, Hsu HL, et al. Evaluating hip labral tears using magnetic resonance arthrography: a prospective study comparing hip arthroscopy and magnetic resonance arthrography diagnosis. Arthroscopy 2005;21:1250.
- 117. Mintz DN, Hooper T, Connell D, Buly R, Padgett DE, Potter HG. Magnetic resonance imaging of the hip: detection of labral and chondral abnormalities using noncontrast imaging. Arthroscopy 2005;21:385-93.
- 118. Toomayan GA, Holman WR, Major NM, Kozlowicz SM, Vail TP. Sensitivity of MR arthrography in the evaluation of acetabular labral tears. AJR Am J Roentgenol 2006;186:449-53.
- 119. Freedman BA, Potter BK, Dinauer PA, Giuliani JR, Kuklo TR, Murphy KP. Prognostic value of magnetic resonance arthrography for Czerny stage II and III acetabular labral tears. Arthroscopy 2006;22:742-7.
- 120. Zlatkin MB, Pevsner D, Sanders TG, Hancock CR, Ceballos CE, Herrera MF. Acetabular labral tears and cartilage lesions of the hip: indirect MR arthrographic correlation with arthroscopy--a preliminary study. AJR Am J Roentgenol 2010;194:709-14.
- 121. Chung CB, Robertson JE, Cho GJ, Vaughan LM, Copp SN, Resnick D. Gluteus medius tendon tears and avulsive injuries in elderly women: imaging findings in six patients. AJR Am J Roentgenol 1999;173:351-3.
- 122. Bird PA, Oakley SP, Shnier R, Kirkham BW. Prospective evaluation of magnetic resonance imaging and physical examination findings in patients with greater trochanteric pain syndrome. Arthritis Rheum 2001;44:2138-45.
- 123. Cvitanic O, Henzie G, Skezas N, Lyons J, Minter J. MRI diagnosis of tears of the hip abductor tendons (gluteus medius and gluteus minimus). AJR Am J Roentgenol 2004;182:137-43.
- 124. Kingzett-Taylor A, Tirman PF, Feller J, et al. Tendinosis and tears of gluteus medius and minimus muscles as a cause of hip pain: MR imaging findings. AJR Am J Roentgenol 1999;173:1123-6.
- 125. Kagan A, 2nd. Rotator cuff tears of the hip. Clin Orthop Relat Res 1999:135-40.
- 126. Demant AW, Kocovic L, Henschkowski J, et al. Hip pain in renal transplant recipients: symptomatic gluteus minimus and gluteus medius tendon abnormality as an alternative MRI diagnosis to avascular necrosis. AJR Am J Roentgenol 2007;188:515-9.
- 127. Twair A, Ryan M, O'Connell M, Powell T, O'Byrne J, Eustace S. MRI of failed total hip replacement caused by abductor muscle avulsion. AJR Am J Roentgenol 2003;181:1547-50.
- 128. Pfirrmann CW, Notzli HP, Dora C, Hodler J, Zanetti M. Abductor tendons and muscles assessed at MR imaging after total hip arthroplasty in asymptomatic and symptomatic patients. Radiology 2005;235:969-76.
- 129. De Smet AA, Best TM. MR imaging of the distribution and location of acute hamstring injuries in athletes. AJR Am J Roentgenol 2000;174:393-9.
- 130. Slavotinek JP, Verrall GM, Fon GT. Hamstring injury in athletes: using MR imaging measurements to compare extent of muscle injury with amount of time lost from competition. AJR Am J Roentgenol 2002;179:1621-8.
- 131. Askling CM, Tengvar M, Saartok T, Thorstensson A. Acute first-time hamstring strains during slow-speed stretching: clinical, magnetic resonance imaging, and recovery characteristics. Am J Sports Med 2007;35:1716-24.
- 132. O'Brien SD, Bui-Mansfield LT. MRI of quadratus femoris muscle tear: another cause of hip pain. AJR Am J Roentgenol 2007;189:1185-9.
- 133. Bui KL, Ilaslan H, Recht M, Sundaram M. Iliopsoas injury: an MRI study of patterns and prevalence correlated with clinical findings. Skeletal Radiol 2008;37:245-9.
- 134. Kozlov DB, Sonin AH. Iliopsoas bursitis: diagnosis by MRI. J Comput Assist Tomogr 1998;22:625-8.
- 135. Nguyen JT, Peterson JS, Biswal S, Beaulieu CF, Fredericson M. Stress-related injuries around the lesser trochanter in long-distance runners. AJR Am J Roentgenol 2008;190:1616-20.
- 136. Ouellette H, Thomas BJ, Nelson E, Torriani M. MR imaging of rectus femoris origin injuries. Skeletal Radiol 2006;35:665-72.
- 137. Torriani M, Souto SC, Thomas BJ, Ouellette H, Bredella MA. Ischiofemoral impingement syndrome: an entity

with hip pain and abnormalities of the quadratus femoris muscle. AJR Am J Roentgenol 2009;193:186-90.

- 138. Zoga AC, Kavanagh EC, Omar IM, et al. Athletic pubalgia and the "sports hernia": MR imaging findings. Radiology 2008;247:797-807.
- 139. Cunningham PM, Brennan D, O'Connell M, MacMahon P, O'Neill P, Eustace S. Patterns of bone and softtissue injury at the symphysis pubis in soccer players: observations at MRI. AJR Am J Roentgenol 2007;188:W291-6.
- 140. Slavotinek JP, Verrall GM, Fon GT, Sage MR. Groin pain in footballers: the association between preseason clinical and pubic bone magnetic resonance imaging findings and athlete outcome. Am J Sports Med 2005;33:894-9.
- 141. Nelson EN, Kassarjian A, Palmer WE. MR imaging of sports-related groin pain. Magn Reson Imaging Clin N Am 2005;13:727-42.
- 142. Brennan D, O'Connell MJ, Ryan M, et al. Secondary cleft sign as a marker of injury in athletes with groin pain: MR image appearance and interpretation. Radiology 2005;235:162-7.
- 143. Anderson SE, Johnston JO, O'Donnell R, Steinbach LS. MR Imaging of sports-related pseudotumor in children: mid femoral diaphyseal periostitis at insertion site of adductor musculature. AJR Am J Roentgenol 2001;176:1227-31.
- 144. Anderson MW, Kaplan PA, Dussault RG. Adductor insertion avulsion syndrome (thigh splints): spectrum of MR imaging features. AJR Am J Roentgenol 2001;177:673-5.
- 145. Robinson P, Barron DA, Parsons W, Grainger AJ, Schilders EM, O'Connor PJ. Adductor-related groin pain in athletes: correlation of MR imaging with clinical findings. Skeletal Radiol 2004;33:451-7.
- 146. Omar IM, Zoga AC, Kavanagh EC, et al. Athletic pubalgia and "sports hernia": optimal MR imaging technique and findings. Radiographics 2008;28:1415-38.
- 147. Albers SL, Spritzer CE, Garrett WE, Jr., Meyers WC. MR findings in athletes with pubalgia. Skeletal Radiol 2001;30:270-7.
- 148. Sher I, Umans H, Downie SA, Tobin K, Arora R, Olson TR. Proximal iliotibial band syndrome: what is it and where is it? Skeletal Radiol 2011;40:1553-6.
- 149. Beaule PE, Zaragoza E, Copelan N. Magnetic resonance imaging with gadolinium arthrography to assess acetabular cartilage delamination. A report of four cases. J Bone Joint Surg Am 2004;86-A:2294-8.
- 150. Nishii T, Nakanishi K, Sugano N, Masuhara K, Ohzono K, Ochi T. Articular cartilage evaluation in osteoarthritis of the hip with MR imaging under continuous leg traction. Magn Reson Imaging 1998;16:871-5.
- 151. Weaver CJ, Major NM, Garrett WE, Urbaniak JE. Femoral head osteochondral lesions in painful hips of athletes: MR imaging findings. AJR Am J Roentgenol 2002;178:973-7.
- 152. Zaragoza E, Lattanzio PJ, Beaule PE. Magnetic resonance imaging with gadolinium arthrography to assess acetabular cartilage delamination. Hip Int 2009;19:18-23.
- 153. Byrd JW, Jones KS. Traumatic rupture of the ligamentum teres as a source of hip pain. Arthroscopy 2004;20:385-91.
- 154. Chang CY, Gill CM, Huang AJ, et al. Use of MR arthrography in detecting tears of the ligamentum teres with arthroscopic correlation. Skeletal Radiol 2015;44:361-7.
- 155. Robinson P, White LM, Agur A, Wunder J, Bell RS. Obturator externus bursa: anatomic origin and MR imaging features of pathologic involvement. Radiology 2003;228:230-4.
- 156. Varma DG, Richli WR, Charnsangavej C, Samuels BI, Kim EE, Wallace S. MR appearance of the distended iliopsoas bursa. AJR Am J Roentgenol 1991;156:1025-8.
- 157. Gierada DS, Erickson SJ. MR imaging of the sacral plexus: abnormal findings. AJR Am J Roentgenol 1993;160:1067-71.
- 158. Tibor LM, Sekiya JK. Differential diagnosis of pain around the hip joint. Arthroscopy 2008;24:1407-21.
- 159. Huang AB, Schweitzer ME, Hume E, Batte WG. Osteomyelitis of the pelvis/hips in paralyzed patients: accuracy and clinical utility of MRI. J Comput Assist Tomogr 1998;22:437-43.
- 160. Lee SK, Suh KJ, Kim YW, et al. Septic arthritis versus transient synovitis at MR imaging: preliminary assessment with signal intensity alterations in bone marrow. Radiology 1999;211:459-65.
- 161. Connolly SA, Connolly LP, Drubach LA, Zurakowski D, Jaramillo D. MRI for detection of abscess in acute osteomyelitis of the pelvis in children. AJR Am J Roentgenol 2007;189:867-72.
- 162. Karmazyn B, Kleiman MB, Buckwalter K, Loder RT, Siddiqui A, Applegate KE. Acute pyomyositis of the pelvis: the spectrum of clinical presentations and MR findings. Pediatr Radiol 2006;36:338-43.

- 163. Karmazyn B, Loder RT, Kleiman MB, et al. The role of pelvic magnetic resonance in evaluating nonhip sources of infection in children with acute nontraumatic hip pain. J Pediatr Orthop 2007;27:158-64.
- 164. Kwack KS, Cho JH, Lee JH, Oh KK, Kim SY. Septic arthritis versus transient synovitis of the hip: gadoliniumenhanced MRI finding of decreased perfusion at the femoral epiphysis. AJR Am J Roentgenol 2007;189:437-45.
- 165. Bollow M, Braun J, Hamm B, et al. Early sacroiliitis in patients with spondyloarthropathy: evaluation with dynamic gadolinium-enhanced MR imaging. Radiology 1995;194:529-36.
- 166. Boutry N, Paul C, Leroy X, Fredoux D, Migaud H, Cotten A. Rapidly destructive osteoarthritis of the hip: MR imaging findings. AJR Am J Roentgenol 2002;179:657-63.
- 167. Cheng XG, You YH, Liu W, Zhao T, Qu H. MRI features of pigmented villonodular synovitis (PVNS). Clin Rheumatol 2004;23:31-4.
- 168. Otake S, Tsuruta Y, Yamana D, Mizutani H, Ohba S. Amyloid arthropathy of the hip joint: MR demonstration of presumed amyloid lesions in 152 patients with long-term hemodialysis. Eur Radiol 1998;8:1352-6.
- 169. Soini I, Kotaniemi A, Kautiainen H, Kauppi M. US assessment of hip joint synovitis in rheumatic diseases. A comparison with MR imaging. Acta Radiol 2003;44:72-8.
- 170. Sugano N, Ohzono K, Nishii T, et al. Early MRI findings of rapidly destructive coxopathy. Magn Reson Imaging 2001;19:47-50.
- 171. Watanabe W, Itoi E, Yamada S. Early MRI findings of rapidly destructive coxarthrosis. Skeletal Radiol 2002;31:35-8.
- 172. Zhai G, Cicuttini F, Srikanth V, Cooley H, Ding C, Jones G. Factors associated with hip cartilage volume measured by magnetic resonance imaging: the Tasmanian Older Adult Cohort Study. Arthritis Rheum 2005;52:1069-76.
- 173. Mamisch TC, Zilkens C, Siebenrock KA, Bittersohl B, Kim YJ, Werlen S. MRI of hip osteoarthritis and implications for surgery. Radiol Clin North Am 2009;47:713-22.
- 174. Taljanovic MS, Graham AR, Benjamin JB, et al. Bone marrow edema pattern in advanced hip osteoarthritis: quantitative assessment with magnetic resonance imaging and correlation with clinical examination, radiographic findings, and histopathology. Skeletal Radiol 2008;37:423-31.
- 175. Moss SG, Schweitzer ME, Jacobson JA, et al. Hip joint fluid: detection and distribution at MR imaging and US with cadaveric correlation. Radiology 1998;208:43-8.
- 176. Neckers AC, Polster JM, Winalski CS, Krebs VE, Sundaram M. Comparison of MR arthrography with arthroscopy of the hip for the assessment of intra-articular loose bodies. Skeletal Radiol 2007;36:963-7.
- 177. Murphey MD, Wetzel LH, Bramble JM, Levine E, Simpson KM, Lindsley HB. Sacroiliitis: MR imaging findings. Radiology 1991;180:239-44.
- 178. Disler DG, Miklic D. Imaging findings in tumors of the sacrum. AJR Am J Roentgenol 1999;173:1699-706.
- 179. Gaeta M, Minutoli F, Pandolfo I, Vinci S, D'Andrea L, Blandino A. Magnetic resonance imaging findings of osteoid osteoma of the proximal femur. Eur Radiol 2004;14:1582-9.
- 180. Llauger J, Palmer J, Amores S, Bague S, Camins A. Primary tumors of the sacrum: diagnostic imaging. AJR Am J Roentgenol 2000;174:417-24.
- 181. Ozaki T, Putzke M, Burger H, Gosheger G, Winkelmann W, Lindner N. Infiltration of sarcomas into the hip joint: comparison of CT, MRI and histologic findings in 67 cases. Acta Orthop Scand 2002;73:220-6.
- 182. American College of Radiology. ACR–SSR practice parameter for the performance and interpretation of magnetic resonance imaging (MRI) of bone and soft tissue tumors. Available at: <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-SoftTissue-Tumors.pdf</u>. Accessed January 14, 2020.
- 183. Hirata T, Konishiike T, Kawai A, Sato T, Inoue H. Dynamic magnetic resonance imaging of femoral head perfusion in femoral neck fracture. Clin Orthop Relat Res 2001:294-301.
- 184. Kamano M, Narita S, Honda Y, Fukushima K, Yamano Y. Contrast enhanced magnetic resonance imaging for femoral neck fracture. Clin Orthop Relat Res 1998:179-86.
- 185. Woodley SJ, Nicholson HD, Livingstone V, et al. Lateral hip pain: findings from magnetic resonance imaging and clinical examination. J Orthop Sports Phys Ther 2008;38:313-28.
- 186. Bencardino JT, Kassarjian A, Palmer WE. Magnetic resonance imaging of the hip: sports-related injuries. Top Magn Reson Imaging 2003;14:145-60.
- 187. Mellado JM, Perez del Palomar L, Diaz L, Ramos A, Sauri A. Long-standing Morel-Lavallee lesions of the trochanteric region and proximal thigh: MRI features in five patients. AJR Am J Roentgenol 2004;182:1289-94.

- 188. Duffy CM, Taylor FN, Coleman L, Graham HK, Nattrass GR. Magnetic resonance imaging evaluation of surgical management in developmental dysplasia of the hip in childhood. J Pediatr Orthop 2002;22:92-100.
- 189. Jaramillo D, Villegas-Medina O, Laor T, Shapiro F, Millis MB. Gadolinium-enhanced MR imaging of pediatric patients after reduction of dysplastic hips: assessment of femoral head position, factors impeding reduction, and femoral head ischemia. AJR Am J Roentgenol 1998;170:1633-7.
- 190. Kawaguchi AT, Otsuka NY, Delgado ED, Genant HK, Lang P. Magnetic resonance arthrography in children with developmental hip dysplasia. Clin Orthop Relat Res 2000:235-46.
- 191. Westhoff B, Wild A, Seller K, Krauspe R. Magnetic resonance imaging after reduction for congenital dislocation of the hip. Arch Orthop Trauma Surg 2003;123:289-92.
- 192. Ranawat V, Rosendahl K, Jones D. MRI after operative reduction with femoral osteotomy in developmental dysplasia of the hip. Pediatr Radiol 2009;39:161-3.
- 193. Yuksel HY, Yilmaz S, Aksahin E, et al. The evaluation of hip muscles in patients treated with one-stage combined procedure for unilateral developmental dysplasia of the hip: part I: MRI evaluation. J Pediatr Orthop 2009;29:872-8.
- 194. Potter HG, Nestor BJ, Sofka CM, Ho ST, Peters LE, Salvati EA. Magnetic resonance imaging after total hip arthroplasty: evaluation of periprosthetic soft tissue. J Bone Joint Surg Am 2004;86-A:1947-54.
- 195. White LM, Kim JK, Mehta M, et al. Complications of total hip arthroplasty: MR imaging-initial experience. Radiology 2000;215:254-62.
- 196. Johnston C, Kerr J, Ford S, O'Byrne J, Eustace S. MRI as a problem-solving tool in unexplained failed total hip replacement following conventional assessment. Skeletal Radiol 2007;36:955-61.
- 197. Toms AP, Marshall TJ, Cahir J, et al. MRI of early symptomatic metal-on-metal total hip arthroplasty: a retrospective review of radiological findings in 20 hips. Clin Radiol 2008;63:49-58.
- 198. Cooper HJ, Ranawat AS, Potter HG, Foo LF, Jawetz ST, Ranawat CS. Magnetic resonance imaging in the diagnosis and management of hip pain after total hip arthroplasty. J Arthroplasty 2009;24:661-7.
- 199. Westrich GH, Salvati EA, Sharrock N, Potter HG, Sanchez PM, Sculco TP. The effect of intraoperative heparin administered during total hip arthroplasty on the incidence of proximal deep vein thrombosis assessed by magnetic resonance venography. J Arthroplasty 2005;20:42-50.
- 200. Stark DD, McCarthy SM, Filly RA, Parer JT, Hricak H, Callen PW. Pelvimetry by magnetic resonance imaging. AJR Am J Roentgenol 1985;144:947-50.
- 201. Tukeva TA, Aronen HJ, Karjalainen PT, Makela PJ. Low-field MRI pelvimetry. Eur Radiol 1997;7:230-4.
- 202. American College of Radiology. ACR practice parameter for performing and interpreting magnetic resonance imaging (MRI). Available at: <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Perf-Interpret.pdf</u>. Accessed January 14, 2020.
- 203. American College of Radiology. ACR guidance document on MR safe practices: 2020. Available at: <u>https://www.acr.org/-/media/ACR/Files/Radiology-Safety/MR-Safety/Manual-on-MR-Safety.pdf</u>. Accessed July 9, 2020.
- 204. Shellock FG. *Reference Manual for Magnetic Resonance Safety, Implants, and Devices.* 2010 ed. Los Angeles: Biomedical Research Publishing Company; 2010.
- 205. Shellock FG, Crues JV. MR procedures: biologic effects, safety, and patient care. Radiology 2004;232:635-52.
- 206. American College of Radiology. ACR–SPR practice parameter for the use of intravascular contrast media. Available at: <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/IVCM.pdf</u>. Accessed January 14, 2020.
- 207. American College of Radiology. ACR–SIR practice parameter for sedation/analgesia. Available at: <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Sed-Analgesia.pdf</u>. Accessed January 14, 2020.
- 208. Ragnarsson JI, Ekelund L, Karrholm J, Hietala SO. Low field magnetic resonance imaging of femoral neck fractures. Acta Radiol 1989;30:247-52.
- 209. Mosher TJ. Musculoskeletal imaging at 3T: current techniques and future applications. Magn Reson Imaging Clin N Am 2006;14:63-76.
- 210. Ramnath RR. 3T MR imaging of the musculoskeletal system (Part II): clinical applications. Magn Reson Imaging Clin N Am 2006;14:41-62.
- 211. Watanabe A, Boesch C, Siebenrock K, Obata T, Anderson SE. T2 mapping of hip articular cartilage in healthy volunteers at 3T: a study of topographic variation. J Magn Reson Imaging 2007;26:165-71.
- 212. Carballido-Gamio J, Link TM, Li X, et al. Feasibility and reproducibility of relaxometry, morphometric, and

geometrical measurements of the hip joint with magnetic resonance imaging at 3T. J Magn Reson Imaging 2008;28:227-35.

- 213. Bogost GA, Lizerbram EK, Crues JV, 3rd. MR imaging in evaluation of suspected hip fracture: frequency of unsuspected bone and soft-tissue injury. Radiology 1995;197:263-7.
- 214. Hayes CW, Balkissoon AA. Magnetic resonance imaging of the musculoskeletal system. II. The hip. Clin Orthop Relat Res 1996:297-309.
- 215. Hayes CE, Dietz MJ, King BF, Ehman RL. Pelvic imaging with phased-array coils: quantitative assessment of signal-to-noise ratio improvement. J Magn Reson Imaging 1992;2:321-6.
- 216. Niitsu M, Mishima H, Miyakawa S, Itai Y. High resolution MR imaging of the bilateral hips with dual phasedarray coil. J Magn Reson Imaging 1996;6:950-3.
- 217. Do-Dai DD, Youngberg RA. MRI of the hip with a shoulder surface coil in off-coronal plane. J Comput Assist Tomogr 1995;19:336-8.
- 218. Cook SM, Pellicci PM, Potter HG. Use of magnetic resonance imaging in the diagnosis of an occult fracture of the femoral component after total hip arthroplasty. A case report. J Bone Joint Surg Am 2004;86-A:149-53.
- 219. Kwok WE, Lo KK, Seo G, Totterman SM. A volume adjustable four-coil phased array for high resolution MR imaging of the hip. MAGMA 1999;9:59-64.
- 220. Rubin SJ, Totterman SM, Meyers SP, Hartley DF. Magnetic resonance imaging of the hip with a pelvic phased-array surface coil: a technical note. Skeletal Radiol 1998;27:77-82.
- 221. Khoury NJ, Birjawi GA, Chaaya M, Hourani MH. Use of limited MR protocol (coronal STIR) in the evaluation of patients with hip pain. Skeletal Radiol 2003;32:567-74.
- 222. Ha AS, Wells L, Jaramillo D. Importance of sagittal MR imaging in nontraumatic femoral head osteonecrosis in children. Pediatr Radiol 2008;38:1195-200.
- 223. Knuesel PR, Pfirrmann CW, Noetzli HP, et al. MR arthrography of the hip: diagnostic performance of a dedicated water-excitation 3D double-echo steady-state sequence to detect cartilage lesions. AJR Am J Roentgenol 2004;183:1729-35.
- 224. Blake LC, Robertson WD, Hayes CE. Sacral plexus: optimal imaging planes for MR assessment. Radiology 1996;199:767-72.
- 225. Toda K, Yoneda S, Urabe A, Hosoi N. Oblique axial and oblique coronal MR imaging of the proximal femur. AJR Am J Roentgenol 1997;168:1622-3.
- 226. Ziegert AJ, Blankenbaker DG, De Smet AA, Keene JS, Shinki K, Fine JP. Comparison of standard hip MR arthrographic imaging planes and sequences for detection of arthroscopically proven labral tear. AJR Am J Roentgenol 2009;192:1397-400.
- 227. Yoon LS, Palmer WE, Kassarjian A. Evaluation of radial-sequence imaging in detecting acetabular labral tears at hip MR arthrography. Skeletal Radiol 2007;36:1029-33.
- 228. Horii M, Kubo T, Hirasawa Y. Radial MRI of the hip with moderate osteoarthritis. J Bone Joint Surg Br 2000;82:364-8.
- 229. Dudda M, Albers C, Mamisch TC, Werlen S, Beck M. Do normal radiographs exclude asphericity of the femoral head-neck junction? Clin Orthop Relat Res 2009;467:651-9.
- 230. Plotz GM, Brossmann J, von Knoch M, Muhle C, Heller M, Hassenpflug J. Magnetic resonance arthrography of the acetabular labrum: value of radial reconstructions. Arch Orthop Trauma Surg 2001;121:450-7.
- 231. Czerny C, Hofmann S, Urban M, et al. MR arthrography of the adult acetabular capsular-labral complex: correlation with surgery and anatomy. AJR Am J Roentgenol 1999;173:345-9.
- 232. Pienkowski D, Resig J, Talwalkar V, Tylkowski C. Novel three-dimensional MRI technique for study of cartilaginous hip surfaces in Legg-Calve-Perthes disease. J Orthop Res 2009;27:981-8.
- 233. Takao M, Sugano N, Nishii T, et al. Application of three-dimensional magnetic resonance image registration for monitoring hip joint diseases. Magn Reson Imaging 2005;23:665-70.
- 234. Li W, Abram F, Beaudoin G, Berthiaume MJ, Pelletier JP, Martel-Pelletier J. Human hip joint cartilage: MRI quantitative thickness and volume measurements discriminating acetabulum and femoral head. IEEE Trans Biomed Eng 2008;55:2731-40.
- 235. Yan K, Xi Y, Sasiponganan C, Zerr J, Wells JE, Chhabra A. Does 3DMR provide equivalent information as 3DCT for the pre-operative evaluation of adult Hip pain conditions of femoroacetabular impingement and Hip dysplasia? Br J Radiol 2018;91:20180474.
- 236. Kheterpal AB, Bunnell KM, Husseini JS, et al. Value of response to anesthetic injection during hip MR

arthrography to differentiate between intra- and extra-articular pathology. Skeletal Radiol 2020;49:555-61.

- 237. Shuman WP, Castagno AA, Baron RL, Richardson ML. MR imaging of avascular necrosis of the femoral head: value of small-field-of-view sagittal surface-coil images. AJR Am J Roentgenol 1988;150:1073-8.
- 238. Ryan M, Cunningham P, Cantwell C, Brennan D, Eustace S. A comparison of fast MRI of hips with and without parallel imaging using SENSE. Br J Radiol 2005;78:299-302.
- 239. Ramnath RR. 3T MR imaging of the musculoskeletal system (Part I): considerations, coils, and challenges. Magn Reson Imaging Clin N Am 2006;14:27-40.
- 240. Kneeland JB, Shimakawa A, Wehrli FW. Effect of intersection spacing on MR image contrast and study time. Radiology 1986;158:819-22.
- 241. Rubin DA, Kneeland JB. MR imaging of the musculoskeletal system: technical considerations for enhancing image quality and diagnostic yield. AJR Am J Roentgenol 1994;163:1155-63.
- 242. Sasiponganan C, Yan K, Pezeshk P, Xi Y, Chhabra A. Advanced MR imaging of bone marrow: quantification of signal alterations on T1-weighted Dixon and T2-weighted Dixon sequences in red marrow, yellow marrow, and pathologic marrow lesions. Skeletal Radiol 2020;49:541-48.
- 243. Aydingoz U, Ozturk MH. MR imaging of the acetabular labrum: a comparative study of both hips in 180 asymptomatic volunteers. Eur Radiol 2001;11:567-74.
- 244. Bush CH. The magnetic resonance imaging of musculoskeletal hemorrhage. Skeletal Radiol 2000;29:1-9.
- 245. De Smet AA, Fisher DR, Heiner JP, Keene JS. Magnetic resonance imaging of muscle tears. Skeletal Radiol 1990;19:283-6.
- 246. De Smet AA. Magnetic resonance findings in skeletal muscle tears. Skeletal Radiol 1993;22:479-84.
- 247. Nguyen B, Brandser E, Rubin DA. Pains, strains, and fasciculations: lower extremity muscle disorders. Magn Reson Imaging Clin N Am 2000;8:391-408.
- 248. Wolf GL, Joseph PM, Goldstein EJ. Optimal pulsing sequences for MR contrast agents. AJR Am J Roentgenol 1986;147:367-71.
- 249. Rybicki FJ, Chung T, Reid J, Jaramillo D, Mulkern RV, Ma J. Fast three-point dixon MR imaging using lowresolution images for phase correction: a comparison with chemical shift selective fat suppression for pediatric musculoskeletal imaging. AJR Am J Roentgenol 2001;177:1019-23.
- 250. Weinberger E, Shaw DW, White KS, et al. Nontraumatic pediatric musculoskeletal MR imaging: comparison of conventional and fast-spin-echo short inversion time inversion-recovery technique. Radiology 1995;194:721-6.
- 251. Bredella MA, Losasso C, Moelleken SC, Huegli RW, Genant HK, Tirman PF. Three-point Dixon chemical-shift imaging for evaluating articular cartilage defects in the knee joint on a low-field-strength open magnet. AJR Am J Roentgenol 2001;177:1371-5.
- 252. Mahnken AH, Staatz G, Ihme N, Gunther RW. MR signal intensity characteristics in Legg-Calve-Perthes disease. Value of fat-suppressed (STIR) images and contrast-enhanced T1-weighted images. Acta Radiol 2002;43:329-35.
- 253. Lamer S, Dorgeret S, Khairouni A, et al. Femoral head vascularisation in Legg-Calvé-Perthes disease: comparison of dynamic gadolinium-enhanced subtraction MRI with bone scintigraphy. Pediatr Radiol 2002;32:580-5.
- 254. Nishii T, Nakanishi K, Sugano N, Naito H, Tamura S, Ochi T. Acetabular labral tears: contrast-enhanced MR imaging under continuous leg traction. Skeletal Radiol 1996;25:349-56.
- 255. Magee T. Comparison of 3.0-T MR vs 3.0-T MR arthrography of the hip for detection of acetabular labral tears and chondral defects in the same patient population. The British Journal of Radiology 2015;88:20140817.
- 256. Llopis E, Cerezal L, Kassarjian A, Higueras V, Fernandez E. Direct MR arthrography of the hip with leg traction: feasibility for assessing articular cartilage. AJR Am J Roentgenol 2008;190:1124-8.
- 257. Kim YJ, Jaramillo D, Millis MB, Gray ML, Burstein D. Assessment of early osteoarthritis in hip dysplasia with delayed gadolinium-enhanced magnetic resonance imaging of cartilage. J Bone Joint Surg Am 2003;85-A:1987-92.
- 258. Boesen M, Jensen KE, Qvistgaard E, et al. Delayed gadolinium-enhanced magnetic resonance imaging (dGEMRIC) of hip joint cartilage: better cartilage delineation after intra-articular than intravenous gadolinium injection. Acta Radiol 2006;47:391-6.
- 259. Nishii T, Tanaka H, Sugano N, Sakai T, Hananouchi T, Yoshikawa H. Evaluation of cartilage matrix disorders by T2 relaxation time in patients with hip dysplasia. Osteoarthritis Cartilage 2008;16:227-33.

- Receive de 2022 (Restatution 2140), Cardenas-Blanco A, Cameron IG, Beaule PE. Can T1-rho MRI detect acetabular cartilage degeneration in femoroacetabular impingement?: a pilot study. J Bone Joint Surg Br 2012;94:1187-92.
 - 261. Rakhra KS, Cardenas-Blanco A, Melkus G, Schweitzer ME, Cameron IG, Beaule PE. Is the t1rho MRI profile of hyaline cartilage in the normal hip uniform? Clin Orthop Relat Res 2015;473:1325-32.
 - 262. Haacke EM, Lenz GW. Improving MR image quality in the presence of motion by using rephasing gradients. AJR Am J Roentgenol 1987;148:1251-8.
 - 263. Peh WC, Chan JH. Artifacts in musculoskeletal magnetic resonance imaging: identification and correction. Skeletal Radiol 2001;30:179-91.
 - 264. Tormanen J, Tervonen O, Koivula A, Junila J, Suramo I. Image technique optimization in MR imaging of a titanium alloy joint prosthesis. J Magn Reson Imaging 1996;6:805-11.
 - 265. Hilfiker P, Zanetti M, Debatin JF, McKinnon G, Hodler J. Fast spin-echo inversion-recovery imaging versus fast T2-weighted spin-echo imaging in bone marrow abnormalities. Invest Radiol 1995;30:110-4.
 - 266. Eustace S, Jara H, Goldberg R, et al. A comparison of conventional spin-echo and turbo spin-echo imaging of soft tissues adjacent to orthopedic hardware. AJR Am J Roentgenol 1998;170:455-8.
 - 267. Fritz J, Lurie B, Miller TT, Potter HG. MR imaging of hip arthroplasty implants. Radiographics 2014;34:E106-32.
 - 268. Lazik A, Landgraeber S, Schulte P, Kraff O, Lauenstein TC, Theysohn JM. Usefulness of metal artifact reduction with WARP technique at 1.5 and 3T MRI in imaging metal-on-metal hip resurfacings. Skeletal Radiol 2015;44:941-51.
 - 269. Schulte-Altedorneburg G, Gebhard M, Wohlgemuth WA, et al. MR arthrography: pharmacology, efficacy and safety in clinical trials. Skeletal Radiol 2003;32:1-12.
 - 270. American College of Radiology. ACR practice parameter for communication of diagnostic imaging findings. Available at: <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CommunicationDiag.pdf</u>. Accessed January 14, 2020.
 - 271. Shellock FG. *Guide to MR Procedures and Metallic Objects*. 7th ed. Philadelphia,PA: Lippincott Williams and Wilkins; 2001.
 - 272. Sawyer-Glover AM, Shellock FG. Pre-MRI procedure screening: recommendations and safety considerations for biomedical implants and devices. J Magn Reson Imaging 2000;12:92-106.
 - 273. American College of Radiology. ACR–AAPM technical standard for diagnostic medical physics performance monitoring of magnetic resonance imaging (MRI) equipment. Available at: <u>https://www.acr.org/-/media/ACR/Files/Practice-Parameters/MR-Equip.pdf</u>. Accessed July 9, 2020.

*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

Development Chronology for this Practice Parameter

2006 (Resolution 6, 35)

2011 (Resolution 21)

Amended 2014 (Resolution 39)

Revised 2016 (Resolution 7)

Revised 2021 (Resolution 40)

Amended 2023 (Resolution 2c)