

ACR–SPR PRACTICE PARAMETER FOR THE PERFORMANCE OF FLUOROSCOPIC AND SONOGRAPHIC VOIDING CYSTOURETHROGRAPHY IN CHILDREN

The American College of Radiology, with more than 40,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care¹. For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

¹ *Iowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing*, 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the "ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures (Revised 2008)" sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, *Stanley v. McCarver*, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

I. INTRODUCTION

This practice parameter was revised collaboratively by the American College of Radiology (ACR) and the Society for Pediatric Radiology (SPR).

Voiding cystourethrography (VCUG) is a radiographic and fluoroscopic study of the lower urinary tract. It requires aseptic bladder catheterization, instillation of iodinated contrast media, fluoroscopic observation, and image documentation of the findings. The purpose of the examination is to assess the bladder, the urethra, other opacified structures, the presence or absence of vesicoureteral reflux (VUR), and micturition. Contrast-enhanced voiding urosonography (ceVUS) is an alternative method of assessment, in which ultrasonography of the kidneys, bladder, and urethra is performed after the intravesical administration of a sonographic contrast agent.

II. INDICATIONS AND CONTRAINDICATIONS [1-10]

Clinical indications for VCUG include, but are not limited to, the following:

- Fetal urinary tract dilation in the increased risk category
- Postnatal hydronephrosis and/or hydroureter
- Abnormal ultrasound (US) (any degree of hydronephrosis, uroepithelial thickening, scarring) after first urinary tract infection (UTI), especially if febrile or non-*Escherichia coli*
- Recurrent UTI
- Congenital anomalies of the urinary tract
- Dysfunctional voiding, such as neurogenic dysfunction of the bladder
- Urinary incontinence
- Bladder outlet obstruction
- Postoperative evaluation of the urinary tract
- Dysuria/difficulty voiding
- Hematuria
- Pelvic Trauma

In some circumstances, a retrograde urethrogram may be preferred over a VCUG, particularly in children with a history of prior urethral operation, trauma, or dysuria. There are no absolute contraindications for VCUG. Potential benefits must outweigh the minor risks of the procedure. However, one should proceed with caution if the child has had a significant reaction to iodinated contrast media, known or suspected latex allergy [11], acute UTI, recent urethral or bladder surgery, potential urethral trauma, or high spinal injury (risk of autonomic dysreflexia). The risk of postprocedure UTI is very low. Routine antibiotic prophylaxis is not necessary, but increased vigilance is warranted in children with significant anatomic abnormalities, particularly high-grade VUR [12]. Use of antibiotic therapy pre- and postprocedure should be made on individual clinical grounds.

ceVUS is a radiation-free alternative to VCUG [13]. See later paragraph.

Nuclear voiding cystography may also be used as an alternative study for the evaluation of reflux in children, especially when detailed anatomic visualization of the urethra, bladder, and kidneys is not required. In the era of modern digital, grid-controlled fluoroscopic equipment with a flat detector, filtration, and low pulse rate, the radiation dose from fluoroscopic VCUG and nuclear VCUG is similar [14,15].

The voiding study chosen will vary depending on locally available equipment, expertise, and personnel.

For the pregnant or potentially pregnant patient, see the [ACR–SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation](#) [16].

III. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

See the [ACR–AAPM–SIIM–SPR Practice Parameter for Digital Radiography](#) [17].

IV. SPECIFICATIONS OF EXAMINATION

A. Patient Selection and Preparation

The study should be performed only for an appropriate clinical indication. Consultation with referring physicians helps to clarify which children may benefit from VCUG.

B. Technique

Preparation and Sedation

VCUG can typically be performed without sedation when parents and children receive adequate preparation and support. When available, Certified Child-Life Specialists may provide education, distraction, and relaxation techniques that are useful in facilitating catheterization as well as patient cooperation during the examination [18,19]. The use of warmed contrast material may decrease patient distress [18,20]. When clinically indicated in select patients, sedation may alleviate distress and can be performed safely, without negatively affecting the examination [18,21,22]. If sedation is used, the child must undergo a presedation evaluation and must be monitored both during and after the examination, as outlined in the current practice parameter (See the [ACR–SIR Practice Parameter for Minimal and/or Moderate Sedation/Analgesia](#) [23]).

Review and Preliminary Imaging

Before the start of the examination, it is helpful to review the child's history and prior imaging findings, particularly any recent studies in which residual contrast may be present. Formal scout radiographs make a significant contribution to the dose of the study while their yield is relatively low [24]. Instead, a precontrast fluoroscopy grab image may be acquired at the start of the study. If the fluoroscopy unit is set with limited resolution to detect anatomy and calcifications from image-hold or image grab technique or in those specific situations requiring superior spatial resolution, a digitally acquired fluorographic spot image or radiograph may be obtained [24-27].

Catheterization

Aseptic bladder catheterization of children should be performed by experienced personnel. Latex precautions should be observed, especially in children with known latex allergy, multiple surgeries, or myelomeningocele.

In males who are old enough to cooperate, to diminish sensation or pain, a topical anesthetic may be instilled retrograde into the urethra with aseptic technique; anesthetic gel may be applied externally in both males and females [28,29].

The catheter size should be appropriate for the child's age or urethral caliber. In premature or extremely small infants, a 5-French catheter is preferred. Above this age, an 8-French catheter is preferred, unless a smaller catheter is appropriate (such as in the case of urethral stricture) or if there is inability to catheterize with the larger catheter. A catheter larger than 8-French may be used in adolescents. Use of a catheter with a tapered, curved end (coude catheter) may help to negotiate catheter passage through strictures/unusual urethral contours. To avoid intravesical looping and knotting of the catheter, which may require invasive retrieval, excessive catheter length should not be inserted into the bladder. Importantly, in uncircumcised males, if the foreskin is retracted at catheterization, it should be repositioned over the glans immediately following catheterization, to avoid secondary paraphimosis.

In girls, the catheter may be secured in place with tape to the perineum or, in older girls, to the thigh. In boys, once the catheter is inserted, a strip of tape may be placed on the catheter extending longitudinally along the dorsum of the penis to the symphysis. Circumferential placement of tape around the penis is discouraged. Inflation of a balloon in the bladder to secure the catheter is discouraged because it may obscure pathology and/or impair free voiding.

After catheter placement, the bladder should be drained before instillation of contrast media. A sterile urine specimen may be retained for culture if clinically indicated. The size of the catheter should be recorded, unless the catheter had been previously placed in another department.

Contrast Media

Iodinated contrast media (typically 12%–18% weight/volume solution) should be administered via the bladder catheter by gravity drip. The height of the bottle controls the infusion pressure, while the diameter of the tubing somewhat limits this pressure, making the exact bottle height relatively unimportant. However, 1 m above the table height is typically sufficient [30]. The amount of contrast agent to be administered depends on the bladder capacity, and can be estimated using the following formula [35,44]:

$(\text{age in years} + 2) \times 30 = \text{capacity (mL)}$.

This formula is a general guide, not a rule, and may not be applicable in all children such as a premature infant or a child with unusually small or large bladder capacity. The bladder filling should continue until voiding, filling with at most twice the estimated bladder capacity if voiding does not occur at the estimated volume.

If recent bladder surgery has been performed, such as augmentation, gravity infusion should be performed with the bottle of contrast positioned at or less than 1 m above the table height to assure low-pressure filling. Contrast infusion should be stopped when the patient has symptoms of pain, when contrast refluxes retrograde into the ureters beyond the ureteral stents (if present), or if contrast leakage is observed. For this indication, typical volumes of contrast infusion do not exceed 75 mL.

Very few patients have allergic reactions to intravesical contrast [32]. However, in the event that a patient has had prior anaphylaxis to contrast, one could consider allergy prophylaxis or an alternative imaging study (See the [ACR–SPR Practice Parameter for the Use of Intravascular Contrast Media](#) [23] and the [ACR Manual on Contrast Media](#) [33]).

Fluoroscopy and Imaging

Standardization of an optimized imaging sequence should be employed as much as possible. Early-filling last-image capture of the bladder with a small amount of contrast may reveal an intravesical ureterocele or other mass, which might be obscured by larger contrast volume. Early oblique last-image capture views can be obtained if necessary. While further bladder filling occurs, continuous imaging is not necessary, although intermittent fluoroscopic monitoring at regular intervals could be helpful to detect intermittent VUR, documenting each fluoroscopic "look" with an image capture of bladder [35].

Oblique image capture images or fluorographic spot images of the bladder are obtained when the bladder is estimated to be full, before voiding to profile each ureterovesical junction in relation to the bladder neck (as delineated by the catheter). Concluding a study when reaching the predicted bladder capacity may miss children with functional or neurological abnormalities (eg, infrequent voider syndrome that is characterized by a large capacity bladder). In some patients, overdistension of the bladder may result in urinary retention/inability to void [36]. If reflux did not occur before voiding and is seen during voiding, the oblique images of the bladder should be repeated with some degree of bladder distention; if the bladder empties before documenting distal ureteral anatomy, refilling the bladder at least partially may be necessary if the catheter is still within the bladder. Lateral imaging of the urinary bladder may be helpful in certain indications such as for evaluation of urachal pathology.

Cyclical filling of the bladder (filling to capacity followed by voiding and refilling 2-3 times with the catheter in place) may be helpful in infants (1 year of age or younger) who void at low volumes and in patients at any age with a high pretest probability of reflux to increase detection of reflux. The latter group includes patients with recent UTI, a prior history of reflux, paraureteric (Hutch) diverticulum, and evidence of pyelonephritis. Cyclical voiding is also helpful in patients who reflux into a very dilated ureter on the first fill; in these patients, the refluxed contrast may be diluted by unopacified urine, and additional voiding cycles will optimize visualization of the ureter and collecting system [37,38]. Cyclical voiding is also helpful in cases of suspected ectopic ureter inserting below the bladder base [39].

When VUR occurs, the degree of reflux should be documented by imaging the renal fossae in the frontal projection, taking care to include the renal parenchyma in the field of view to assess for intrarenal reflux into the ducts of Bellini. Additionally, bladder volume at the onset of reflux may be recorded. Postvoid residual may be estimated [31].

For optimal imaging of the urethra, the field of view and patient positioning should be prepared before the child begins to void. Visual inspection of the perineum will reveal when the patient begins to void and avoids unnecessary and excessive fluoroscopy. Older males might be able to void more easily if the fluoroscopic table is tilted with a foot stand in place to 30–45° or up to 90° if they are able to stand.

Once voiding is detected, the male urethra may be imaged before removal of the catheter, because pertinent pathology may be demonstrated with the catheter in place [40]. However, it is preferable to also obtain images of the urethra after the catheter has been removed, especially in boys. In neonates and young infants who may void sporadically, if the bladder has moderately emptied before catheter removal, it is best to wait until it refills before removing the catheter because the child may not void again without a full bladder.

The entire urethra should be demonstrated during the voiding phase. To image the urethra, boys should be positioned slightly obliquely from the lateral position during voiding, with slight offset of the hips and legs

such that they are not superimposed. In most young boys, the entire urethra will be visible on a single voiding image. In adolescent boys, separate images of the posterior and anterior urethra may be necessary. In girls, the urethra is generally imaged in the frontal projection, and catheter removal is not necessary for optimal views. Lateral imaging of the female urethra is performed in special circumstances, such as evaluation of suspected urogenital sinus anomalies.

Images of the renal fossae immediately after voiding should be obtained to document the presence and grade of reflux or its absence. The maximal degree of VUR should be accurately described and graded (see Appendix A). When an obstructive process, such as obstruction at the ureteropelvic or ureterovesical junction, coexists with reflux, refluxed contrast will be diluted by the indwelling unopacified urine, with decreased density of the refluxed contrast. In such situations, one should refrain from grading the reflux because the degree of urinary tract dilation is not necessarily secondary to the reflux alone. An attempt at reflux grading in this situation may result in overestimation in the contribution of reflux to urinary tract pathology [41].

If there is concern for coexistent obstruction, the rate of contrast drainage from the pelvicalyceal system and ureter may be estimated by obtaining a delayed image. In patients with high-grade reflux, a screen capture image can be obtained after sitting upright for 5–10 mins following complete or close to complete voiding to document at least partial drainage of the refluxed contrast and exclude significant obstruction. If a toilet-trained patient is unable to void during the fluoroscopic portion of the examination, after adequate bladder distension and after a reasonable amount of time and coaxing, they may be allowed to void in the restroom if it is sufficiently adjacent to the fluoroscopy room. Postvoid images should be obtained immediately after voiding in the restroom, with an estimate of the time interval between completion of voiding and imaging recorded. As in postvoid imaging when voiding is directly observed on the fluoroscopy table, images should be obtained over the renal fossae and bladder to document the presence or absence of reflux and degree of bladder emptying. This limitation of the voiding portion of the examination must be documented.

Alternate strategies to achieve voiding in the reluctant child include the following: The male patient may be given a urinal and put into the upright position to try to induce voiding. The female patient may be asked to sit upright on the fluoroscopy table on a bedpan.

Study interpretation should include analysis of the following:

- Appearance of the spine and pelvic bones. Evaluation may be limited on a precontrast fluoroscopic image. If an abnormality is suspected, a radiographic exposure of that abnormality may be obtained.
- Documentation of opaque calculi, calcifications, or foreign bodies when present. Evaluation may be limited on a precontrast fluoroscopic image. If an abnormality is suspected, a radiographic exposure of that abnormality may be obtained.
- Bladder contour, location, capacity, diverticula, and residual volume.
- Bladder lumen filling defects, such as ureterocele, clot, or other masses.
- Absence or presence and greatest degree of reflux including intrarenal reflux.
- When during the examination reflux, if present, first occurred.
- Site of insertion of ureter(s) when visualized, due to refluxed contrast.
- Appearance of the entire urethra.
- Presence or absence of extravasation or fistula.
- Documentation of drainage of refluxed material in high grade reflux.

V. DOCUMENTATION

The findings of the voiding cystourethrogram should be reported in accordance with the [ACR Practice Parameter for Communication of Diagnostic Imaging Findings](#) [42].

VI. EQUIPMENT SPECIFICATIONS

Equipment performance monitoring should be in accordance with the [ACR–AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Fluoroscopic Equipment](#) [43].

VII. RADIATION SAFETY IN IMAGING

Radiologists, medical physicists, non-physician radiology providers, radiologic technologists, and all supervising physicians have a responsibility for safety in the workplace by keeping radiation exposure to staff, and to society as a whole, "as low as reasonably achievable" (ALARA) and to assure that radiation doses to individual patients are appropriate, taking into account the possible risk from radiation exposure and the diagnostic image quality necessary to achieve the clinical objective. All personnel who work with ionizing radiation must understand the key principles of occupational and public radiation protection (justification, optimization of protection, application of dose constraints and limits) and the principles of proper management of radiation dose to patients (justification, optimization including the use of dose reference levels). https://www-pub.iaea.org/MTCD/Publications/PDF/PUB1775_web.pdf

Nationally developed guidelines, such as the [ACR's Appropriateness Criteria®](#), should be used to help choose the most appropriate imaging procedures to prevent unnecessary radiation exposure.

Facilities should have and adhere to policies and procedures that require ionizing radiation examination protocols (radiography, fluoroscopy, interventional radiology, CT) to vary according to diagnostic requirements and patient body habitus to optimize the relationship between appropriate radiation dose and adequate image quality. Automated dose reduction technologies available on imaging equipment should be used, except when inappropriate for a specific exam. If such technology is not available, appropriate manual techniques should be used.

Additional information regarding patient radiation safety in imaging is available from the following websites – Image Gently® for children (www.imagegently.org) and Image Wisely® for adults (www.imagewisely.org). These advocacy and awareness campaigns provide free educational materials for all stakeholders involved in imaging (patients, technologists, referring providers, medical physicists, and radiologists).

Radiation exposures or other dose indices should be periodically measured by a Qualified Medical Physicist in accordance with the applicable ACR Technical Standards. Monitoring or regular review of dose indices from patient imaging should be performed by comparing the facility's dose information with national benchmarks, such as the ACR Dose Index Registry and relevant publications relying on its data, applicable ACR Practice Parameters, NCRP Report No. 172, Reference Levels and Achievable Doses in Medical and Dental Imaging: Recommendations for the United States or the Conference of Radiation Control Program Director's National Evaluation of X-ray Trends; 2006, 2009, amended 2013, revised 2023 (Res. 2d).

VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *Position Statement on QC & Improvement, Safety, Infection Control, and Patient Education* on the ACR website (<https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement>).

IX. CONTRAST ENHANCED VOIDING UROSONOGRAPHY (ceVUS) INTRODUCTION

The intravesical administration of US contrast agent and the demonstration of refluxing microbubbles into a ureter or renal collecting system is known as ceVUS [44-46]. In 2016, the Food and Drug Administration of the United States approved the use of an US contrast agent (Lumason, Bracco) for use in vesicoureteric reflux detection in children. Studies of ceVUS versus VCUG have demonstrated that ceVUS is more sensitive in detecting vesicoureteric reflux and is comparable in the evaluation of the urethra [45,46]. Unlike VCUG, there is no exposure to radiation.

X. CONTRAST ENHANCED VOIDING UROSONOGRAPHY (ceVUS) INDICATIONS AND CONTRAINDICATIONS

The indications for ceVUS are the same as for VCUG. The study is contraindicated if one or both kidneys are difficult to visualize sonographically, which may occur in a morbidly obese child or a child with severe lumbar scoliosis [13]. There are no known side effects related to intravesical administration of US contrast agents [47].

XI. CONTRAST ENHANCED VOIDING UROSONOGRAPHY (ceVUS) QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL

ceVUS is performed by sonographers or radiologists trained in bladder catheterization and sonography of the bladder, ureters, kidneys and urethra. It is best to have two persons present for the examination; one performing the catheterization and administration of intravesical US contrast and the other performing the US itself. The ceVUS is documented with static images and clips.

XII. CONTRAST ENHANCED VOIDING UROSONOGRAPHY (ceVUS) SPECIFICATIONS OF EXAMINATION

The request for the study, patient selection and preparation, review of prior imaging, and catheterization process are similar to those for VCUGs. The study is ideally performed with an US scanner equipped with a contrast-specific mode, which optimizes contrast visualization during the examination. Scout sonographic images of the kidneys and bladder should be obtained before catheterization. The US contrast agent is prepared as per the instruction of the manufacturer. The microbubble solution is mixed immediately before its use to avoid microbubble degradation/loss of suspension over time. The bladder should be decompressed after catheterization.

A 0.2% suspension of the US contrast agent and normal saline may be administered into the bladder catheter via gravity drip. Alternatively, 0.5–1.0 mL of the contrast agent may be administered via the bladder catheter followed by a normal saline infusion [48]. The height of the infusion and the volume of suspension or normal saline administered are similar to those for VCUG.

Bladder filling is carried out under US monitoring with the patient supine. This is followed by alternate scanning of the right and left kidneys (supine or oblique if necessary) and the bladder, especially obliquely at the ureterovesical junctions. During voiding, additional suprapubic and/or transperineal scanning of the urethra are performed with the catheter in place as well as once the catheter has been removed. Patients may be scanned from the back while sitting on a potty/toilet or, in males, while standing and using a urinal, thus allowing a more physiological position for voiding.

Additional postvoid scanning of the bladder and kidneys is subsequently obtained. The appearance of echogenic microbubbles within a ureter or renal collecting systems indicates vesicoureteric reflux. Intraparenchymal reflux of echogenic microbubbles should be documented. Vesicoureteric reflux is graded in a similar way to VCUG [44]. As with a VCUG, cyclical filling may be necessary in neonates and infants [48].

Assessment and reporting is similar to VCUG with the exception of structures not applicable to US such as the bony pelvis and fluoroscopic radiation. Additionally, the degree of urinary tract dilation, if present, and quality of the renal parenchyma can be described.

ACKNOWLEDGEMENTS

This practice parameter was revised according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (<https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards>) by the Committee on Practice Parameters – Pediatric Radiology of the ACR Commission on Pediatric Radiology in collaboration with the SPR.

Writing Committee

Members represent their societies in the initial and final revision of this practice parameter.

ACR

John Amodio, MD, Chair

SPR

Susan J. Back, MD

Jane Sun Kim, MD

Carol Barnewolt, MD

Terry L. Levin, MD, FACR

Jeanne Chow, MD

Sakura Noda, MD

Steven Kraus, MD

Akshay Saxena, MD

Committee on Practice Parameters – Pediatric Radiology

(ACR Committee responsible for sponsoring the draft through the process)

Terry L. Levin, MD, FACR, Chair

Zofia Lasiecka, MD, PhD

John B. Amodio, MD, FACR

Arleen Li, MD

Bradford W. Betz, MD, FACR

Lauren P. Nicola, MD

Harris L. Cohen, MD, FACR

Sakura Noda, MD

Michael Collard, MD

Erica Poletto, MD

Adam Goldman-Yassen, MD

Summit Shah, MD

Jane Sun Kim, MD

Andrew T. Trout, MD

Jessica Kurian, MD

Esben S. Vogelius, MD

Hollie A. Lai, MD

Jason Wright, MD

Adina Laufer, MD

Richard A. Barth, MD, FACR, Chair, Commission on Pediatric Radiology

David B. Larson, MD, MBA, FACR, Chair, Commission on Quality and Safety

Mary S. Newell, MD, FACR, Chair, Committee on Practice Parameters and Technical Standards

Comments Reconciliation Committee

Sonia Gupta, MD, Chair

Paul A Larson, MD, FACR

Kevin Smith, MD, FACR, Co-Chair

Terry Levin, MD, FACR

Richard A. Barth, MD, FACR

Mary S. Newell, MD, FACR

Jacqueline A. Bello, MD, FACR

Beverley Newman, MB, BCh, BSc, FACR

Daniel W. Brown, MD

Hansel J. Otero, MD

Ellen Chung, MD

Matthew S. Pollack, MD, FACR

Kassa Darge, MD, PhD

Anil G. Rao, DMRD, DNB

Richard Duszak, Jr., MD, FACR

Timothy L. Swan, MD, FACR

REFERENCES

1. Estrada CR, Peters CA, Retik AB, Nguyen HT. Vesicoureteral reflux and urinary tract infection in children with a history of prenatal hydronephrosis--should voiding cystourethrography be performed in cases of postnatally persistent grade II hydronephrosis? *The Journal of urology* 2009;181:801-6; discussion 06-7.
2. Lee JH, Kim MK, Park SE. Is a routine voiding cystourethrogram necessary in children after the first febrile urinary tract infection? *Acta Paediatrica* 2012;101:105-9.
3. Spencer JD, Bates CM, Mahan JD, et al. The accuracy and health risks of a voiding cystourethrogram after a febrile urinary tract infection. *Journal of pediatric urology* 2012;8:72-6.
4. Lee NG, Rushton HG, Peters CA, Groves DS, Pohl HG. Evaluation of prenatal hydronephrosis: novel criteria for predicting vesicoureteral reflux on ultrasonography. *The Journal of urology* 2014;192:914-8.
5. Kovanlikaya A, Kazam J, Dunning A, et al. The role of ultrasonography in predicting vesicoureteral reflux. *Urology* 2014;84:1205-10.
6. Gordon ZN, McLeod DJ, Becknell B, Bates DG, Alpert SA. Uroepithelial Thickening on Sonography Improves Detection of Vesicoureteral Reflux in Children with First Febrile Urinary Tract Infection. *The Journal of urology* 2015;194:1074-9.
7. Ristola MT, Loyttyniemi E, Hurme T. Factors Associated with Abnormal Imaging and Infection Recurrence after a First Febrile Urinary Tract Infection in Children. *Eur J Pediatr Surg* 2017;27:142-49.
8. Ristola MT, Hurme T. Consequences of following the new American Academy of Pediatrics guidelines for imaging children with urinary tract infection. *Scand J Urol* 2015;49:419-23.
9. Roberts KB. Urinary tract infection: clinical practice guideline for the diagnosis and management of the initial UTI in febrile infants and children 2 to 24 months. *Pediatrics* 2011;128:595-610.
10. Nguyen HT, Benson CB, Bromley B, et al. Multidisciplinary consensus on the classification of prenatal and postnatal urinary tract dilation (UTD classification system). *Journal of pediatric urology* 2014;10:982-98.
11. Zerlin JM, McLaughlin K, Kerchner S. Latex allergy in patients with myelomeningocele presenting for imaging studies of the urinary tract. *Pediatric radiology* 1996;26:450-4.
12. Johnson EK, Malhotra NR, Shannon R, et al. Urinary tract infection after voiding cystourethrogram. *Journal of pediatric urology* 2017;13:384.e1-84.e7.
13. Darge K. Voiding urosonography with ultrasound contrast agents for the diagnosis of vesicoureteric reflux in

children. I. Procedure. *Pediatric radiology* 2008;38:40-53.

14. Haid B, Becker T, Koen M, et al. Lower radiation burden in state of the art fluoroscopic cystography compared to direct isotope cystography in children. *Journal of pediatric urology* 2015;11:35 e1-6.
15. Linke SY, Tsiflikas I, Herz K, Szavay P, Gatidis S, Schafer JF. Ultra low-dose VCUG in children using a modern flat detector unit. *European radiology* 2016;26:1678-85.
16. American College of Radiology. ACR–SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Pregnant-Pts.pdf>. Accessed January 13, 2023.
17. American College of Radiology. ACR–AAPM–SIIM–SPR Practice Parameter for Digital Radiography. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Rad-Digital.pdf>. Accessed January 13, 2023.
18. Rao J, Kennedy SE, Cohen S, Rosenberg AR. A systematic review of interventions for reducing pain and distress in children undergoing voiding cystourethrography. *Acta Paediatr* 2012;101:224-9.
19. Sandy NS, Nguyen HT, Ziniel SI, et al. Assessment of parental satisfaction in children undergoing voiding cystourethrography without sedation. *The Journal of urology* 2011;185:658-62.
20. Goodman TR, Kilborn T, Pearce R. Warm or cold contrast medium in the micturating cystourethrogram (MCUG): which is best? *Clinical radiology* 2003;58:551-4.
21. Herd DW, McAnulty KA, Keene NA, Sommerville DE. Conscious sedation reduces distress in children undergoing voiding cystourethrography and does not interfere with the diagnosis of vesicoureteric reflux: a randomized controlled study. *AJR. American journal of roentgenology* 2006;187:1621-6.
22. Stokland E, Andreasson S, Jacobsson B, Jodal U, Ljung B. Sedation with midazolam for voiding cystourethrography in children: a randomised double-blind study. *Pediatric radiology* 2003;33:247-9.
23. American College of Radiology. ACR–SIR Practice Parameter for Sedation/Analgesia. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Sed-Analgesia.pdf>. Accessed January 9, 2018.
24. Darge K, Grattan-Smith JD, Riccabona M. Pediatric uro-radiology: state of the art. *Pediatric radiology* 2011;41:82-91.
25. Hernanz-Schulman M, Goske MJ, Bercha IH, Strauss KJ. Pause and pulse: ten steps that help manage radiation dose during pediatric fluoroscopy. *AJR. American journal of roentgenology* 2011;197:475-81.
26. Creeden SG, Rao AG, Eklund MJ, Hill JG, Thacker PG. Pre-procedural scout radiographs are unnecessary for routine pediatric fluoroscopic examinations. *Pediatric radiology* 2017;47:290-93.
27. Rao AG, Simmons CE, Sr., Thacker PG, Collins H, Ritenour ER, Hill JG. Radiation exposure contribution of the scout abdomen radiograph in common pediatric fluoroscopic procedures. *Pediatric radiology* 2016;46:1241-8.
28. Mularoni PP, Cohen LL, DeGuzman M, Mennuti-Washburn J, Greenwald M, Simon HK. A randomized clinical trial of lidocaine gel for reducing infant distress during urethral catheterization. *Pediatric emergency care* 2009;25:439-43.
29. Vaughan M, Paton EA, Bush A, Pershad J. Does lidocaine gel alleviate the pain of bladder catheterization in young children? A randomized, controlled trial. *Pediatrics* 2005;116:917-20.
30. Kuta AJ, Benator RM. Intussusception: hydrostatic pressure equivalents for barium and meglumine sodium diatrizoate. *Radiology* 1990;175:125-6.
31. Frimberger D, Bauer SB, Cain MP, et al. Establishing a standard protocol for the voiding cystourethrography. *Journal of pediatric urology* 2016;12:362-66.
32. Weese DL, Greenberg HM, Zimmern PE. Contrast media reactions during voiding cystourethrography or retrograde pyelography. *Urology* 1993;41:81-4.
33. American College of Radiology. ACR Manual on Contrast Media. Available at: https://www.acr.org/-/media/ACR/Files/Clinical-Resources/Contrast_Media.pdf. Accessed January 13, 2023.
34. Ward VL. Patient dose reduction during voiding cystourethrography. *Pediatric radiology* 2006;36 Suppl 2:168-72.
35. Fernbach SK, Feinstein KA, Schmidt MB. Pediatric voiding cystourethrography: a pictorial guide. *Radiographics : a review publication of the Radiological Society of North America, Inc* 2000;20:155-68; discussion 68-71.
36. Choong S, Emberton M. Acute urinary retention. *BJU international* 2000;85:186-201.
37. Paltiel HJ, Rupich RC, Kiruluta HG. Enhanced detection of vesicoureteral reflux in infants and children with use of cyclic voiding cystourethrography. *Radiology* 1992;184:753-5.

38. Papadopoulou F, Efremidis SC, Oiconomou A, et al. Cyclic voiding cystourethrography: is vesicoureteral reflux missed with standard voiding cystourethrography? *European radiology* 2002;12:666-70.
39. Wyly JB, Lebowitz RL. Refluxing urethral ectopic ureters: recognition by the cyclic voiding cystourethrogram. *AJR. American journal of roentgenology* 1984;142:1263-7.
40. Chaumoitre K, Merrot T, Petit P, Sayegh-Martin Y, Alessandrini P, Panuel M. Voiding cystourethrography in boys. Does the presence of the catheter during voiding alter the evaluation of the urethra? *The Journal of urology* 2004;171:1280-1.
41. Lebowitz RL, Blickman JG. The coexistence of ureteropelvic junction obstruction and reflux. *AJR. American journal of roentgenology* 1983;140:231-8.
42. American College of Radiology. ACR Practice Parameter for Communication of Diagnostic Imaging Findings. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CommunicationDiag.pdf>. Accessed January 13, 2023.
43. American College of Radiology. ACR-AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Fluoroscopic Equipment. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/Fluoro-Equip.pdf>. Accessed January 13, 2023.
44. Duran C, Beltran VP, Gonzalez A, Gomez C, Riego JD. Contrast-enhanced Voiding Urosonography for Vesicoureteral Reflux Diagnosis in Children. *Radiographics : a review publication of the Radiological Society of North America, Inc* 2017;37:1854-69.
45. Darge K. Voiding urosonography with US contrast agents for the diagnosis of vesicoureteric reflux in children. II. Comparison with radiological examinations. *Pediatric radiology* 2008;38:54-63; quiz 126-7.
46. Papadopoulou F, Anthopoulou A, Siomou E, Efremidis S, Tsamboulas C, Darge K. Harmonic voiding urosonography with a second-generation contrast agent for the diagnosis of vesicoureteral reflux. *Pediatric radiology* 2009;39:239-44.
47. Darge K, Papadopoulou F, Ntoulia A, et al. Safety of contrast-enhanced ultrasound in children for non-cardiac applications: a review by the Society for Pediatric Radiology (SPR) and the International Contrast Ultrasound Society (ICUS). *Pediatric radiology* 2013;43:1063-73.
48. Ntoulia A, Anupindi SA, Darge K, Back SJ. Applications of contrast-enhanced ultrasound in the pediatric abdomen. *Abdom Radiol (NY)* 2018;43:948-59.

*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

Development Chronology for this Practice Parameter

1995 (Resolution 8)

Amended 1995 (Resolution 24, 53)

Revised 1999 (Resolution 34)

Revised 2004 (Resolution 9)

Amended 2006 (Resolution 17, 35)

Revised 2009 (Resolution 33)

Revised 2014 (Resolution 13)

Revised 2019 (Resolution 10)

Revised 2024 (Resolution 31)

APPENDIX A

Grading System (Resolution 21) International Reflux Study of 1985 [31] (See Figure 7 from Pediatric Voiding Cystourethrography, a Pictorial Guide) [35]

1. Reflux only into the ureter.
2. Reflux into the entire ureter and pelvicalyceal system, no dilatation.
3. Mild pelvic or ureteral dilatation, with mild or no blunting of the fornices.
4. Moderate dilatation of the pelvis and ureter, with moderate dilation of the calyces.
5. Massive ureteral or pyelocalyceal dilatation.

Intrarenal reflux is NOT a component of the grading system and its presence does not alter the reflux grade (35, Fig8)

Grading system for ceVUS is similar to VCUG [44].