

# ACR-ASNR-SPR PRACTICE PARAMETER FOR THE PERFORMANCE OF COMPUTED TOMOGRAPHY ANGIOGRAPHY (CTA) OF THE HEAD AND NECK

## ACR-ASNR-SPR PRACTICE PARAMETER FOR THE PERFORMANCE OF COMPUTED TOMOGRAPHY ANGIOGRAPHY (CTA) OF THE HEAD AND NECK

The American College of Radiology, with more than 40,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

### PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care.<sup>1</sup> For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question. The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

---

<sup>1</sup> *Iowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing*, 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the "ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures (Revised 2008)" sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, *Stanley v. McCarver*, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

### I. INTRODUCTION

This practice parameter was revised collaboratively by the American College of Radiology (ACR), the American Society of Neuroradiology (ASNR), and the Society for Pediatric Radiology (SPR).

Computed tomography angiography (CTA) of the head and neck is a proven and useful procedure for the detection and characterization of vascular diseases and of vascular anatomy relevant to the treatment of extravascular disorders [1]. CTA may be used as the primary modality for detecting disease or as an adjunctive tool for characterizing known disease or assessing changes over time. When possible, magnetic resonance angiography (MRA) can be considered as an alternative to CTA to reduce radiation exposure, especially in the pediatric and vulnerable populations [2, 3]. Although it is not possible to detect all cerebrovascular abnormalities using CTA, adherence to the following practice parameter will maximize the probability of their detection and optimize patient safety.

CTA is a medical imaging technology that exposes patients to ionizing radiation. It should only be performed under the supervision of a physician with the necessary training in radiation biology and protection to optimize patient safety. Medical physicists and trained technical staff must be available.

CTA should be performed only for a valid medical indication and with the minimum exposure that provides the image quality necessary for adequate diagnostic information.

## **II. INDICATIONS**

Indications for CTA of the head and neck vessels include, but are not limited to, the diagnosis, characterization, and/or surveillance of:

1. Arterial aneurysms or pseudoaneurysms, venous varices, and dissections [2-10]
2. Ischemic stroke, transient ischemic attacks, vasospasm, and thromboembolism, including collateral assessment [9, 11-24]
3. Acute hemorrhage in the head or neck [25-29]
4. Atherosclerotic steno-occlusive disease, including atherosclerotic plaque localization and characterization [1, 30-39]
5. Nonatherosclerotic, noninflammatory vasculopathy, including radiation vasculopathy
6. Vasculitis and collagen vascular diseases [40]
7. Traumatic vascular injuries [3, 35, 41-50]
8. Venous and dural sinus thrombosis [51-53]
9. Vascular malformations and fistulas [54]
10. Pulsatile tinnitus [55]
11. Vascular anatomic variants [35, 56]
12. Evaluation for vascular intervention and follow-up (percutaneous and surgical) [57-71]
13. Tumors of vascular origin, with rich vascular supply or involving vascular structures [69, 72-76]
14. Surgical and radiation therapy localization, planning, and neuronavigation [71, 77]
15. To assess for vascular compression (eg bow-hunter's syndrome and Eagle syndrome) [78, 79]
16. Brain death in certain jurisdictions, especially when used in conjunction with other testing [80]
17. Postsurgical/posttreatment vascular complications

For certain indications, such as cerebral aneurysms and vasospasm, it may be appropriate to limit CTA to include only the head to avoid unnecessary radiation to the patient.

For the pregnant or potentially pregnant patient, see the [ACR-SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation](#) [81].

## **III. QUALIFICATIONS AND RESPONSIBILITIES OF PERSONNEL**

See the [ACR Practice Parameter for Performing and Interpreting Diagnostic Computed Tomography \(CT\)](#) [82].

### **A. Physician**

Examinations must be performed under the supervision of and interpreted by a physician who has the following qualifications:

The physician should meet the criteria listed in the [ACR Practice Parameter for Performing and Interpreting](#)

[Diagnostic Computed Tomography \(CT\)](#) and in the [ACR-SPR Practice Parameter for the Use of Intravascular Contrast Media](#) and should be trained in radiation safety [82, 83].

1. The physician is responsible for reviewing indications for the examination and for specifying the parameters of image acquisition; the route, volume, timing, type, and rate of contrast injection; and the method of image reconstruction and archival. The physician should monitor the quality of the images, be aware of potential artifacts [84], and interpret the study. Interpreting physicians must have knowledge of the benefits and risks of the procedures. Knowledge of the head and neck anatomy, including the vascular anatomy, and diseases of the intracranial and extracranial cerebrovascular system and their treatment is required.
2. Physicians meeting the aforementioned criteria additionally must have knowledge of the spectrum of nonvascular abnormalities presenting on CT scans. They should be capable of identifying and characterizing important nonvascular abnormalities that may be visualized on CTA, such as neoplasia, sequelae of infection, trauma, noninfectious inflammatory diseases, congenital anomalies, and normal anatomic variants, and any other abnormalities that may affect patient care and might necessitate treatment or further characterization through additional diagnostic testing.
3. The physician should be familiar with the use of 3-D processing workstations and be capable of performing or directing creation of 3-D renderings, multiplanar reformations, and measurements of vessel dimensions.
4. The physician should work with a Qualified Medical Physicist to optimize site-specific CTA scan protocols, when possible.

#### B. Technologist

The technologist should have the responsibility of patient comfort, preparing and positioning the patients for the CT examination, monitoring the patient during the examination, and obtaining the CT data in a manner prescribed by the supervising physician. For the intravenous (IV) administration of contrast material for CTA, qualifications for technologists performing IV injections should be in compliance with current ACR policy and existing operating procedures or manuals at the imaging facility. The technologist should perform the regular quality control testing of the CT system under the supervision of a medical physicist ([ACR-SPR Practice Parameter for the Use of Intravascular Contrast Media](#)) [83].

1. The technologist performing CT examinations should be certified by the American Registry of Radiologic Technologists or have an unrestricted state license with documented training and experience in CT.

### IV. SPECIFICATIONS OF THE EXAMINATION

CTA is a broad term. When referring to evaluation of arterial vessels, CTA is commonly used as an acronym. When CTA is used for evaluation of venous structures, it is typically denoted as a CT venogram (CTV). The equipment and contrast used for these examinations is the same. The scan protocols differ in the time delay to scanning following the injection of contrast.

The written or electronic request for a CTA of the head and neck should provide sufficient information to demonstrate the medical necessity of the examination and allow for the proper performance and interpretation of the examination.

Documentation that satisfies medical necessity includes 1) signs and symptoms and/or 2) relevant history (including known diagnoses). The provision of additional information regarding the specific reason for the examination or a provisional diagnosis would be helpful and may at times be needed to allow for the proper performance and interpretation of the examination.

The request for the examination must be originated by a physician or other appropriately licensed health care

provider. The accompanying clinical information should be provided by a physician or other appropriately licensed health care provider familiar with the patient's clinical problem or question and consistent with the state scope of practice requirements. (ACR Resolution 35 adopted in 2006 – revised in 2016, Resolution 12-b)

#### **IV. SPECIFICATIONS OF THE EXAMINATION**

##### **A. Patient Selection and Preparation**

Patients without absolute contraindication to the administration of iodinated contrast media are candidates for CTA of the head and neck. In cases of relative contraindication to the administration of iodinated contrast medium, measures to reduce the possibility of contrast medium reactions or nephrotoxicity should be followed to the extent that the patient's condition allows, as defined in the [ACR-SPR Practice Parameter for the Use of Intravascular Contrast Media](#), or an alternative vascular imaging modality should be considered, for example, MRA [83, 85].

When possible, patients should be well hydrated, and IV access should be established. A 20-gauge or larger antecubital IV catheter should be placed ideally on the right side (and preferably proximal to the wrist) to accommodate an optimal rate of 4–5 mL/s of iodinated contrast media. Small catheters that can withstand the prescribed injection rates can be used, and lower injection rates may be used for pediatric patients. All catheters used for the CTA examination should first be tested with a rapidly injected bolus of sterile saline to ensure that the venous access is secure and can accommodate the rapid bolus, minimizing the risk of contrast medium extravasations. The injection site should be monitored by medical personnel trained in the rapid recognition of IV extravasations. Department procedures for care of IV extravasations should be documented and applied if necessary.

#### **IV. SPECIFICATIONS OF THE EXAMINATION**

##### **B. CT Equipment**

The use of a multidetector-row CT scanner is required for CTA. The scanner must be capable of detecting and reliably displaying pathology in the adjacent structures and end organs of the vessels. Please refer to section VI for further equipment details.

A contrast medium power injector that allows programming of both the volume and flow rate must be used for head and neck CTA examinations.

Capability of creating multiplanar reformations, curved planar reformations, maximum-intensity projections, volume renderings, and/or shaded surface displays should be available for CTAs and applied to the appropriate study. The direct measurement of vascular diameters and, when appropriate, path lengths should also be available.

#### **IV. SPECIFICATIONS OF THE EXAMINATION**

##### **C. Examination Technique**

Before acquiring the CTA, a noncontrast head CT and/or noncontrast neck CT may be obtained, depending on the clinical suspicion, presentation, and acuity, for detecting hemorrhage, mapping of arterial calcification, or localization of the anatomy of interest. Similarly, once contrast has already been administered for the CTA, a delayed contrast-enhanced head and/or neck CT can be of value to detect areas of delayed/parenchymal enhancement, slow-flow lesions, spot sign not captured on the CTA, and/or for collateral assessment. Section thickness for these additional CT acquisitions is application-dependent but should not exceed 5 mm. The radiation exposure to the patient should be minimized within the limits of acceptable image quality, including optimization of peak kilovoltage (kVp) and mAs [86, 87]. In infants and children, weight- or age-appropriate guidelines should be used for acceptable CT radiation exposure to reflect the "as low as reasonably achievable" (ALARA) principle. If available, dose modulation and iterative reconstruction approaches should be used, with appropriate targeted signal-to-noise ratio [88, 89].

Because of substantial variations in the time required for an IV injection of nonionic contrast medium (iodine, 300–370 mg/mL) to reach the target vascular anatomy, an assessment of patient-specific circulation time is frequently required, especially for arterial imaging, although not mandatory. Circulation timing can be performed

using one of the following techniques [90]:

1. IV injection of a small test bolus (eg, 10-15 mL) of contrast medium at the same rate and through the same access that will be used for the CTA followed by acquisition of sequential cine CT images at the level of the artery or vein of interest. The rate and intensity of enhancement of the lumen of interest are then used to create a time density curve. The peak of the curve is used to calculate the scanning delay postinjection. A perfusion CT series performed before the CTA can be used similarly to a test bolus for determining the timing of the CTA acquisition.
2. The use of automated or semiautomated triggering software based on monitoring of the attenuation within the vessel of interest (or a great vessel such as the aorta) by the CT scanner following initiation of the full dose of contrast media injection may be used. The CT automatically starts when the enhancement in the vessel reaches a predetermined operator-selected level. An individualized posttrigger delay may optimize vessel opacification [91].
3. For CTV, administration of nonionic contrast medium with a 40–50-second prescanning delay, or up to a 30-second delay after the arterial bolus time, should allow adequate opacification of the venous structures minimizing flow artifacts.

Ideally, the administration of iodinated contrast media for the CTA should be performed with a minimum flow rate of 4 mL/s in any patient weighing 50 kg or more. Higher flow rates up to 6 mL/s are frequently required for larger patients, and in general, higher flow rates are required for shorter acquisitions. In children, contrast medium dosing should be scaled to body weight. Injection rate should be scaled similarly and preferably delivered via powered injection. For young children and infants, a 22- or 24-gauge IV catheter may be the only option, and a 2 mL/s injection rate may be reasonable for these patients. For patients under 50 kg, a dose of up to 2 mL/kg should be considered. In summary, contrast injection parameters should be modified on an individual patient basis, and the volume of contrast medium should be selected with consideration of the patient's weight and comorbidities that might increase the risk of contrast associated acute kidney injury. When performing CTA of the head and neck, a right-arm injection is preferable to a left-arm injection to avoid artifacts from undiluted contrast medium in the left brachiocephalic vein, and reflux of contrast due to venous narrowing at the level of the sternum. When possible, a bolus of saline following the iodinated contrast medium injection may reduce the volume of contrast medium required to achieve adequate vascular opacification.

The CTA head and neck acquisition should be performed with a section thickness of 1.5 mm or less, depending on the vascular territory to be assessed. The scan should be reconstructed with overlapping sections. For many indications, such as intracranial aneurysms, vasospasm, and cerebral venous thrombosis, imaging only needs to include the head. When CTA imaging of the neck is performed, such as in the setting of trauma, the acquisition should at least cover the aortic arch, the origin and cervical course of the subclavian and carotid arteries, and proximal subclavian arteries, through the skull base (eg, the floor of the sella). For many indications, such as stroke imaging, the acquisition should be extended through the circle of Willis and may be extended up to the cranial vertex. In the pediatric population, anatomic coverage should be strictly limited to the vascular segments of interest. Automated tube voltage selection can also be employed in conjunction with tube current modulation when available.

Utilization of iterative image reconstruction techniques, when available, is recommended to reduce image noise and artifacts, thereby allowing significant dose reduction [92]. Utilization of AI/deep learning techniques for image reconstruction may allow further dose reductions [93].

Postprocessing of the CTA by either physicians, radiologic technologists, or appropriately trained staff to provide multiplanar reformations and/or 3-D renderings is recommended [94]. Volume renderings, maximum-intensity projections, shaded surface displays, and curved planar reformations must be created by a person with knowledge of both head and neck vascular anatomy and pathology to avoid misrepresenting normal regions as diseased and vice versa. Segmentation of the CT data through a variety of manual and automated means may facilitate vascular visualization and measurement of stenosis, but it must be performed with care to avoid excluding key regions of the anatomy or creating pseudolesions. Pertinent measurements of vascular dimensions should be performed [95].

## IV. SPECIFICATIONS OF THE EXAMINATION

### D. Interpretation

CTAs of the head and neck are preferentially interpreted on equipment that allows stacked dynamic paging of the primary transverse and the reformatted CTA sections. A complete interpretation includes review of all images, including the scout and the transverse CT sections (source images) and, as indicated, multiplanar/curved reformations, volume renderings, maximum-intensity projections, and other images produced during postprocessing. The use of 3D reconstruction techniques (such as volume rendering or MIPs) is a required component of the examination. On occasion, the interpreting physician will personally create postprocessed images documenting important findings that are essential to the interpretation of the study [96]. These images should be archived with the patient's original study or other postprocessed images.

Interpretation of the head and neck CTA includes an assessment of the patency and caliber of the carotid and vertebral arteries, their origins, the carotid bifurcations, the intracranial arteries, possible occlusion, dissection, stenosis, and aneurysmal dilatation. To the extent that venous structures are adequately opacified on CTA images, as opposed to a dedicated delayed CTV, evaluation of images for venous pathology is also necessary. The visible and adequately opacified veins should be commented on when appropriate. Interpretation of dedicated head and neck CTV includes an assessment of the patency and caliber of the dural venous sinuses, cortical veins, and internal jugular veins. The visible and adequately opacified arteries should be commented on when appropriate.

The visible regional anatomy and pathology should be commented on when appropriate. The soft tissues and bony structures in the cervical region should be assessed in addition to the vessels. Bone kernel reconstruction images can be generated when appropriate. Comparison with previous studies should be performed when appropriate.

When applying the North American Symptomatic Carotid Endarterectomy Trial (NASCET) method for evaluation of cervical internal carotid artery stenosis, it is important for the interpreting physician to take into consideration that the denominator measurement needs to be done well beyond the tapering bulb, which tapers over a long distance, and should only be done where the vessel walls are parallel. An alternate method uses the residual lumen diameter measured in millimeters. This approach has been validated against the NASCET methodology and has been shown to be reproducible, to be easy to implement, and to provide similar information [94, 97-102].

## V. DOCUMENTATION

Reporting should be in accordance with the [ACR Practice Parameter for Communication of Diagnostic Imaging Findings](#) [103].

In addition to examining the head and neck vascular structures of interest, the CTA sections should be examined for clinically relevant extravascular abnormalities. These abnormalities should be described in the formal report of the examination.

## VI. EQUIPMENT SPECIFICATIONS

Equipment performance monitoring should be in accordance with the [ACR-AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Computed Tomography \(CT\) Equipment](#) [104].

For diagnostic quality CTA, the CT scanner should meet or exceed the following specifications:

1. Head and neck CTA should be performed on a multidetector CT (MDCT) scanner, with at least 16 active detector rows, with 64 or more detector rows being ideal.
2. Gantry rotation: 1 second or less for head and neck CTA.
3. Tube heat capacity that allows for a single  $\geq 10$ -second acquisition.
4. Capability of acquiring a section thickness of less than or equal to 1.5 mm.
5. Capability of helical or volume acquisitions with variable pitch.
6. Ability to create multiplanar images.
7. Image acquisition and reconstruction capabilities that allow for spatial resolutions of  $\geq 8$  lp/cm.

To maximize information available from the CT scan and thus derive the full diagnostic benefit for the patient

following X-ray irradiation, any CT scanner used for CTA must allow display and interpretation of the full 12 bits (from -1,000 to 3,095 Hounsfield units) of attenuation information (or possibly 16 bits for scans obtained on photon counting CT scanners). Additionally, the display field of view must be sufficient to allow an assessment of the vasculature of interest, the end-organ, and adjacent tissues. Dual-energy CTA can be obtained when available to decrease total patient radiation dose, lower contrast administration, distinguish contrast from hemorrhage and calcium, and reduce hardware artifacts [105-108]. Photon counting CT, when available, may provide increased quality by decreased noise and fewer artifacts in addition to the capability of material decomposition [109, 110].

Appropriate emergency equipment and medications must be immediately available to treat adverse reactions associated with administered medications. The equipment and medications should be monitored for inventory and drug expiration dates on a regular basis. The equipment, medications, and other emergency support must also be appropriate for the range of ages and sizes in the patient population.

## **VII. RADIATION SAFETY IN IMAGING**

Radiologists, medical physicists, non-physician radiology providers, radiologic technologists, and all supervising physicians have a responsibility for safety in the workplace by keeping radiation exposure to staff, and to society as a whole, "as low as reasonably achievable" (ALARA) and to assure that radiation doses to individual patients are appropriate, taking into account the possible risk from radiation exposure and the diagnostic image quality necessary to achieve the clinical objective. All personnel who work with ionizing radiation must understand the key principles of occupational and public radiation protection (justification, optimization of protection, application of dose constraints and limits) and the principles of proper management of radiation dose to patients (justification, optimization including the use of dose reference levels). [https://www-pub.iaea.org/MTCD/Publications/PDF/PUB1775\\_web.pdf](https://www-pub.iaea.org/MTCD/Publications/PDF/PUB1775_web.pdf)

Facilities and their responsible staff should consult with the radiation safety officer to ensure that there are policies and procedures for the safe handling and administration of radiopharmaceuticals in accordance with ALARA principles. These policies and procedures must comply with all applicable radiation safety regulations and conditions of licensure imposed by the Nuclear Regulatory Commission (NRC) and by applicable state, local, or other relevant regulatory agencies and accrediting bodies, as appropriate. Quantities of radiopharmaceuticals should be tailored to the individual patient by prescription or protocol, using body habitus or other customized method when such guidance is available.

Nationally developed guidelines, such as the [ACR's Appropriateness Criteria](#)®, should be used to help choose the most appropriate imaging procedures to prevent unnecessary radiation exposure.

Additional information regarding patient radiation safety in imaging is available from the following websites – Image Gently® for children ([www.imagegently.org](http://www.imagegently.org)) and Image Wisely® for adults ([www.imagewisely.org](http://www.imagewisely.org)). These advocacy and awareness campaigns provide free educational materials for all stakeholders involved in imaging (patients, technologists, referring providers, medical physicists, and radiologists).

Radiation exposures or other dose indices should be periodically measured by a Qualified Medical Physicist in accordance with the applicable ACR Technical Standards. Monitoring or regular review of dose indices from patient imaging should be performed by comparing the facility's dose information with national benchmarks, such as the ACR Dose Index Registry and relevant publications relying on its data, applicable ACR Practice Parameters, NCRP Report No. 172, Reference Levels and Achievable Doses in Medical and Dental Imaging: Recommendations for the United States or the Conference of Radiation Control Program Director's National Evaluation of X-ray Trends; 2006, 2009, amended 2013, revised 2023 (Res. 2d).

## **VIII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION**

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *ACR Position Statement on Quality Control and Improvement, Safety, Infection Control and Patient Education* on the ACR website (<https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement>).

## **ACKNOWLEDGEMENTS**

This practice parameter was developed according to the process described under the heading *The Process for Developing ACR Practice Parameters and Technical Standards* on the ACR website (<https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards>) by the Committee on Practice Parameters – Neuroradiology of the ACR Commission on Neuroradiology and the Committee on Practice Parameters – Pediatric Imaging of the ACR Commission on Pediatric Radiology in collaboration with the ASNR, and the SPR. Writing Committee – members represent their societies in the initial and final revision of this practice parameter

ACR

Joyner, David A MD, Chair  
Isikbay, Masis BS, MD  
Lasiecka, Zofia M MD, PhD  
Maloney, John A MD  
Ormsby, Jacob MBA, MD

ASNR

Lin, Doris D.M. MD, PhD  
Nemeth, Alexander J MD  
Segovis, Colin MD, PhD  
Singhal, Aparna MD

SPR

Narayanan, Srikala MD  
Powers, Andria MD  
Radhakrishnan, Rupa MBBS, MS

Committee on Practice Parameters – Neuroradiology

(ACR Committee responsible for sponsoring the draft through the process)

Shah, Lubdha M MD, Chair  
Amrhein, Timothy J MD  
Becker, Jennifer L MD  
Chaudhry, Umar S MD  
Hutchins, Troy A MD  
Ormsby, Jacob MBA, MD  
Sio, Terence Tai-Weng MD, MS

Aiken, Ashley H MD  
Austin, Matthew J MD  
Blackham, Kristine A MD  
Chiang, Gloria C MD  
Mazaheri, Parisa MD  
Riley, Kalen MD

Committee on Practice Parameters and Technical Standards

(ACR Committee responsible for sponsoring the draft through the process)

Caplin, Drew M MD, Chair

Committee on Practice Parameters – Pediatric Imaging

(ACR Committee responsible for sponsoring the draft through the process)

Amodio, John B MD, Chair  
Bhimaniya, Sudhir X MBBS, MD  
Collard, Michael MD  
Lai, Hollie A MD  
Lasiecka, Zofia M MD, PhD  
Li, Arleen MD  
Noda, Sakura MD  
Vatsky, Seth DO

Alizai, Hamza MD  
Blumfield, Einat MD  
Goldman-Yassen, Adam MD  
Lala, Shailee V MD  
Laufer, Adina MD  
Maloney, John A MD  
Shah, Summit MD

Bulas, Dorothy I MD, Chair, Commission on Pediatric Radiology

Jordan, John E MD, Chair, Commission on Neuroradiology

Larson, David B MBA, MD, Chair, Commission on Quality and Safety

Comments Reconciliation Committee

Chu, Sammy MD - CSC, Co-Chair  
Bulas, Dorothy I MD  
Isikbay, Masis BS, MD  
Joyner, David A MD  
Lasiecka, Zofia M MD, PhD

Amodio, John B MD  
Caplin, Drew M MD  
Jordan, John E MD  
Larson, David B MBA, MD  
Lin, Doris D.M. MD, PhD

Maloney, John A MD  
Nemeth, Alexander J MD  
Powers, Andria MD  
Segovis, Colin MD, PhD  
Singhal, Aparna MD

Narayanan, Srikala MD  
Ormsby, Jacob MBA, MD  
Radhakrishnan, Rupa MBBS, MS  
Shah, Lubdha M MD

## REFERENCES

1. Brown DL, Hoffman SN, Jacobs TL, Gruis KL, Johnson SL, Chernew ME. CT angiography is cost-effective for confirmation of internal carotid artery occlusions. *J Neuroimaging*. 2008 Oct;18(4):355-9.
2. Hellinger JC, Pena A, Poon M, Chan FP, Epelman M. Pediatric computed tomographic angiography: imaging the cardiovascular system gently. *Radiol Clin North Am*. 2010 Mar;48(2):439-67, x.
3. Stence NV, Fenton LZ, Goldenberg NA, Armstrong-Wells J, Bernard TJ. Craniocervical arterial dissection in children: diagnosis and treatment. *Curr Treat Options Neurol*. 2011;13(6):636-648.
4. Chen W, Wang J, Xin W, Peng Y, Xu Q. Accuracy of 16-row multislice computed tomographic angiography for assessment of small cerebral aneurysms. *Neurosurgery*. 2008 Jan;62(1):113-21; discussion 121-2.
5. Dammert S, Krings T, Moller-Hartmann W, et al. Detection of intracranial aneurysms with multislice CT: comparison with conventional angiography. *Neuroradiology*. 2004;46(6):427-434.
6. Jayaraman MV, Mayo-Smith WW, Tung GA, et al. Detection of intracranial aneurysms: multi-detector row CT angiography compared with DSA. *Radiology*. 2004;230(2):510-518.
7. McKinney AM, Palmer CS, Truwit CL, Karagulle A, Teksam M. Detection of aneurysms by 64-section multidetector CT angiography in patients acutely suspected of having an intracranial aneurysm and comparison with digital subtraction and 3D rotational angiography. *AJNR Am J Neuroradiol*. 2008;29(3):594-602.
8. Teksam M, McKinney A, Casey S, Asis M, Kieffer S, Truwit CL. Multi-section CT angiography for detection of cerebral aneurysms. *AJNR Am J Neuroradiol*. 2004 Oct;25(9):1485-92.
9. Mehan WA Jr, Romero JM, Hirsch JA, et al. Unruptured intracranial aneurysms conservatively followed with serial CT angiography: could morphology and growth predict rupture?. *J Neurointerv Surg*. 6(10):761-6, 2014 Dec.
10. Wang GX, Gong MF, Wen L, et al. Computed Tomography Angiography Evaluation of Risk Factors for Unstable Intracranial Aneurysms. *World Neurosurg*. 2018 Jul;115():S1878-8750(18)30561-8.
11. Chaudhary SR, Ko N, Dillon WP, et al. Prospective evaluation of multidetector-row CT angiography for the diagnosis of vasospasm following subarachnoid hemorrhage: a comparison with digital subtraction angiography. *Cerebrovasc Dis*. 2008;25(1-2):144-50.
12. Coutts SB, Lev MH, Eliasziw M, et al. ASPECTS on CTA source images versus unenhanced CT: added value in predicting final infarct extent and clinical outcome. *Stroke*. 2004 Nov;35(11):2472-6.
13. Josephson SA, Bryant SO, Mak HK, Johnston SC, Dillon WP, Smith WS. Evaluation of carotid stenosis using CT angiography in the initial evaluation of stroke and TIA. *Neurology*. 2004 Aug 10;63(3):457-60.
14. Tan JC, Dillon WP, Liu S, Adler F, Smith WS, Wintermark M. Systematic comparison of perfusion-CT and CT-angiography in acute stroke patients. *Ann Neurol*. 2007 Jun;61(6):533-43.
15. Torres-Mozqueda F, He J, Yeh IB, et al. An acute ischemic stroke classification instrument that includes CT or MR angiography: the Boston Acute Stroke Imaging Scale. *AJNR Am J Neuroradiol*. 2008 Jun;29(6):1111-7.
16. Wintermark M, Uske A, Chalaron M, et al. Multislice computerized tomography angiography in the evaluation of intracranial aneurysms: a comparison with intraarterial digital subtraction angiography. *J Neurosurg*. 2003 Apr;98(4):828-36.
17. Yoon DY, Lim KJ, Choi CS, Cho BM, Oh SM, Chang SK. Detection and characterization of intracranial aneurysms with 16-channel multidetector row CT angiography: a prospective comparison of volume-rendered images and digital subtraction angiography. *AJNR Am J Neuroradiol*. 2007 Jan;28(1):60-7.
18. Chatzikonstantinou A, Krissak R, Flüchter S, et al. CT angiography of the aorta is superior to transesophageal echocardiography for determining stroke subtypes in patients with cryptogenic ischemic stroke. *Cerebrovasc Dis*. 2012;33(4):322-8.
19. Furtado AD, Adraktas DD, Brasic N, et al. The triple rule-out for acute ischemic stroke: imaging the brain, carotid arteries, aorta, and heart. *AJNR Am J Neuroradiol*. 2010 Aug;31(7):1290-6.
20. Greenberg ED, Gold R, Reichman M, et al. Diagnostic accuracy of CT angiography and CT perfusion for cerebral vasospasm: a meta-analysis. *AJNR Am J Neuroradiol*. 2010;31(10):1853-1860.
21. Ritter MA, Poeplau T, Schaefer A, et al. CT angiography in acute stroke: does it provide additional information on occurrence of infarction and functional outcome after 3 months?. *Cerebrovasc Dis*. 2006;22(5-6):362-7.

**22.** Wintermark M, Sanelli PC, Albers GW, et al. Imaging recommendations for acute stroke and transient ischemic attack patients: A joint statement by the American Society of Neuroradiology, the American College of Radiology, and the Society of NeuroInterventional Surgery. *AJNR Am J Neuroradiol* 2013;34:E117-27.

**23.** Sundaram S, Kannoth S, Thomas B, Sarma PS, Sylaja PN. Collateral Assessment by CT Angiography as a Predictor of Outcome in Symptomatic Cervical Internal Carotid Artery Occlusion. *AJNR Am J Neuroradiol*. 38(1):52-57, 2017 Jan.

**24.** Varadharajan S, Saini J, Acharya UV, Gupta AK. Computed tomography angiography in acute stroke (revisiting the 4Ps of imaging). *Am J Emerg Med*. 2016 Feb;34(2):S0735-6757(15)00949-3.

**25.** Kim J, Smith A, Hemphill JC, et al. Contrast extravasation on CT predicts mortality in primary intracerebral hemorrhage. *AJNR Am J Neuroradiol*. 2008 Mar;29(3):520-5.

**26.** Romero JM, Artunduaga M, Forero NP, et al. Accuracy of CT angiography for the diagnosis of vascular abnormalities causing intraparenchymal hemorrhage in young patients. *Emerg Radiol*. 2009 May;16(3):195-201.

**27.** Chakraborty S, Alhazzaa M, Wasserman JK, et al. Dynamic characterization of the CT angiographic 'spot sign'. *PLoS One*. 2014;9(3):e90431.

**28.** Demchuk AM, Dowlatshahi D, Rodriguez-Luna D, et al. Prediction of haematoma growth and outcome in patients with intracerebral haemorrhage using the CT-angiography spot sign (PREDICT): a prospective observational study. *Lancet Neurol*. 2012 Apr;11(4):307-14.

**29.** Rodriguez-Luna D, Dowlatshahi D, Aviv RI, et al. Venous phase of computed tomography angiography increases spot sign detection, but intracerebral hemorrhage expansion is greater in spot signs detected in arterial phase. *Stroke*. 2014 Mar;45(3):734-9.

**30.** Bartlett ES, Walters TD, Symons SP, Fox AJ. Diagnosing carotid stenosis near-occlusion by using CT angiography. *AJNR Am J Neuroradiol*. 2006 Mar;27(3):632-7.

**31.** Prokop M, Waaijer A, Kreuzer S. CT angiography of the carotid arteries. *JBR-BTR*. 2004;87(1):23-9.

**32.** Randoux B, Marro B, Marsault C. Carotid Artery Stenosis: Competition between CT Angiography and MR Angiography. *AJNR Am J Neuroradiol*. 2004 Apr;25(4):663-4; author reply 664.

**33.** Wintermark M, Ko NU, Smith WS, Liu S, Higashida RT, Dillon WP. Vasospasm after subarachnoid hemorrhage: utility of perfusion CT and CT angiography on diagnosis and management. *AJNR Am J Neuroradiol*. 2006 Jan;27(1):26-34.

**34.** Wintermark M, Meuli R, Browaeys P, et al. Comparison of CT perfusion and angiography and MRI in selecting stroke patients for acute treatment. *Neurology*. 2007 Feb 27;68(9):694-7.

**35.** Barazangi N, Wintermark M, Lease K, Rao R, Smith W, Josephson SA. Comparison of computed tomography angiography and transesophageal echocardiography for evaluating aortic arch disease. *J Stroke Cerebrovasc Dis*. 2011;20(5):436-42.

**36.** Magge R, Lau BC, Soares BP, et al. Clinical risk factors and CT imaging features of carotid atherosclerotic plaques as predictors of new incident carotid ischemic stroke: a retrospective cohort study. *AJNR Am J Neuroradiol*. 2013 Feb;34(2):402-9.

**37.** Romero JM, Babiarz LS, Forero NP, et al. Arterial wall enhancement overlying carotid plaque on CT angiography correlates with symptoms in patients with high grade stenosis. *Stroke*. 2009 May;40(5):1894-6.

**38.** Wintermark M, Jawadi SS, Rapp JH, et al. High-resolution CT imaging of carotid artery atherosclerotic plaques. *AJNR Am J Neuroradiol*. 2008;29(5):875-882.

**39.** Sun PP, Feng PY, Wang Q, Shen SS. Angiography with the 256-multislice spiral computed tomography and its application in evaluating atherosclerotic plaque and cerebral ischemia. *Medicine (Baltimore)*. 97(30):e11408, 2018 Jul.

**40.** Schmidt WA. Imaging in vasculitis. *Best Pract Res Clin Rheumatol*. 2013 Feb;27(1):S1521-6942(13)00003-X.

**41.** Berne JD, Norwood SH, McAuley CE, Villareal DH. Helical computed tomographic angiography: an excellent screening test for blunt cerebrovascular injury. *J Trauma*. 57(1):11-7; discussion 17-9, 2004 Jul.

**42.** Nguyen-Huynh MN, Wintermark M, English J, et al. How accurate is CT angiography in evaluating intracranial atherosclerotic disease?. *Stroke*. 2008 Apr;39(4):1184-8.

**43.** van Prehn J, Muhs BE, Pramanik B, et al. Multidimensional characterization of carotid artery stenosis using CT imaging: a comparison with ultrasound grading and peak flow measurement. *Eur J Vasc Endovasc Surg*. 2008 Sep;36(3):267-72.

**44.** Methodius-Ngwodo WC, Burkett AB, Kochupura PV, Wellons ED, Fuhrman G, Rosenthal D. The role of CT angiography in the diagnosis of blunt traumatic thoracic aortic disruption and unsuspected carotid artery injury. *Am Surg*. 2008;74(7):580-585; discussion 585-586.

**45.** Langner S, Fleck S, Kirsch M, Petrik M, Hosten N. Whole-body CT trauma imaging with adapted and optimized CT angiography of the craniocervical vessels: do we need an extra screening examination?. *AJNR Am J Neuroradiol*. 2014;35(10):1933-1938.

Neuroradiol. 29(10):1902-7, 2008 Nov.

**46.** Roberts DJ, Chaubey VP, Zygun DA, et al. Diagnostic accuracy of computed tomographic angiography for blunt cerebrovascular injury detection in trauma patients: a systematic review and meta-analysis. [Review]. Annals of Surgery. 257(4):621-32, 2013 Apr.

**47.** Tessler RA, Nguyen H, Newton C, Betts J. Pediatric penetrating neck trauma: Hard signs of injury and selective neck exploration. J Trauma Acute Care Surg. 2017 Jun;82(6):989-994.

**48.** Todnem N, Hardigan T, Banerjee C, Alleyne CH. Cephalad Migration of Intradural Bullet from Thoracic Spine to Cervical Spine. World Neurosurg. 2018 Nov;119():S1878-8750(18)31414-1.

**49.** Ugalde IT, Claiborne MK, Cardenas-Turanzas M, Shah MN, Langabeer JR 2nd, Patel R. Risk Factors in Pediatric Blunt Cervical Vascular Injury and Significance of Seatbelt Sign. The Western Journal of Emergency Medicine. 19(6):961-969, 2018 Nov.

**50.** Ibraheem K, Wong S, Smith A, et al. Computed tomography angiography in the "no-zone" approach era for penetrating neck trauma: A systematic review. The Journal of Trauma and Acute Care Surgery. 89(6):1233-1238, 2020 12.

**51.** Leach JL, Fortuna RB, Jones BV, Gaskill-Shipley MF. Imaging of cerebral venous thrombosis: current techniques, spectrum of findings, and diagnostic pitfalls. Radiographics. 2006;26 Suppl 1:S19-41; discussion S42-13.

**52.** Taschner C, Leclerc X, Lucas C, Pruvost J. Computed tomography angiography for the evaluation of carotid artery dissections. Front Neurol Neurosci. 2005;20():119-128.

**53.** Vertinsky AT, Schwartz NE, Fischbein NJ, Rosenberg J, Albers GW, Zaharchuk G. Comparison of multidetector CT angiography and MR imaging of cervical artery dissection. AJNR Am J Neuroradiol. 2008;29(9):1753-1760.

**54.** Singh R, Gupta V, Ahuja C, Kumar A, Mukherjee KK, Khandelwal N. Role of time-resolved-CTA in intracranial arteriovenous malformation evaluation at 128-slice CT in comparison with digital subtraction angiography. Neuroradiol J. 2018 Jun;31(3):235-243.

**55.** Xu S, Ruan S, Liu S, Xu J, Gong R. CTA/V detection of bilateral sigmoid sinus dehiscence and suspected idiopathic intracranial hypertension in unilateral pulsatile tinnitus. Neuroradiology. 60(4):365-372, 2018 Apr.

**56.** Sivaraju L, Mani S, Prabhu K, Daniel RT, Chacko AG. Three-dimensional computed tomography angiographic study of the vertebral artery in patients with congenital cranivertebral junction anomalies. Eur Spine J. 2017 Apr;26(4):1028-1038.

**57.** Chen W, Yang Y, Xing W, Qiu J, Peng Y. Sixteen-row multislice computed tomography angiography in the diagnosis and characterization of intracranial aneurysms: comparison with conventional angiography and intraoperative findings. J Neurosurg. 2008 Jun;108(6):1184-91.

**58.** Dehdashti AR, Binaghi S, Uske A, Regli L. Comparison of multislice computerized tomography angiography and digital subtraction angiography in the postoperative evaluation of patients with clipped aneurysms. J Neurosurg. 2006 Mar;104(3):395-403.

**59.** Katano H, Kato K, Umemura A, Yamada K. Perioperative evaluation of carotid endarterectomy by 3D-CT angiography with refined reconstruction: preliminary experience of CEA without conventional angiography. Br J Neurosurg. 2004 Apr;18(2):138-48.

**60.** Lev MH, Farkas J, Rodriguez VR, et al. CT angiography in the rapid triage of patients with hyperacute stroke to intraarterial thrombolysis: accuracy in the detection of large vessel thrombus. J Comput Assist Tomogr. 2001;25(4):520-8.

**61.** Ozsvath RR, Casey SO, Lustrin ES, Alberico RA, Hassankhani A, Patel M. Cerebral venography: comparison of CT and MR projection venography. AJR Am J Roentgenol 1997;169:1699-707.

**62.** Papke K, Kuhl CK, Fruth M, et al. Intracranial aneurysms: role of multidetector CT angiography in diagnosis and endovascular therapy planning. Radiology. 2007 Aug;244(2):532-40.

**63.** Rosenthal ES, Schwamm LH, Roccatagliata L, et al. Role of recanalization in acute stroke outcome: rationale for a CT angiogram-based "benefit of recanalization" model. AJNR Am J Neuroradiol. 2008 Sep;29(8):1471-5.

**64.** Wetzel SG, Kirsch E, Stock KW, Kolbe M, Kaim A, Radue EW. Cerebral veins: comparative study of CT venography with intraarterial digital subtraction angiography. AJNR Am J Neuroradiol. 1999 Feb;20(2):249-55.

**65.** Bal S, Bhatia R, Shobha N, et al. Stroke- on- Awakening: Safety of CT-CTA Based Selection for Reperfusion Therapy. Can J Neurol Sci. 41(2):182-6, 2014 Mar.

**66.** Fiebach JB, Al-Rawi Y, Wintermark M, et al. Vascular occlusion enables selecting acute ischemic stroke patients for treatment with desmoteplase. Stroke. 2012 Jun;43(6):1561-6.

**67.** González RG, Furie KL, Goldmacher GV, et al. Good outcome rate of 35% in IV-tPA-treated patients with computed tomography angiography confirmed severe anterior circulation occlusive stroke. Stroke. 2013 Nov;44(11):3109-13.

**68.** Nambiar V, Sohn SI, Almekhlafi MA, et al. CTA collateral status and response to recanalization in patients with

acute ischemic stroke. *AJNR Am J Neuroradiol.* 35(5):884-90, 2014 May.

**69.** Schaefer PW, Yoo AJ, Bell D, et al. CT angiography-source image hypoattenuation predicts clinical outcome in posterior circulation strokes treated with intra-arterial therapy. *Stroke.* 2008 Nov;39(11):3107-9.

**70.** Storey C, Barry J, Adkins W, Nanda A, Saenz-Cuellar H. A Morphometric Analysis for the Feasibility of Percutaneous Translacerum Access of the Internal Carotid Artery Based on Computed Tomography Angiography. *World Neurosurg.* 121:e925-e930, 2019 Jan.

**71.** Wada K, Nawashiro H, Ohkawa H, Arimoto H, Takeuchi S, Mori K. Feasibility of the combination of 3D CTA and 2D CT imaging guidance for clipping microsurgery of anterior communicating artery aneurysm. *Br J Neurosurg.* 2015 Apr;29(2):229-36.

**72.** Goodman DN, Hoh BL, Rabinov JD, Pryor JC. CT angiography before embolization for hemorrhage in head and neck cancer. *AJNR Am J Neuroradiol.* 2003 Jan;24(1):140-2.

**73.** Kramer M, Vairaktaris E, Nkenke E, Schlegel KA, Neukam FW, Lell M. Vascular mapping of head and neck: computed tomography angiography versus digital subtraction angiography. *J Oral Maxillofac Surg.* 2008 Feb;66(2):302-7.

**74.** Sims JR, Rordorf G, Smith EE, et al. Arterial occlusion revealed by CT angiography predicts NIH stroke score and acute outcomes after IV tPA treatment. *AJNR Am J Neuroradiol.* 2005 Feb;26(2):246-51.

**75.** Swieton D, Kaszubowski M, Szyndler A, et al. Visualizing Carotid Bodies With Doppler Ultrasound Versus CT Angiography: Preliminary Study. *AJR Am J Roentgenol.* 2017 Dec;209(6):1348-1352.

**76.** Xiao Z, Zheng Y, Li J, Chen D, Liu F, Cao D. Four-dimensional CT angiography (4D-CTA) in the evaluation of juvenile nasopharyngeal angiofibromas: comparison with digital subtraction angiography (DSA) and surgical findings. *Dentomaxillofac Radiol.* 2017 Dec;46(8):20170171.

**77.** Srinivasan VM, Schafer S, Ghali MG, Arthur A, Duckworth EA. Cone-beam CT angiography (Dyna CT) for intraoperative localization of cerebral arteriovenous malformations. *J Neurointerv Surg.* 8(1):69-74, 2016 Jan.

**78.** Jost GF, Dailey AT. Bow hunter's syndrome revisited: 2 new cases and literature review of 124 cases. *Neurosurg Focus* 2015;38:E7.

**79.** Chuang WC, Short JH, McKinney AM, Anker L, Knoll B, McKinney ZJ. Reversible left hemispheric ischemia secondary to carotid compression in Eagle syndrome: surgical and CT angiographic correlation. *AJNR Am J Neuroradiol.* 2007 Jan;28(1):143-5.

**80.** Suarez-Kelly LP, Patel DA, Britt PM, et al. Dead or alive? New confirmatory test using quantitative analysis of computed tomographic angiography. *J Trauma Acute Care Surg.* 2015 Dec;79(6):995-1003; discussion 1003.

**81.** American College of Radiology. ACR-SPR Practice Parameter for Imaging Pregnant or Potentially Pregnant Patients with Ionizing Radiation. Available at: <https://gravitas.acr.org/PPTS/GetDocumentView?docId=23+&releaseld=2>.

**82.** American College of Radiology. ACR-SPR Practice Parameter for Performing and Interpreting Diagnostic Computed Tomography (CT). Available at <https://gravitas.acr.org/PPTS/GetDocumentView?docId=132+&releaseld=2>

**83.** American College of Radiology. ACR-SPR Practice Parameter for the use of Intravascular Contrast Media. Available at <https://gravitas.acr.org/PPTS/GetDocumentView?docId=142+&releaseld=2>

**84.** Kim JJ, Dillon WP, Glastonbury CM, Provenzale JM, Wintermark M. Sixty-four-section multidetector CT angiography of carotid arteries: a systematic analysis of image quality and artifacts. *AJNR Am J Neuroradiol.* 2010 Jan;31(1):91-9.

**85.** Oleinik A, Romero JM, Schwab K, et al. CT angiography for intracerebral hemorrhage does not increase risk of acute nephropathy. *Stroke.* 40(7):2393-7, 2009 Jul.

**86.** Loftus ML, Minkowitz S, Tsioris AJ, Min RJ, Sanelli PC. Utilization guidelines for reducing radiation exposure in the evaluation of aneurysmal subarachnoid hemorrhage: A practice quality improvement project. *AJR Am J Roentgenol.* 2010 Jul;195(1):176-80.

**87.** Siegel MJ, Hildebolt C, Bradley D. Effects of automated kilovoltage selection technology on contrast-enhanced pediatric CT and CT angiography. *Radiology.* 2013 Aug;268(2):538-47.

**88.** Komlosi P, Zhang Y, Leiva-Salinas C, et al. Adaptive statistical iterative reconstruction reduces patient radiation dose in neuroradiology CT studies. *Neuroradiology.* 2014 Mar;56(3):187-93.

**89.** Singh S, Kalra MK, Shenoy-Bhangle AS, et al. Radiation dose reduction with hybrid iterative reconstruction for pediatric CT. *Radiology.* 2012 May;263(2):537-46.

**90.** Lell M, Tomandl BF, Anders K, Baum U, Nkenke E. Computed tomography angiography versus digital subtraction angiography in vascular mapping for planning of microsurgical reconstruction of the mandible. *Eur Radiol.* 2005 Aug;15(8):1514-20.

**91.** Yuan D, Li L, Zhang Y, et al. Image quality improvement in head and neck CT angiography: Individualized post-

trigger delay versus fixed delay. *Eur J Radiol*. 2023 Nov;168():S0720-048X(23)00456-4.

**92.** Dieckmeyer M, Sollmann N, Kupfer K, et al. Computed Tomography of the Head : A Systematic Review on Acquisition and Reconstruction Techniques to Reduce Radiation Dose. *Clin Neuroradiol*. 2023 Sep;33(3):591-610.

**93.** Huang X, Zhao W, Wang G, et al. Improving image quality with deep learning image reconstruction in double-low-dose head CT angiography compared with standard dose and adaptive statistical iterative reconstruction. *Br J Radiol*. 2023 Mar;96(1143):20220625.

**94.** Sparacia G, Bencivinni F, Banco A, Sarno C, Bartolotta TV, Lagalla R. Imaging processing for CT angiography of the cervicocranial arteries: evaluation of reformatting technique. *Radiol Med*. 2007 Mar;112(2):224-38.

**95.** Wintermark M, Glastonbury C, Tong E, et al. Semi-automated computer assessment of the degree of carotid artery stenosis compares favorably to visual evaluation. *J Neurol Sci*. 2008 Jun 15;269(1-2):74-9.

**96.** Leong JL, Batra PS, Citardi MJ. Three-dimensional computed tomography angiography of the internal carotid artery for preoperative evaluation of sinonasal lesions and intraoperative surgical navigation. *Laryngoscope*. 115(9):1618-23, 2005 Sep.

**97.** Bartlett ES, Symons SP, Fox AJ. Correlation of carotid stenosis diameter and cross-sectional areas with CT angiography. *AJNR Am J Neuroradiol*. 2006 Mar;27(3):638-42.

**98.** Bartlett ES, Walters TD, Symons SP, Aviv RI, Fox AJ. Classification of carotid stenosis by millimeter CT angiography measures: effects of prevalence and gender. *AJNR Am J Neuroradiol*. 2008 Oct;29(9):1677-83.

**99.** Bartlett ES, Walters TD, Symons SP, Fox AJ. Quantification of carotid stenosis on CT angiography. *AJNR Am J Neuroradiol*. 2006 Jan;27(1):13-9.

**100.** Bartlett ES, Walters TD, Symons SP, Fox AJ. Carotid stenosis index revisited with direct CT angiography measurement of carotid arteries to quantify carotid stenosis. *Stroke*. 2007 Feb;38(2):286-91.

**101.** Bucek RA, Puchner S, Haumer M, Reiter M, Minar E, Lammer J. CTA quantification of internal carotid artery stenosis: application of luminal area vs. luminal diameter measurements and assessment of inter-observer variability. *J Neuroimaging*. 2007 Jul;17(3):219-26.

**102.** Cademartiri F, Nieman K, van der Lugt A, et al. Intravenous contrast material administration at 16-detector row helical CT coronary angiography: test bolus versus bolus-tracking technique. *Radiology*. 2004 Dec;233(3):817-23.

**103.** American College of Radiology. ACR Practice Parameter for Communication of Diagnostic Imaging Findings. Available at <https://gravitas.acr.org/PPTS/GetDocumentView?docId=74+&releaseld=2>

**104.** American College of Radiology. ACR–AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Computed Tomography (CT) Equipment. Available at <https://gravitas.acr.org/PPTS/GetDocumentView?docId=131+&releaseld=2>

**105.** Zhao L, Li F, Zhang Z, et al. Assessment of an advanced virtual monoenergetic reconstruction technique in cerebral and cervical angiography with third-generation dual-source CT: Feasibility of using low-concentration contrast medium. *Eur Radiol*. 2018 Oct;28(10):4379-4388.

**106.** Ferda J, Novák M, Mírka H, et al. The assessment of intracranial bleeding with virtual unenhanced imaging by means of dual-energy CT angiography. *Eur Radiol*. 2009 Oct;19(10):2518-22.

**107.** Jiang XY, Zhang SH, Xie QZ, et al. Evaluation of Virtual Noncontrast Images Obtained from Dual-Energy CTA for Diagnosing Subarachnoid Hemorrhage. *AJNR Am J Neuroradiol*. 2015 May;36(5):855-60.

**108.** Kamalian S, Lev MH, Pomerantz SR. Dual-Energy Computed Tomography Angiography of the Head and Neck and Related Applications. *Neuroimaging Clin N Am*. 2017 Aug;27(3):S1052-5149(17)30044-8.

**109.** Symons R, Reich DS, Bagheri M, et al. Photon-Counting Computed Tomography for Vascular Imaging of the Head and Neck: First In Vivo Human Results. *Invest Radiol*. 2018 Mar;53(3):135-142.

**110.** Madhavan AA, Bathla G, Benson JC, Diehn FE, Nagelschneider AA, Lehman VT. High yield clinical applications for photon counting CT in neurovascular imaging. *Br J Radiol*. 2024 May 07;97(1157):894-901.

\*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

#### Development Chronology for this Practice Parameter

2010 (Resolution 20)

Amended 2014 (Resolution 39)

Revised 2015 (Resolution 19)

Revised 2020 (Resolution 42)

Amended 2023 (Resolution 2c, 2d)

Revised 2025 (Resolution 16)

Revised 2025 (Resolution 16)