

ACR–ACOG–AIUM–SPR–SRU PRACTICE PARAMETER FOR THE PERFORMANCE OF ULTRASOUND OF THE FEMALE PELVIS

The American College of Radiology, with more than 40,000 members, is the principal organization of radiologists, radiation oncologists, and clinical medical physicists in the United States. The College is a nonprofit professional society whose primary purposes are to advance the science of radiology, improve radiologic services to the patient, study the socioeconomic aspects of the practice of radiology, and encourage continuing education for radiologists, radiation oncologists, medical physicists, and persons practicing in allied professional fields.

The American College of Radiology will periodically define new practice parameters and technical standards for radiologic practice to help advance the science of radiology and to improve the quality of service to patients throughout the United States. Existing practice parameters and technical standards will be reviewed for revision or renewal, as appropriate, on their fifth anniversary or sooner, if indicated.

Each practice parameter and technical standard, representing a policy statement by the College, has undergone a thorough consensus process in which it has been subjected to extensive review and approval. The practice parameters and technical standards recognize that the safe and effective use of diagnostic and therapeutic radiology requires specific training, skills, and techniques, as described in each document. Reproduction or modification of the published practice parameter and technical standard by those entities not providing these services is not authorized.

PREAMBLE

This document is an educational tool designed to assist practitioners in providing appropriate radiologic care for patients. Practice Parameters and Technical Standards are not inflexible rules or requirements of practice and are not intended, nor should they be used, to establish a legal standard of care¹. For these reasons and those set forth below, the American College of Radiology and our collaborating medical specialty societies caution against the use of these documents in litigation in which the clinical decisions of a practitioner are called into question.

The ultimate judgment regarding the propriety of any specific procedure or course of action must be made by the practitioner considering all the circumstances presented. Thus, an approach that differs from the guidance in this document, standing alone, does not necessarily imply that the approach was below the standard of care. To the contrary, a conscientious practitioner may responsibly adopt a course of action different from that set forth in this document when, in the reasonable judgment of the practitioner, such course of action is indicated by variables such as the condition of the patient, limitations of available resources, or advances in knowledge or technology after publication of this document. However, a practitioner who employs an approach substantially different from the guidance in this document may consider documenting in the patient record information sufficient to explain the approach taken.

The practice of medicine involves the science, and the art of dealing with the prevention, diagnosis, alleviation, and treatment of disease. The variety and complexity of human conditions make it impossible to always reach the most appropriate diagnosis or to predict with certainty a particular response to treatment. Therefore, it should be recognized that adherence to the guidance in this document will not assure an accurate diagnosis or a successful outcome. All that should be expected is that the practitioner will follow a reasonable course of action based on current knowledge, available resources, and the needs of the patient to deliver effective and safe medical care. The purpose of this document is to assist practitioners in achieving this objective.

¹ *Iowa Medical Society and Iowa Society of Anesthesiologists v. Iowa Board of Nursing*, 831 N.W.2d 826 (Iowa 2013) Iowa Supreme Court refuses to find that the "ACR Technical Standard for Management of the Use of Radiation in Fluoroscopic Procedures (Revised 2008)" sets a national standard for who may perform fluoroscopic procedures in light of the standard's stated purpose that ACR standards are educational tools and not intended to establish a legal standard of care. See also, *Stanley v. McCarver*, 63 P.3d 1076 (Ariz. App. 2003) where in a concurring opinion the Court stated that "published standards or guidelines of specialty medical organizations are useful in determining the duty owed or the standard of care applicable in a given situation" even though ACR standards themselves do not establish the standard of care.

I. INTRODUCTION

The clinical aspects contained in specific sections of this practice parameter (Introduction, Indications, Specifications of the Examination, and Equipment Specifications) were developed collaboratively by the American College of Radiology (ACR), the American Institute of Ultrasound in Medicine (AIUM), the American College of Obstetricians and Gynecologists (ACOG), the Society for Pediatric Radiology (SPR), and the Society of Radiologists in Ultrasound (SRU). Recommendations for Qualifications and Responsibilities of Personnel, Written Requests for the Examination, Documentation, and Quality Control and Improvement, Safety, Infection Control, and Patient Education vary among the four organizations and are addressed by each separately.

This practice parameter has been developed to assist physicians and other health care providers performing sonographic studies of the female pelvis. Ultrasound of the female pelvis should be performed only when there is a valid medical reason, and the lowest possible ultrasonic exposure settings should be used to gain the necessary diagnostic information. In some cases, additional or specialized examinations may be necessary. Although it is not possible to detect every abnormality, adherence to the following practice parameter will maximize the probability of detecting most abnormalities. For ultrasound of the urinary bladder, see [ACR–AIUM–SPR–SRU Practice Parameter for the Performance of an Ultrasound Examination of the Abdomen and/or Retroperitoneum \[1\]](#).

II. INDICATIONS

Indications for pelvic sonography include, but are not limited to, the following:

1. Evaluation of pelvic pain
2. Evaluation of pelvic masses
3. Evaluation of dyspareunia
4. Evaluation of pregnancy of unknown location or ectopic pregnancy
5. Evaluation of endocrine abnormalities, including polycystic ovaries
6. Evaluation of dysmenorrhea
7. Evaluation of amenorrhea
8. Evaluation of abnormal uterine bleeding (AUB)
9. Evaluation of postmenopausal bleeding
10. Evaluation of delayed menses
11. Follow-up of a previously detected abnormality
12. Evaluation, monitoring, and/or treatment of patients with infertility
13. Evaluation when there is limited clinical examination of the pelvis
14. Evaluation for signs or symptoms of pelvic infection
15. Further characterization of a pelvic abnormality noted on another imaging study
16. Evaluation of congenital uterine, gonadal, and lower genital tract anomalies
17. Evaluation of excessive bleeding, pain, or signs of infection after pelvic surgery, delivery, or abortion
18. Localization of an intrauterine device (IUD)
19. Surveillance for malignancy in high-risk patients
20. Evaluation of incontinence or pelvic organ prolapse
21. Guidance for interventional or surgical procedures
22. Preoperative and postoperative evaluation of pelvic structures

III. QUALIFICATIONS OF PERSONNEL

See the [ACR–SPR–SRU Practice Parameter for the Performance and Interpretation of Diagnostic Ultrasound Examinations \[2\]](#).

IV. SPECIFICATIONS OF THE EXAMINATION

The following section details the examination to be performed for each organ and anatomic region in the female pelvis. All relevant structures should be identified by the transabdominal and/or transvaginal approach. A

transrectal or transperineal approach may be useful in patients who are not candidates for introduction of a vaginal transducer and in assessing the patient with pelvic organ prolapse. More than one approach may be necessary [3,4].

IV. SPECIFICATIONS OF THE EXAMINATION

A. General Pelvic Preparation

For a transabdominal pelvic sonogram a full bladder is typically needed to displace the bowel from the field of view and to provide an optimal acoustic window to better visualize the pelvic structures, particularly if a transvaginal examination cannot be performed. Occasionally, overdistention of the bladder may compromise the evaluation. When this occurs, imaging may be repeated after partial bladder emptying. If any abnormalities of the urinary bladder are detected, these findings should be documented in accordance with the [ACR-AIUM-SPR-SRU Practice Parameter for the Performance of an Ultrasound Examination of the Abdomen and/or Retroperitoneum](#) [1].

For a transvaginal sonogram, the urinary bladder is preferably empty. The patient, the sonographer, or the clinician may introduce the vaginal transducer, preferably under real-time monitoring. Consideration of having a chaperone present should be in accordance with local policy [5,6]. Two-handed technique (one hand on probe and one hand on external lower abdomen) is helpful to assess mobility of structures and may help move bowel away from the adnexa or to stabilize adnexal structures to aid in their visualization. It is recommended that the examiner inform the patient before a dynamic maneuver such as direct manual pressure on the lower abdomen or sliding of the probe is performed.

IV. SPECIFICATIONS OF THE EXAMINATION

B. Uterus

The vagina and uterus provide anatomic landmarks that can be used as reference points for the other pelvic structures, whether normal or abnormal. When examining the uterus, the following should be evaluated: (a) the uterine size, shape, and orientation; (b) the endometrium; (c) the myometrium; and (d) the cervix. In children and adolescents, note should be made whether the uterine configuration is prepubertal or postpubertal. The vagina may be imaged while introducing the transducer and can be a landmark for the cervix [7,8]. Although not part of the standard examination, if evaluation of the vaginal mucosa and rectovaginal septum are desired, instillation of 20 mL of gel into the vagina with distension of the vaginal fornices may be helpful [9].

Overall uterine length is evaluated in sagittal view from the fundus to the cervix (to the external os, if it can be identified). The length can be measured as a straight line from the fundus to the external os using outer-to-outer technique or by measuring from the fundal region along the endometrial lining and endocervical canal (trace method) using outer-to-outer technique [10]. The depth of the uterus (anteroposterior dimension) is measured in the same sagittal view from its anterior to posterior walls, perpendicular to the longitudinal axis of the endometrium. The maximum width is measured in the transverse or coronal view. If volume measurements of the uterine corpus are performed, the cervical component should be excluded from the uterine length measurement. Note is to be made that a volume measurement of the corpus in the nonpregnant state is an estimate because there is no sonographic anatomic landmark for where the cervix ends and the uterine corpus begins.

Abnormalities of the uterus should be documented [11-13]. The myometrium and cervix should be evaluated for contour changes, echogenicity, masses, and cysts as well as symmetry between anterior and posterior myometrium. The myometrial echogenicity is reported as either homogenous or heterogeneous. If the myometrium is heterogeneous due to shadowing or hyperechogenic islands, that should be specified. Myometrial lesions need to be assessed and described. When an abnormality of the myometrium is noted, the objective finding leading to this conclusion must be documented. For example, simply documenting adenomyosis as a subjective finding is insufficient and has poor reproducibility and reliability [14,15]. Note should be made when the uterus is not mobile or tenderness is elicited during the examination. Fixed retroflexion of the uterus, particularly in the presence of posterior adenomyosis, or absence of sliding between the uterus and adjacent rectum or

adnexa, should be recognized as a possible indicator of pelvic adhesion seen in deep endometriosis (DE) in the posterior cul-de-sac [16]. Size and location of clinically relevant lesions should be documented. Masses that may require follow-up or intervention should be measured in at least two dimensions, acknowledging that it is not usually necessary to measure all uterine fibroids.

The endometrium should be evaluated for thickness, focal abnormality, echogenicity and echotexture (homogeneous vs heterogeneous), and the presence and characteristics of fluid or masses in the uterine cavity. The thickest part of the endometrium should be measured perpendicular to its longitudinal plane in the anteroposterior diameter from echogenic to echogenic border, using outer-to-outer technique [10] (see Figure 1). When fluid is present in the cavity, the endometrial thickness should be measured on either side of the fluid in the same plane and the measurements added together to report the total thickness; the fluid should be excluded in this measurement (see Figure 2). In reproductive-aged postmenarchal patients, assessment of the endometrium should allow for variations expected with phases of the menstrual cycle and with hormonal supplementation [13,17,18]. It must be reported if the endometrium is not adequately seen in its entirety or is ill defined; in this circumstance, measurement may not be included in the report. The endometrium should be evaluated using power Doppler to assess the vascular pattern. The location of any areas of focal hypervascularity or feeding vessels should be documented. Sonohysterography may be a useful to further evaluate the patient with AUB, an abnormal appearing endometrium, and to further evaluate or an incompletely visualized endometrium. (See the [ACR-ACOG-AIUM-SRU Practice Parameter for the Performance of Sonohysterography](#) [19]). If the patient has an IUD, its location should be documented.



Figure 1. Measurement of endometrial thickness.

The endometrial thickness measured in its thickest portion from echogenic to echogenic border (calipers) perpendicular to the midline longitudinal plane of the uterus.



Figure 2. Measurement of endometrium with fluid in cavity.

In the presence of endometrial fluid, the measurement of the two separate layers of the endometrium (calipers), excluding the fluid, are added to determine the endometrial thickness.

The addition of 3-D ultrasound (transabdominal, transvaginal, transperineal, and/or transrectal) can be helpful in many circumstances, including, but not limited to, evaluating the relationship of masses to the endometrial cavity, identifying uterine congenital anomalies and thickened and/or heterogenous endometrium, identifying uterine synechia, and evaluating the location and orientation of an IUD and the integrity of the pelvic floor [16,20-27]. Performing 3-D ultrasound of the uterus may be enhanced if done in the luteal phase because the endometrium is hyperechoic and enables visualization of uterine cavity abnormalities.

IV. SPECIFICATIONS OF THE EXAMINATION

C. Adnexae Including Ovaries and Fallopian Tubes

When evaluating the adnexa, an attempt should be made to identify the ovaries first because they can serve as a major point of reference for assessing the presence of adnexal pathology. Ovarian size may be determined by measuring the ovary in three dimensions (longitudinal, transverse, and anteroposterior diameters) on views obtained in two orthogonal planes [28,29] with calculation of ovarian volume as necessary. Any ovarian abnormalities should be documented [30-35].

The ovaries may not be identifiable in some individuals. This issue occurs most frequently before puberty and after menopause when the ovaries are smaller and/or follicles are not consistently present to serve as a landmark [36]. The adnexal region should be surveyed for abnormalities, particularly masses and dilated tubular structures.

If an adnexal abnormality is noted, its relationship to the ovaries and uterus should be assessed. The size and sonographic characteristics of adnexal masses should be documented. Any ovarian lesion should be fully documented with both gray scale and color; gray scale alone cannot determine solid versus debris containing cystic mass. The addition of 3-D to 2-D ultrasound can be helpful to differentiate multilocular ovarian cysts from hydrosalpinges. The use of the sliding organ sign technique can demonstrate the presence or absence of mobility

of the adnexal structures [37,38]. Abnormal ovarian location, such as in the posterior cul-de-sac with adhesion, particularly to the uterus, pelvic side wall, or contralateral ovary, should be documented because this may indicate endometriosis, other sources of adhesions, or displacement of the ovary in the setting of adnexal torsion (ovarian torsion, isolated tubal torsion, or both ovarian and tubal torsion). Asymmetrical enlargement of the ovary and peripheral location of the follicles are suggestive of ovarian torsion. Identifying the twisted vessel ("whirlpool" sign) is also helpful in making this diagnosis [39]. The presence of Doppler signal does not exclude ovarian torsion.

All ovarian lesions should be documented according to a validated standardized risk stratification system. A lesion is defined as a finding judged to be inconsistent with normal physiologic function. The size of the ovary and the lesion are measured in mm as the largest 3 diameters in 2 perpendicular planes. If using the Ovarian-Adnexal Imaging Reporting Data System (O-RADS), lesions are described as unilocular, unilocular-solid, multilocular, multilocular-solid, or solid. Internal contents of cysts should be described as anechoic, low level internal echoes, ground glass internal echoes, or mixed internal echoes. Papillary projections are solid projections at least 3 mm in height when measured from the cyst wall. Papillary projections and/or the largest solid portion should be measured in 3 planes. Color score is measured subjectively: 1 is no vascular flow, 2 is minimal, 3 is moderate, 4 is highly vascular. If the mass has typical features which suggest a specific diagnosis, such as an endometrioma or teratoma, this information should be provided [35,40,41]. If sonographic characteristics are suggestive of a specific diagnosis, such as hemorrhagic cyst, endometrioma, mature teratoma, hydrosalpinx, peritoneal inclusion cyst or pedunculated fibroid, this information should also be provided [40-44, 54-55].

Spectral, color, and/or power Doppler ultrasound may be useful to evaluate the vascular characteristics of pelvic lesions [45-48].

IV. SPECIFICATIONS OF THE EXAMINATION

D. Cul-de-Sac

The cul-de-sac and bowel posterior to the uterus should be evaluated for the presence of free or loculated fluid, or mass. If a mass is detected, its size, position, shape, sonographic characteristics, and relationship to the ovaries and uterus should be documented. Differentiation of normal loops of bowel from a mass may be difficult if only a transabdominal examination is performed. The rectosigmoid colon wall may be imaged from the posterior vaginal fornix [49]. Special attention to the posterior cul-de-sac should be made in women with pelvic pain, with fixed retroflexion of the uterus, with sonographic evidence of posterior adenomyosis and with known or clinically suspected endometriosis [16,49]. Hypoechoic masses with tapering ends in the rectosigmoid wall may be seen in DE [49,50]. The presence of adhesions in the cul-de-sac may be inferred in the absence of a normal uterine sliding sign [49,51] during dynamic imaging. Any tenderness during the ultrasound is helpful to be documented.

V. DOCUMENTATION

Reporting should be in accordance with the [ACR Practice Parameter for Communication of Diagnostic Imaging Findings](#) [52].

Adequate documentation is essential for high-quality patient care. There should be a permanent record of the ultrasound examination and its interpretation. Cine clips may be useful. Comparison with prior relevant imaging studies should be made and is helpful when available. Images of all appropriate areas, both normal and abnormal, should be recorded. Variations from normal size should generally be accompanied by measurements. Images should be labeled with the patient identification, facility identification, examination date, anatomic landmarks, and image orientation. An official interpretation (final report) of the ultrasound examination should be included in the patient's medical record. Retention of the ultrasound examination images should be consistent both with clinical need and with relevant legal and local health care facility requirements.

VI. EQUIPMENT SPECIFICATIONS

Equipment performance monitoring should be in accordance with the [ACR–AAPM Technical Standard for Diagnostic Medical Physics Performance Monitoring of Real Time Ultrasound Equipment \[53\]](#).

The sonographic examination of the female pelvis should be conducted with a real-time scanner, preferably using sector, curved linear, and/or endocavitary transducers. The transducer should be adjusted to operate at the highest frequency appropriate for clinical circumstances, realizing that there is a trade-off between resolution and beam penetration.

VII. QUALITY CONTROL AND IMPROVEMENT, SAFETY, INFECTION CONTROL, AND PATIENT EDUCATION

All transducers should be cleaned after use. Any transducer in contact with mucosa should be covered by a protective sheath prior to use. Following the examination, the sheath should be disposed of and the transducer cleaned with high-level disinfectant. The method of high-level disinfection may depend on manufacturer’s specifications and infectious disease recommendations.

Policies and procedures related to quality, patient education, infection control, and safety should be developed and implemented in accordance with the ACR Policy on Quality Control and Improvement, Safety, Infection Control, and Patient Education appearing under the heading *Position Statement on QC & Improvement, Safety, Infection Control, and Patient Education* on the ACR website (<https://www.acr.org/Advocacy-and-Economics/ACR-Position-Statements/Quality-Control-and-Improvement>).

ACKNOWLEDGEMENTS

This practice parameter was revised according to the process described under the heading *The Process for Developing ACR Practice Guidelines and Technical Standards* on the ACR website (<https://www.acr.org/Clinical-Resources/Practice-Parameters-and-Technical-Standards>) by the Committee on Practice Parameters – Ultrasound of the ACR Commission on Ultrasound and the Committee on Practice Parameters – Pediatric Radiology of the ACR Commission on Pediatric Radiology, in collaboration with the ACOG, the AIUM, the SPR, and the SRU.

Writing Committee – members represent their societies in the initial and final revision of this practice parameter

ACR

Bohm-Velez, Marcela MD, Chair
Cohen, Harris L MD
Itani, Malak MD
Li, Arleen MD
Melany, Michelle MD
Sohaey, Roya MD

SRU

Andreotti, Rochelle E MD
Khati, Nadia J MD

ACOG

Brewster, Wendy MD, PhD
Skinner, Bethany MD

AIUM

Detti, Laura MD
Groszmann, Yvette MD, MPH
Hochberg, Lauri Silver MD
Puscheck, Elizabeth E MBA, MD, MS

SPR

Estroff, Judy MD
Tutman, Jeffrey MD

Committee on Practice Parameters - Ultrasound

(ACR Committee responsible for sponsoring the draft through the process)

Dahiya, Nirvikar MD, Chair
Bohm-Velez, Marcela MD

Azadi, Javad MD
Burgan, Constantine MD

Carnahan, Molly MD
Gabriel, Helena MD
Itani, Malak MD
Penna, Rupinder MD
Squires, Judy H MD

Fung, Christopher MD
Gunabushanam, Gowthaman MBBS, MD
Melany, Michelle MD
Sohaey, Roya MD

Committee on Practice Parameters – Pediatric Imaging

(ACR Committee responsible for sponsoring the draft through the process)

Levin, Terry L MD, Chair
Amodio, John B MD
Blumfield, Einat MD
Goldman-Yassen, Adam MD
Lala, Shailee V MD
Laufer, Adina MD
Maloney, John A MD
Shah, Summit MD
Vatsky, Seth DO

Alizai, Hamza MD
Betz, Bradford W MD
Collard, Michael MD
Lai, Hollie A MD
Lasiacka, Zofia M MD, PhD
Li, Arleen MD
Noda, Sakura MD
Trout, Andrew T MD

Barth, Richard MD, Chair, Commission on Pediatric Radiology
Larson, David B MBA, MD, Chair, Commission on Quality and Safety
Nicola, Lauren MD, Chair, Commission on Ultrasound

Comments Reconciliation Committee

Gupta, Yasha - CSC, Co-Chair
Amodio, John B MD
Barth, Richard MD
Brewster, Wendy MD, PhD
Crummy, Timothy MD, MHA - CSC
Detti, Laura MD
Gabriel, Helena MD
Hochberg, Lauri Silver MD
Khatri, Nadia J MD
Levin, Terry L MD
Melany, Michelle MD
Puscheck, Elizabeth E MBA, MD, MS
Skinner, Bethany MD
Tutman, Jeffrey MD

Winsor, Kimberly MD - CSC, Chair
Andreotti, Rochelle E MD
Bohm-Velez, Marcela MD
Cohen, Harris L MD
Dahiya, Nirvikar MD
Estroff, Judy MD
Groszmann, Yvette MD, MPH
Itani, Malak MD
Larson, David B MBA, MD
Li, Arleen MD
Nicola, Lauren MD
Schoppe, Kurt MD - CSC
Sohaey, Roya MD

REFERENCES

1. American College of Radiology. ACR–AIUM–SPR–SRU practice parameter for the performance of an ultrasound examination of the abdomen and/or retroperitoneum. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-Abd-Retro.pdf>. Accessed January 8, 2018.
2. American College of Radiology. ACR–SPR–SRU practice parameter for the performance and interpretation of diagnostic ultrasound examinations. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-Perf-Interpret.pdf>. Accessed July 20, 2018.
3. Langer JE, Oliver ER, Lev-Toaff AS, Coleman BG. Imaging of the female pelvis through the life cycle.

Radiographics 2012;32:1575-97.

4. Timor-Tritsch IE, Monteagudo A, Rebarber A, Goldstein SR, Tsybmal T. Transrectal scanning: an alternative when transvaginal scanning is not feasible. *Ultrasound Obstet Gynecol* 2003;21:473-9.
5. Stagno SJ, Forster H, Belinson J. Medical and osteopathic boards' positions on chaperones during gynecologic examinations. *Obstet Gynecol* 1999;94:352-4.
6. Davenport MS, Brimm D, Rubin JM, Kazerooni EA. Patient preferences for chaperone use during transvaginal sonography. *Abdom Radiol (NY)* 2016;41:324-33.
7. Walker DK, Salibian RA, Salibian AD, Belen KM, Palmer SL. Overlooked diseases of the vagina: a directed anatomic-pathologic approach for imaging assessment. *Radiographics* 2011;31:1583-98.
8. Wildenberg JC, Yam BL, Langer JE, Jones LP. US of the Nongravid Cervix with Multimodality Imaging Correlation: Normal Appearance, Pathologic Conditions, and Diagnostic Pitfalls. *Radiographics* 2016;36:596-617.
9. Young SW, Saphier NB, Dahiya N, et al. Sonographic evaluation of deep endometriosis: protocol for a US radiology practice. *Abdom Radiol (NY)* 2016;41:2364-79.
10. Abuhamad A, Minton KK, Benson CB, et al. Obstetric and gynecologic ultrasound curriculum and competency assessment in residency training programs: consensus report. *Am J Obstet Gynecol* 2018;218:29-67.
11. Ascher SM, Imaoka I, Lage JM. Tamoxifen-induced uterine abnormalities: the role of imaging. *Radiology* 2000;214:29-38.
12. Behr SC, Courtier JL, Qayyum A. Imaging of mullerian duct anomalies. *Radiographics* 2012;32:E233-50.
13. Williams PL, Laifer-Narin SL, Ragavendra N. US of abnormal uterine bleeding. *Radiographics* 2003;23:703-18.
14. Van den Bosch T, de Bruijn AM, de Leeuw RA, et al. Sonographic classification and reporting system for diagnosing adenomyosis. *Ultrasound Obstet Gynecol* 2019;53:576-82.
15. Harmsen MJ, Van den Bosch T, de Leeuw RA, et al. Consensus on revised definitions of Morphological Uterus Sonographic Assessment (MUSA) features of adenomyosis: results of modified Delphi procedure. *Ultrasound Obstet Gynecol* 2022;60:118-31.
16. Di Donato N, Bertoldo V, Montanari G, Zannoni L, Caprara G, Seracchioli R. Question mark form of uterus: a simple sonographic sign associated with the presence of adenomyosis. *Ultrasound Obstet Gynecol* 2015;46:126-7.
17. Nalaboff KM, Pellerito JS, Ben-Levi E. Imaging the endometrium: disease and normal variants. *Radiographics* 2001;21:1409-24.
18. American College of Obstetricians and Gynecologists. ACOG Committee Opinion No. 734 Summary: The Role of Transvaginal Ultrasonography in Evaluating the Endometrium of Women With Postmenopausal Bleeding. *Obstetrics & Gynecology* 2018;131:945-46.
19. American College of Radiology. ACR–ACOG–AIUM–SRU practice parameter for the performance of sonohysterography. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-Hysterog.pdf>. Accessed January 8, 2018.
20. Fong K, Kung R, Lytwyn A, et al. Endometrial evaluation with transvaginal US and hysterosonography in asymptomatic postmenopausal women with breast cancer receiving tamoxifen. *Radiology* 2001;220:765-73.
21. Benacerraf BR, Shipp TD, Bromley B. Which patients benefit from a 3D reconstructed coronal view of the uterus added to standard routine 2D pelvic sonography? *AJR Am J Roentgenol* 2008;190:626-9.
22. Abuhamad AZ, Singleton S, Zhao Y, Bocca S. The Z technique: an easy approach to the display of the mid-coronal plane of the uterus in volume sonography. *J Ultrasound Med* 2006;25:607-12.
23. Graupera B, Pascual MA, Hereter L, et al. Accuracy of three-dimensional ultrasound compared with magnetic resonance imaging in diagnosis of Mullerian duct anomalies using ESHRE-ESGE consensus on the classification of congenital anomalies of the female genital tract. *Ultrasound Obstet Gynecol* 2015;46:616-22.
24. Mavrelou D, Naftalin J, Hoo W, Ben-Nagi J, Holland T, Jurkovic D. Preoperative assessment of submucous fibroids by three-dimensional saline contrast sonohysterography. *Ultrasound Obstet Gynecol* 2011;38:350-4.
25. Sakhel K, Benson CB, Platt LD, Goldstein SR, Benacerraf BR. Begin with the basics: role of 3-dimensional sonography as a first-line imaging technique in the cost-effective evaluation of gynecologic pelvic disease. *J*

- Ultrasound Med 2013;32:381-8.
26. Santoro GA, Wiczorek AP, Dietz HP, et al. State of the art: an integrated approach to pelvic floor ultrasonography. *Ultrasound Obstet Gynecol* 2011;37:381-96.
 27. Shipp TD, Bromley B, Benacerraf BR. The width of the uterine cavity is narrower in patients with an embedded intrauterine device (IUD) compared to a normally positioned IUD. *J Ultrasound Med* 2010;29:1453-6.
 28. Cohen HL, Tice HM, Mandel FS. Ovarian volumes measured by US: bigger than we think. *Radiology* 1990;177:189-92.
 29. Bodelon C, Pfeiffer RM, Buys SS, Black A, Sherman ME. Analysis of serial ovarian volume measurements and incidence of ovarian cancer: implications for pathogenesis. *J Natl Cancer Inst* 2014;106.
 30. Brown DL, Zou KH, Tempany CM, et al. Primary versus secondary ovarian malignancy: imaging findings of adnexal masses in the Radiology Diagnostic Oncology Group Study. *Radiology* 2001;219:213-8.
 31. Jarvela IY, Sladkevicius P, Kelly S, Ojha K, Nargund G, Campbell S. Three-dimensional sonographic and power Doppler characterization of ovaries in late follicular phase. *Ultrasound Obstet Gynecol* 2002;20:281-5.
 32. Kinkel K, Hricak H, Lu Y, Tsuda K, Filly RA. US characterization of ovarian masses: a meta-analysis. *Radiology* 2000;217:803-11.
 33. Sato S, Yokoyama Y, Sakamoto T, Futagami M, Saito Y. Usefulness of mass screening for ovarian carcinoma using transvaginal ultrasonography. *Cancer* 2000;89:582-8.
 34. Levine D, Brown DL, Andreotti RF, et al. Management of asymptomatic ovarian and other adnexal cysts imaged at US: Society of Radiologists in Ultrasound Consensus Conference Statement. *Radiology* 2010;256:943-54.
 35. Timmerman D, Valentin L, Bourne TH, et al. Terms, definitions and measurements to describe the sonographic features of adnexal tumors: a consensus opinion from the International Ovarian Tumor Analysis (IOTA) Group. *Ultrasound Obstet Gynecol* 2000;16:500-5.
 36. Langer JE, Oliver ER, Lev-Toaff AS, Coleman BG. Imaging of the Female Pelvis through the Life Cycle. *RadioGraphics* 2012;32:1575-97.
 37. Timor-Tritsch IE, Monteagudo A, Tsymbal T. Three-dimensional ultrasound inversion rendering technique facilitates the diagnosis of hydrosalpinx. *J Clin Ultrasound* 2010;38:372-6.
 38. Piessens S, Edwards A. Sonographic Evaluation for Endometriosis in Routine Pelvic Ultrasound. *J Minim Invasive Gynecol* 2020;27:265-66.
 39. Dawood MT, Naik M, Bharwani N, Sudderuddin SA, Rockall AG, Stewart VR. Adnexal Torsion: Review of Radiologic Appearances. *Radiographics* 2021;41:609-24.
 40. Andreotti RF, Timmerman D, Strachowski LM, et al. O-RADS US Risk Stratification and Management System: A Consensus Guideline from the ACR Ovarian-Adnexal Reporting and Data System Committee. *Radiology* 2020;294:168-85.
 41. Andreotti RF, Timmerman D, Benacerraf BR, et al. Ovarian-Adnexal Reporting Lexicon for Ultrasound: A White Paper of the ACR Ovarian-Adnexal Reporting and Data System Committee. *J Am Coll Radiol* 2018;15:1415-29.
 42. Timmerman D, Testa AC, Bourne T, et al. Simple ultrasound-based rules for the diagnosis of ovarian cancer. *Ultrasound Obstet Gynecol* 2008;31:681-90.
 43. Van Calster B, Van Hoorde K, Valentin L, et al. Evaluating the risk of ovarian cancer before surgery using the ADNEX model to differentiate between benign, borderline, early and advanced stage invasive, and secondary metastatic tumours: prospective multicentre diagnostic study. *Bmj* 2014;349:g5920.
 44. Timmerman D, Van Calster B, Testa A, et al. Predicting the risk of malignancy in adnexal masses based on the Simple Rules from the International Ovarian Tumor Analysis group. *Am J Obstet Gynecol* 2016;214:424-37.
 45. Funt SA, Hann LE. Detection and characterization of adnexal masses. *Radiol Clin North Am* 2002;40:591-608.
 46. Kaakaji Y, Nghiem HV, Nodell C, Winter TC. Sonography of obstetric and gynecologic emergencies: Part II, Gynecologic emergencies. *AJR Am J Roentgenol* 2000;174:651-6.
 47. Laing FC, Brown DL, DiSalvo DN. Gynecologic ultrasound. *Radiol Clin North Am* 2001;39:523-40.
 48. Polat P, Suma S, Kantarcy M, Alper F, Levent A. Color Doppler US in the evaluation of uterine vascular abnormalities. *Radiographics* 2002;22:47-53.
 49. Guerriero S, Condous G, van den Bosch T, et al. Systematic approach to sonographic evaluation of the pelvis

- Revised 2024 (Resolution 21)
- Revised 2004 (Resolution 23)
- Revised 2019 (Resolution 27)
- Revised 2014 (Resolution 23)
- Revised 2009 (Resolution 19)
- Amended 2006 (Resolution 35)
- Revised 2004 (Resolution 23)
- Revised 1999 (Resolution 36)
- Revised 1995 (Resolution 37)
- 1991 (Resolution 10)
- Development Chronology for this Practice Parameter
- Revised 2024 (Resolution 21) updated endometriosis, including terms, definitions and measurements: a consensus opinion from the International Deep Endometriosis Analysis (IDEA) group. *Ultrasound Obstet Gynecol* 2016;48:318-32.
50. Benacerraf BR, Groszmann Y, Hornstein MD, Bromley B. Deep infiltrating endometriosis of the bowel wall: the comet sign. *J Ultrasound Med* 2015;34:537-42.
51. Hudelist G, Fritzer N, Staettner S, et al. Uterine sliding sign: a simple sonographic predictor for presence of deep infiltrating endometriosis of the rectum. *Ultrasound Obstet Gynecol* 2013;41:692-5.
52. American College of Radiology. ACR practice parameter for communication of diagnostic imaging findings. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/CommunicationDiag.pdf>. Accessed January 8, 2018.
53. American College of Radiology. ACR technical standard for diagnostic medical physics performance monitoring of real time ultrasound equipment. Available at: <https://www.acr.org/-/media/ACR/Files/Practice-Parameters/US-Equip.pdf>. Accessed January 8, 2018.
54. International Working Group of AAGL, ESGE, ESHRE and WES, Tomassetti C, Johnson NP, Petrozza J, Abrao MS, Einarsson JI, Horne AW, Lee TT, Missmer S, Vermeulen N. An international terminology for endometriosis, 2021. *Facts Views Vis Obgyn* 2021; 13: 295 – 304.
55. Koninckx PR, Ussia A, Adamyan L, Wattiez A, Donnez J. Deep endometriosis: definition, diagnosis, and treatment. *Fertil Steril* 2012; 98: 564 – 571.

*Practice parameters and technical standards are published annually with an effective date of October 1 in the year in which amended, revised or approved by the ACR Council. For practice parameters and technical standards published before 1999, the effective date was January 1 following the year in which the practice parameter or technical standard was amended, revised, or approved by the ACR Council.

Development Chronology for this Practice Parameter

1991 (Resolution 10)

Revised 1995 (Resolution 37)

Revised 1999 (Resolution 36)

Revised 2004 (Resolution 23)

Amended 2006 (Resolution 35)

Revised 2009 (Resolution 19)

Revised 2014 (Resolution 23)

Revised 2019 (Resolution 27)

Revised 2024 (Resolution 21)